

12.5

TURF WEEDGRASS CONTROL XXXX
(S-1280)
05/22/84 RF 07/02/84 RF
06/11/84 RF 07/11/84 VS
06/21/84 RF


FRONT:

(Scotts(R))
ProTurf(R)
BRAND

TURF
WEEDGRASS CONTROL

- . Preemergent control of crabgrass, goosegrass, and other grassy weeds
- . Control of 7 sprouting broadleaf weeds

Net Weight 44 1/4 lbs (20.07 kg)



LEFT COLUMN

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION. Causes eye irritation. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This product is toxic to fish. Do not apply directly to any body of water. Apply this product only as specified on the label. Do not contaminate water by cleaning of equipment or disposal of wastes.

ACTIVE INGREDIENT:

Pendimethalin, N-(1-Ethylpropyl) 3,4 dimethyl- 2,6-dinitrobenzenamine	1.71%
INERT INGREDIENTS:	98.29%
	Total 100.00%

EPA Reg No 538 AXX
Product of USA

EPA Est 538 OH-1

Licensed under one or more of the following U.S. patents 3,920,742; 4,066,441

ProTurf Division, The O M Scott & Sons Company
Marysville, Ohio 43041
(c)1985. The O M Scott & Sons Company. All rights reserved.

Recommended for use by Professionals

BEST DOCUMENT AVAILABLE

XXXXXX

GUSSETS:

RIGHT GUSSET

XXXXX

(Scotts(R))
ProTurf(R)
BRAND

Turf Weedgrass Control

LEFT GUSSET

Scotts(R)
ProTurf(R)
BRAND

Turf Weedgrass Control

XXXXX

BEST DOCUMENT AVAILABLE.

BACK:

(Scotts(R))
ProTurf(R)
BRAND

TURF WEEDGRASS CONTROL

Directions for use

LEFT COLUMN

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This product provides preemergent control of sprouting and grassy broadleaf weeds.

At the LIGHT RATE

For use on most turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bermudagrass, St Augustinegrass, centipedegrass, bahiagrass and zoysiagrass. NOT for use on bentgrasses.

Applied in the Spring this product provides control of:

Sprouting Grassy Weeds

crabgrass
foxtail
goosegrass*
barnyardgrass
fall panicum

Sprouting Broadleaf Weeds

chickweed
oxalis
prostrate spurge

* A repeat application after 6-8 weeks is necessary at the normal rate

Poa annua control at the LIGHT RATE

Applied in mid-August to mid-October before germination, this product prevents the emergence of Poa annua seedlings. Do not make a repeat application until the following spring.

At the HEAVY RATE

For use on southern grasses including established bermudagrass (non overseeded), St Augustinegrass, centipedegrass, bahiagrass, zoysiagrass, and tall fescue.

BEST DOCUMENT AVAILABLE

4
BACK CONTINUED:

Applied in the Spring this product provides control of:

Sprouting Grassy Weeds

crabgrass
goosegrass
foxtail
barnyardgrass
fall panicum

Sprouting Broadleaf Weeds

chickweed
cudweed
hop clover
henbit
oxalis
eveningprimrose
prostrate spurge

Applied in the Fall this product also controls annual dicot weeds germinating during this time of year, such as henbit and chickweed.

HOW TO USE:

Apply product to dry foliage. Water after application to remove granules from foliage and move product to the surface of the soil. Do not destroy the chemical barrier created at the soil surface by raking or aerifying subsequent to application.

PRECAUTIONS:

Do not use where Poa annua or bentgrass is the desired species.

Application of this product to bermudagrass overseeded with cool season turfgrass species will injure and significantly thin the overseeding.

Delay seeding by 4 months and sprigging by 5 months after application.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in.

Application should not exceed twice yearly.

Do not apply to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water.

STORAGE AND DISPOSAL:

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ProLurf Division, The O M Scott & Sons Company
Marysville, Ohio 43041

BEST DOCUMENT AVAILABLE

BACK CONTINUED:

Suggested Spreader Settings

44 1/4 lbs treats 22,000 ft² (1/2 acre) at LIGHT RATE
44 1/4 lbs treats 11,000 ft² (1/4 acre) at HEAVY RATE

<u>SPREADER</u>	<u>GROUND OR PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>LIGHT RATE</u>	<u>SPREADER SETTINGS</u>	
				<u>HEAVY RATE</u>	<u>HALF RATE with 1/2 SWATH WIDTH LIGHT HEAVY**</u>
Scotts Drop Type Gandy (30") Scott Rotary* (R-7, R-7X) Lely Models (see below)					not recommended not recommended

*Set pattern slide to 1/4 closed position.

**At the half rate setting, apply at a swath width of XXXX ft. for the Scott Rotary and X ft. for the Lely to achieve the overall LIGHT or HEAVY RATE.

Metric: 20.07 kg treats 0.2 ha at the LIGHT RATE
20.07 kg treats 0.1 ha at the HEAVY RATE
Lely models WTR, WRF, HR and 1250. PTO at 450 rpm

For proper distribution calibrate your spreader before application.

XXXXX

BEST DOCUMENT AVAILABLE

4830