


10F 2
pm 2i

REG. # 538-193



**Turf
Weedgrass
Control**

Turf Weedgrass Control

**Turf
Weedgrass
Control**



- preemergent control of crabgrass, goosegrass, and other grassy weeds
- control of 7 sprouting broadleaf weeds

Net Weight 44 1/4 lbs (20.07 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION. Causes eye irritation. Avoid contact with eyes, skin or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water. Runoff from treated areas may be hazardous to aquatic organisms in nearby water areas. Do not contaminate water by cleaning of equipment or disposal of

wastes. In case of spills, either collect for use or dispose of properly.

ACTIVE INGREDIENT

Pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenesulfonamide

INERT INGREDIENTS

1.71%

98.29%

Total 100.00%

EPA Reg No 538-193

EPA Est 538-04-1

Product of USA

*Licensed under one or more of the following U.S. patents 3,920,742; 4,086,441

ProTurf Division, The O M Scott & Sons Company
Marysville, Ohio 43041

© 1985 The O M Scott & Sons Company. All rights reserved.

83212

83212

Recommended For Use By Professionals

83212

**Turf
Weedgrass
Control**

by

**Turf
Weedgrass
Control**

83212



**Turf
Weedgrass
Control**

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This product provides preemergent control of sprouting and grassy broadleaf weeds.

At the LIGHT RATE

For use on most turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bermudagrass, St. Augustinegrass, centipedegrass, bahiagrass, and zoysiagrass. NOT for use on bentgrasses.

Applied in the Spring this product provides control of:

Sprouting Grassy Weeds:
crabgrass, bermudagrass, foxtail, tall panicum, goosegrass*

Sprouting Broadleaf Weeds:
chickweed, oxalis, prostrate spurge

*A repeat application after 6-8 weeks is necessary at this light rate.

Poa annua control at the LIGHT RATE

Applied in mid-August to mid-October before germination, this product prevents the emergence of Poa annua seedlings. Do not make a repeat application until the following spring.

At the HEAVY RATE

For use on southern grasses including established bermudagrass (non overseeded), St. Augustinegrass, centipedegrass, bahiagrass, zoysiagrass, and tall fescue.

Applied in the Spring this product provides control of:

Sprouting Grassy Weeds:
crabgrass, bermudagrass, goosegrass, tall panicum, foxtail

Sprouting Broadleaf Weeds:
chickweed, oxalis, cudweed, eveningprimrose, hop clover, prostrate spurge, henbit

Applied in the Fall this product also controls annual dicot weeds germinating during this time of year, such as henbit and chickweed.

How to use

Apply product to dry foliage. Water after application to remove granules from foliage and move product to the surface of the soil. Do not destroy the chemical barrier created at the soil surface by raking or aerifying subsequent to application.

Precautions

Do not use where Poa annua or bentgrass is the desired species.

Application of this product to bermudagrass overseeded with cool season turfgrass species will injure and significantly thin the overseeding.

Delay seeding by 4 months and sprigging by 5 months after application.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in.

Application should not exceed twice yearly.

Do not apply to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water.

Storage and disposal

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ProTurf Division
The O. M. Scott & Sons Company
Marysville, Ohio 43041

Suggested Spreader Settings
To provide distribution calibrate spreader before application.
44% lbs treats 22,000 ft² (1/2 acre) at LIGHT RATE
44% lbs treats 11,000 ft² (1/4 acre) at HEAVY RATE

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVER (ft) overlap wheels overlap wheels	SPREADER SETTINGS LIGHT RATE HEAVY RATE
Scotts Drop Type Gandy (30)	3 mph	5 feet	5 1/2 27
Scott's Rotaries R-7, R-7X, PT-10, PT-11, R-8 (cone 8)	3 mph	8 feet	8 4 1/2
Lely Models (see below)	3 mph	20 feet	20 10 1/2

*Use discharge hose cover #1188 (sold separately) on the PT-10 and PT-11 models. For the R-72 model use the discharge pattern plate #1189.
Lely models PT-10, 1000, 1200, 1400, 1600, 1800, 2000, 2200, 2400, 2600, 2800, 3000, 3200, 3400, 3600, 3800, 4000, 4200, 4400, 4600, 4800, 5000, 5200, 5400, 5600, 5800, 6000, 6200, 6400, 6600, 6800, 7000, 7200, 7400, 7600, 7800, 8000, 8200, 8400, 8600, 8800, 9000, 9200, 9400, 9600, 9800, 10000.
Lely models PT-10, 1000, 1200, 1400, 1600, 1800, 2000, 2200, 2400, 2600, 2800, 3000, 3200, 3400, 3600, 3800, 4000, 4200, 4400, 4600, 4800, 5000, 5200, 5400, 5600, 5800, 6000, 6200, 6400, 6600, 6800, 7000, 7200, 7400, 7600, 7800, 8000, 8200, 8400, 8600, 8800, 9000, 9200, 9400, 9600, 9800, 10000.