FRONT:

(S-1194) (PENDING EPA APPROVAL) FERTILIZER PLUS FUNGICIDE VIII 83196 10/10/86 JF 11/04/86 JF


(Scotts(R)) ProTurf(R) BRAND

23-3-3 Fertilizer Plus Fungicide VIII

broad spectrum fungicide
 controlled release nitrogen

Net Weight 33 1/2 1bs (15.20 kg)

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS



. .

<u>Hazards to Humans and Domestic Animals</u>: CAUTION, causes eye irritation. Do not get in eyes, on skin, or on clothing. In case of contact immediately wash eyes or skin with plenty of water. Get medic: attention if irritation persists. May be harmful if swallowed or absorbeu through skin. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.


<u>Environmental Hazards</u>: Do not apply directly to water.

٩,

EPA Reg No 538-194

EPA Est 538-OH-1

FRONT CONTINUED:



Recommended for Use by Professional Jurfgrass Managers

١.

83196

23-3-3 Fertilizer Plus Fungicide VIII GUSSETS;

RIGHT GUSSET

83196

((

.

N


(Scotts(R)) ProTurf(R) BRAND

23~3-3 Fertilizer Plus Fungicide VIII 3/6


LEFT GUSSET

(Scotts(R)) ProTurf(R) BRAND

23-3-3 Fertilizer Plus Fungicide VIII 83196



416



(Scotts_(R)) ProTurf_(R) BRAND

((

23-3-3 Fertilizer Plus Fungicide VIII

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For use on <u>subfactorized consisting</u> of bentgrass, bluegrass, <u>Poa annua</u>, bermudagrass, ryegrass, fine fescue or tall fescue or mixtures where these grasses predominate.

Controls these diseases while fertilizing the turfgrass:

Dollar spot
Brown patch
Fusarium (Gerlachia) patch*
Leaf spot/melting out
 (Helminthosporium spp/Drechslera spp)

*Washington, Oregon and California only

A STREET THE THIRE STREET
Allens - Dates
Marcole (Printle) - Julio
THE REAL PROPERTY OF THE PARTY

Apply to moist or dry foliage. To avoid product pickup, lightly irrigate treated areas soon after application.

For prevention or control of dollar spot, leaf spot, melting out, pink patch (red thread), Fusarium patch (in Washington, Oregon and California) and brown patch (light to moderate disease), apply at the NORMAL RATE when disease activity is expected or noted. Repeat at two to three week intervals to prevent disease through the season. Repeat at two to three week intervals to (30-days) use the DOUBLE RAYS.

<u>To control</u> severe leaf spot, melting out, dollar spot, brown patch and pink **Patch: ned thready apply** it the DOUBLE RATE every 10-30 days until the disease is brought under control, then revert back to the preventive schedule.

<u>For suppression of summer patch diseases</u> apply at the DOUBLE RATE every 20 days (beginning in late spring to early summer). Three to five applications may be necessary for seasonlong suppression. Water immediately after application and keep turf from becoming drought stressed by watering (1") weekly. Under extreme hot weather conditions on sandy and clay soils, additional watering may be required.

83196

BEST AVAILABLE COPY

BACK CONTINUED:

To control severe brown patch infestations, use a non-fertilizer fungicide such as ProTurf Systemic Fungicide, ProTurf Fungicide VI, or ProTurf Fungicide 7 to limit the amount of fertilizer applied. This lower apply, for the Suppression of Summer great and Cate great (Fractions of Suppression of Summer great and Cate great (Fractions of Ford). Suppression of Summer great and Cate great (Fractions of Ford), suppression of Summer great and Cate great (Fractions of Ford). Suppression of Summer great and Cate great (Fractions of Ford), suppression of Summer great further infestations by following up with ProTurf fungicide or combination fertilizer plus fungicide products.

Note to User

On short cut bentgrass (less than 3/4 inch) when temperatures are above 80° F, apply only to dry foliage.

Do not expose product to atmospheric moisture. For example, after application fold down and seal top of bag. This is particularly important under humid conditions.

Use only in clean, dry hopper. Do <u>not</u> leave product in hopper after application.

Storage and disposal

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, fcod, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ProTurf Division The O M Scott & Sons Company Marysville, OH 43041

١.

.(

BEST AVAILABLE COPY

BACK CONTINUED:

6/6

Suggested Spreader Settings to provide proper distribution calibrate spreader before application

Do not adjust rates of application in order to meet fertilizer requirements for your soil.

33 1/2 lbs (15.20 kg) treats 22,000 ft² (1/2 acre/2044 m²) at NORMAL RATE 33 1/2 lbs (15.20 kg) treats 11,000 ft² (1/4 acre/1022 m²) at DOUBLE RATE

			SPREADER SETTING	
SPREADER	GROUND OR <u>Pto speed</u>	WIDTH OF <u>COVERAGE</u>	NORMAL <u>RATE</u>	DOUBLE <u>RATE</u>
Scotts Drop Scotts Rotaries*	3 mph	overlap wheels	5	6 1/4
R-7, R-7X	3 mph	7 feet .	Ð	G
R-8 (cone 9) Lely WTR &	3 mph	8 feet	3	L
WFR Models Lely-HR	4.5 mph 450 rpm	16 feet	3 1/2111	4 1/2111
(PTO Model)	04.5 mph	16 feet	3 1/2111	4 1/2III 🔻
Gandy (30")	3 mph	overlap wheels	26	32

*Set pattern slide to 1/4 closed position.

١,

08690

(