

Product Description:

ProTurf
Logo

Stock #83220

Weedgrass Control
60 WDG


Major Features

- o Preemergent control of crabgrass, goosegrass, other grassy weeds and broadleaf weeds

Analysis

Net Weight 10 lbs (4.54 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS



Hazards to Humans and Domestic Animals: CAUTION. May cause eye and skin irritation. Wash thoroughly after handling. Avoid contact with eyes, skin or clothing. FIRST AID: in case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This product is toxic to fish. Do not apply directly to any body of water or wet lands. Apply this product only as specified on the label. Do not apply when weather conditions favor drift from target areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

ACTIVE INGREDIENT:

Pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl-
2,6-dinitrobenzenamine*

	.60%
INERT INGREDIENTS:	.40%
TOTAL	100%

*Licensed under one or more of the following U.S. patents:
3,920,742; 4,066,441

EPA Reg No 538-XXX

EPA Est XXX

Product of USA

Sold By: ProTurf Division The O M Scott & Sons Company Marysville, Ohio 43041

(c) 1985. The O M Scott & Sons Company. All rights reserved.

Recommended For Use By Professionals
83220

Directions:Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This product provides preemergent control of selected sprouting grassy and broadleaf weeds.

At the LIGHT RATE

For use on most established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bermudagrass, St Augustinegrass, centipedegrass, bahiagrass and zoysiagrass. NOI for use on bentgrasses, Poa annua where it is the desired species, or on dichondra. See "Precautions" for use on winter overseeding of bermudagrass.

Applied in the spring at the light rate this product provides control of:

Sprouting Grassy Weeds

crabgrass
foxtail
goosegrass*
barnyardgrass
fall panicum
Poa annua (see below)

* A repeat application after 6-8 weeks is necessary at the light rate

Poa annua control

Applied at the light rate in mid-August to mid-October, this product prevents the emergence of Poa annua seedlings. Do not make a repeat application until the following spring.

At the HEAVY RATE

For use on established southern grasses only. This includes bermudagrass (non overseeded), St Augustinegrass, centipedegrass, bahiagrass, zoysiagrass, and tall fescue.

Applied in the spring at the heavy rate this product provides control of:

Sprouting Grassy Weeds

crabgrass
goosegrass
foxtail
barnyardgrass
fall panicum

Sprouting Broadleaf Weeds

prostrate spurge
oxalis
hop clover
cudweed
evening primrose

Applied in the late summer or early fall before germination, this product prevents the sprouting of chickweed and henbit.

Coverage
By Weight

	1 1/4 pound	2 1/2 pounds	5 pounds	10 pounds
Light Rate	1/2 Acre	1 Acre	2 Acres	4 Acres
Heavy Rate	1/4 Acre	1/2 Acre	1 Acre	2 Acres

Usage Instructions

Slowly add the required amount of product to a partially-filled spray tank while agitating. Fill the remainder of the tank with water to the desired level. Continue agitating the mixture during application. Use 3 to 5 gallons of water for every 1,000 sq. ft. of coverage. Do not let mixture stand in the tank for more than 12 hours.

For best results water after application to move the product off the foliage to the soil surface.

Do not mow the treated area until the product has been watered in.

PRECAUTIONS:

At the LIGHT RATE this product may cause yellowing and thinning of the cool season grasses in winter overseeded turf.

Delay seeding by 4 months and sprigging by 5 months after application.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before applying product.

On severely thinned or damaged turf wait until filling-in and rooting of stolons is complete before applying.

This product may stain. If contact with the spray mixture occurs to objects (sidewalks, driveways, etc.), rinse with water.

Not for use on bentgrass and Poa annua where it is the desired species, or on dichondra.

STORAGE AND DISPOSAL:

STORAGE: Do not store above 120°F. Store in a clean, dry place. Reseal opened container securely.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

1990r