

538-195

PM25

1064

ACCEPTED

NOV 26 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 538-195

PROTURF WEEDGRASS CONTROL 60 WDG
 (BOTTLE LABEL) 86017S
 08/14/85 JF 11/04/85 JF
 08/21/85 JF 11/12/85 JF
 08/28/85 JF

FRONT:

(Scotts(R))
 ProTurf(R)
 BRAND

(Scotts(R))
 ProTurf(R)
 BRAND

(Scotts(R))
 ProTurf(R)
 BRAND

Weedgrass
 Control
 60 WDG

Weedgrass Control
 60 WDG

Weedgrass
 Control
 60 WDG

- o Preemergent control of crabgrass, goosegrass, other grassy weeds and broadleaf weeds

Net Weight 8 lbs (3.63 kg)

KEEP OUT OF REACH OF CHILDREN
 CAUTION
 PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION. May cause eye and skin irritation. Wash thoroughly after handling. Avoid contact with eyes, skin or clothing. FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This product is toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to fish in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label. Do not apply when weather conditions favor drift from target areas.

ACTIVE INGREDIENT:

Pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl-
 2,6-dinitrobenzenamine*

INERT INGREDIENTS: 40%
 Total 100%

*Licensed under one or more of the following U.S. patents:
 3,920,742; 4,066,441.

EPA Reg No 538-195


EPA Est 9779-AR-13

Product of USA

Sold By: ProTurf Division
 The O M Scott & Sons Company
 Marysville, Ohio 43041

(c) 1986. The O M Scott & Sons Company. All rights reserved.

86017S



Recommended For Use By Professional Turfgrass Managers

SIDE PANEL:

Preemergent control of crabgrass, goosegrass, other grassy weeds and broadleaf weeds.

BACK:

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

This product provides preemergent control of selected sprouting grassy and broadleaf weeds.

At the LIGHT RATE

For use on most established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bermudagrass (non-overseeded), St Augustinegrass, centipedegrass, bahiagrass and zoysiagrass. NOT for use on bentgrasses and Poa annua where these are the desired species, or on dichondra. See "Precautions" for use on winter overseeding of bermudagrass.

Applied in the spring at the light rate this product provides control of:

Sprouting Grassy Weeds

crabgrass
foxtail
goosegrass*

barnyardgrass
fall panicum
Poa annua (see below)


Sprouting Broadleaf Weeds

prostrate spurge*

oxalis

* A repeat application after 6-8 weeks is necessary at the Light Rate

Poa annua and chickweed control: Applied at the Light Rate in the late summer to early fall before germination, this product prevents the emergence of Poa annua and chickweed seedlings. Do not make a repeat application until the following spring.



BACK CONTINUED:

At the HEAVY RATE

For use on established southern grasses only. This includes bermudagrass (non overseeded), St Augustinegrass, centipedegrass, bahiagrass, zoysiagrass, and tall fescue.

Applied in the spring at the Heavy Rate this product provides control of:

Sprouting Grassy Weeds

- crabgrass
- gousetgrass
- foxtail
- barnyardgrass
- fall panicum

Sprouting Broadleaf Weeds

- prostrate spurge
- oxalis
- hop clover
- cudweed
- evening primrose

Applied at the Heavy Rate in the late summer to early fall before germination, this product prevents the emergence of Poa annua, chickweed and henbit seedlings.

Coverage By Weight

	<u>LIGHT RATE</u>	<u>HEAVY RATE</u>
1 pound	17,000 sq ft	8,500 sq ft
2 pounds	34,000 sq ft	17,000 sq ft
4 pounds	68,000 sq ft	34,000 sq ft
8 pounds	136,000 sq ft	68,000 sq ft

Usage instructions


Slowly add the required amount of product to a partially-filled spray tank while agitating. Fill the remainder of the tank with water to the desired level. Continue agitating the mixture during application. For best results, use 3 to 5 gallons of water for every 1,000 sq. ft. of coverage. Use strainer screens of 30 mesh or coarser. When the tank mixture is allowed to stand, vigorous agitation should be provided to resuspend the materials before spraying.

For best results water after application to move the product off the foliage to the soil surface.

Do not mow the treated area until the product has been watered in.

Precautions:

This product may cause yellowing and thinning of cool season grasses in winter overseeded turf.



BACK CONTINUED:

Delay seeding by 4 months and sprigging by 5 months after application.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before applying product.

On severely thinned or damaged turf wait until filling-in and rooting of stolons is complete before applying.

This product may stain. If contact with the spray mixture occurs to objects (sidewalks, driveways, etc.) rinse with water.

Not for use on bentgrasses and Poa annua where these are the desired species, or on dichondra.

Storage and Disposal:

STORAGE: Do not store above 120°F. Store in a clean, dry place. Reseal opened container securely.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY

This product has been researched to provide necessary data to support its use as described on the label. However, the user should understand that tests have not been carried out under all possible use conditions, and that Scotts cannot and does not warrant that the product is compatible with all other chemicals, or that user's mechanical equipment is appropriate to maintain the product in suspension. User should conduct its own compatibility testing, and should test its mixes and equipment thoroughly before using the product on customers' property. Accordingly, SCOTTS MAKES NO WARRANTY, EXPRESS OR IMPLIED, CONCERNING THE RESULTS TO BE OBTAINED FROM USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH THESE DIRECTIONS AND THE DIRECTIONS ON THE LABEL. The exclusive remedy of the user or buyer and limit of liability of The O M Scott & Sons Company, or any other seller, for any and all losses, damages, and injuries resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. The buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by an officer of The O M Scott & Sons Company.

ProTurf Division
The O M Scott & Sons Company
Marysville, Ohio 43041

