



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

APR 13 1994

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

Vincent Snyder, Jr.  
O. M. SCOTT & SONS CO.  
14111 Scottslawn Road  
Marysville, OH 430416

SUBJECT: Label Amendment Submissions of 07/08/93 in Response to PR Notice 93-7  
EPA Reg. No.: SEE ATTACHED 538-196


Dear Registrant:

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, successfully **REMOVES THE PRODUCT FROM THE SCOPE** of PR Notice 93-7 one the comments (if any) on the attached sheet are addressed. Other than the alterations (if any) listed on the attached sheet, no further alterations to the labeling are necessary to reflect the requirements of the Worker Protection Standard.

Since your amendment proposes to delete uses directly and/or to delete uses implicitly through the use of exclusionary statements), the Agency will publish in the Federal Register a notice of receipt of your amendment to delete uses, as required by section 6(f)(1) of FIFRA. The Agency will defer action for 90 days from the date of the Federal Register notice to give affected minor users an opportunity to try to persuade you to retain the uses(s). When the comment period has expired, the Agency will approve the amendment unless you have withdrawn it, and stamp the amended labeling "accepted." Your stamped-accepted labeling will then be sent to you.

As described in Supplement A of PR Notice 93-11 (self-verification option), you may sell and distribute product bearing the submitted labeling before you receive a copy of your stamped-accepted labeling. Note that your registration is not amended until the comment period required by section 6(f)(1) of FIFRA has expired and the Agency has approved the amendment. Therefore, you may withdraw your proposed amendment at any time during the comment period and until you receive your stamped-accepted labeling.

Sincerely,

  
James Tompkins, Deputy Chief  
Registration Support Branch  
Registration Division (7505W)

Enclosures



Recycled/Recyclable  
Printed with Soy/Canola ink on paper that  
contains at least 50% recycled fiber

1022

O. M. SCOTT & SONS CO.  
538 07/08/93

Original Submissions

- PROFURF INSECTICIDE III - 538-111
- SCOTTS PROFURF NEW K-O-G WEED CONTROL - 538-112
- PROFURF FUNGICIDE VI - 538-159
- PROFURF FERTILIZER PLUS TURF WEEDGRASS CONTROL - 538-196
- PROFURF FLUID FUNGICIDE II - 538-203
- PROFURF FLUID FUNGICIDE III - 538-216
- FERTILIZER PLUS INSECTICIDE/PREEMERGENT WEED CONTROL - 538-226

RANGE LABELS

30422

FRONT

(S-1206T) - LOW END OF RANGE  
PROTURF FERTILIZER PLUS TURF  
WEEDGRASS CONTROL XXXXX

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

10/03/85 JF 01/31/90 VS  
07/07/93 VS/cr  
01/28/94 VS/cr

APR 13 1994

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

(Scotts(R))  
ProTurf(R)  
BRAND

538-196

High Density  
30-3-9  
Fertilizer Plus  
Turf Weedgrass  
Control

- o preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, and Poa annua
- o provides controlled-release nitrogen feeding

Net Weight 75 3/4 lbs (34.36 kg)

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. In case of spills, either collect for use or dispose of properly.

ACTIVE INGREDIENT:

pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl  
-2,6-dinitrobenzenamine\*

INERT INGREDIENTS:	0.50%
	99.50%
	Total 100.00%

EPA Reg. No. 538-196

EPA Est 538-OH-1

Fertilizer Plus Turf Weedgrass Control

30-3-9

XXX-XXXX

Guaranteed Analysis

Total nitrogen (N)	30%
0.6% ammoniacal nitrogen	
22.9% urea, methylene ureas nitrogen	
6.5% water insoluble nitrogen	
Available phosphoric acid (P <sub>2</sub> O <sub>5</sub> )	3%
Soluble potash (K <sub>2</sub> O)	9%

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA

U.S. Patent No's 3,705,794; 3,989,470

Licensed under one or more of the following U.S. Patents: 3,920,742; 4,066,441

ProTurf Division, The O M Scott & Sons Company

14111 Scottslawn Road

Marysville, Ohio 43041

(c) 1993. The O M Scott & Sons Company. All rights reserved.

Recommended for Use by Professional Turfgrass Managers

83816

30-3-9  
Fertilizer Plus  
Turf Weedgrass  
Control

GUESSETS:

RIGHT GUESSET

83816

(Scotts(R))  
ProTurf(R)

30-3-9  
Fertilizer Plus  
Turf Weedgrass  
Control

LEFT GUESSET

(Scotts(R))  
ProTurf(R)  
BRAND

30-3-9  
Fertilizer Plus  
Turf Weedgrass  
Control

83816

BACK:

60722

30-3-9  
Fertilizer Plus  
Turf Weedgrass Control

(Scotts(R))  
ProTurf(R)  
BRAND

30-3-9  
Fertilizer Plus  
Turf Weedgrass Control

**Directions for use**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for reasearch purposes.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass and Poa annua (except putting greens and tees), bermudagrass, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass, Bentgrass and Poa annua Precautions".

Not for use on bentgrass and Poa annua putting greens and tees, or on dichondra.

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

Grassy Weeds

crabgrass  
foxtail  
goosegrass\*  
Poa annua  
barnyardgrass  
fall panicum

Dicot Weeds

chickweed  
oxalis  
prostrate spurge\*  
knotweed  
cudweed  
hop clover  
eveningprimrose  
henbit  
purslane

To prevent crabgrass, foxtail, barnyardgrass, fall panicum, Poa annua, cudweed, eveningprimrose, chickweed, oxalis, spurge, henbit and knotweed, apply product to dry foliage at the recommended rate range in the spring or fall before they sprout.

\*To prevent goosegrass and heavy infestations of spurge, repeat application at the suggested rate after 6-8 weeks.

To prevent Poa annua, henbit and fall germinating chickweed, apply product to dry foliage at the suggested rate in late summer-early fall before germination. A repeat application after Poa annua treatment should not be made until the following spring.

**BACK CONTINUED:**

Apply product uniformly to dry foliage at the suggested rate with a properly calibrated spreader. For best results, water after application (1/4-1/2 inch) to remove granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying subsequent to application. Allow at least a four month span between any two applications except when repeat treatments are necessary after 6-8 weeks.

Do not mow treated area until product has been watered in and turf foliage has dried.

**Precautions**

Do not use on bentgrass and Poa annua putting greens and tees where it is the desired species, or on dichondra.

Do not apply this product to cool season turfgrasses when temperatures are regularly in th 80's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, scalping, and excessive soil moisture).

**Bermudagrass, 1/2 Inch Height or Taller Bentgrass and Poa annua Precautions**

Do not apply to dormant bentgrass and Poa annua (turf that is not actively growing or has experienced frost).

For dormant bermudagrass, an application can be made if it is maintained at a high cut (1/2 inch or more) on medium-fine textured soils such as clays and clay loams.

Where turf thinning or damage has occurred due to winter injury, wait until greenup, fill-in and good rooting before making an application.

Do not apply to bentgrass or Poa annua after late summer.

Application of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseeding.

**Other Precautions**

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly seeded areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

BACK CONTINUED:

Do not apply more than a cumulative total of 802 lbs. of product/acre per year on cool season grasses and 1203 lbs of product/acre per year on warm season grasses.

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water. If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 MPH) due to its staining potential.

Where turf has been severely thinned due to winter damage, insects, disease, scalping, etc., an application should be withheld until filling-in and rooting of stolons/rhizomes is complete and desired turf density is achieved.



<u>Turfgrass Species</u>	<u>Weeds</u>	<u>Rates of Fertilizer Plus Turf Weedgrass Control</u>	<u>Comments</u>
Cool Season Grasses Kentucky bluegrass Perennial ryegrass Fine fescue Tall fescue	crabgrass foxtail <u>Poa annua</u> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	301-348 lbs./acre as the initial application prior to weed germination.	Apply a repeat application of 248-301 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected.
	goosegrass	301-348 lbs./acre in the spring prior to weed germination. Apply a repeat application after 6-8 weeks at 301-348 lbs./acre	
Bentgrass and established <u>Poa annua</u> (1/2 inch height or taller. Not for use on greens and tees)	crabgrass foxtail <u>Poa annua</u> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed	301 lbs./acre in the spring prior to weed germination.	Apply a repeat application of 248-301 lbs./acre after 6-8 weeks where heavy infestations are expected.
	goosegrass	301 lbs./acre in the spring prior to weed germination with a repeat application of 301 lbs/acre after 6-8 weeks.	

<u>Turfgrass Species</u>	<u>Weeds</u>	<u>Rates of Turf Fertilizer Plus Preemergent Weed Control</u>	<u>Comments</u>
Warm Season Grasses Bermudagrass St. Augustinegrass Bahagrass Zoysiagrass Centipedegrass Tall fescue	crabgrass foxtail <u>Poa annua</u> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	301-401 lbs./acre prior to weed germination.	Apply a repeat application of 248-301 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected.
	goosegrass cudweed hop clover evening-primrose henbit	301-348 lbs./acre prior to weed germination.	Apply a repeat application of 301-348 lbs./acre after 6-8 weeks.
Bermudagrass Greens	crabgrass foxtail <u>Poa annua</u> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	301 lbs./acre prior to weed germination.	Apply a repeat application of 301 lbs./acre after 6-8 weeks is necessary.
	goosegrass cudweed hop clover evening-primrose henbit	301 lbs./acre prior to weed germination with a repeat application of 301 lbs./acre after 6-8 weeks.	

BACK CONTINUED

Storage and disposal

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application. Do not adjust rates in order to meet fertilizer requirements of your soil.

75 3/4 lbs treats 11,000 ft<sup>2</sup> (1/4 acre) at the 301 lbs product/acre rate (equivalent to 1.5 lb active ingredient/acre and 2.0 lb N/1,000 sq ft)

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	RECOMMENDED RATE RANGE			
			Lbs product/acre			
			248	301	348	401
Scotts Drop Type	3 mph	overlap wheels	X	X	X	X
Gandy - 30"	3 mph	overlap wheels	X	X	X	X
Scotts Rotaries*						
R-7, R-7X	3 mph	12 feet	X	X	X	X
R-8, (cone 8)	3 mph	11 feet	X	X	X	X
Lely Models	4.5 mph	30 feet	X	X	X	X

\*Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide 1/4 closed.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm. Metric Coverage: 34.36 kg treats 1022 m<sup>2</sup> at the 301 lbs product/acre rate

**LIMITATION OF LIABILITY**

It must be understood that this product has not been evaluated on all bentgrass varieties and Poa annua under all possible growing conditions. The user should exercise judgement and caution when using this product on a given variety until familiar with the performance under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT ON BENTGRASS OR POA ANNUA WHEN NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES. The exclusive remedy of the user or Buyer, and the limit of liability of The O M Scott & Sons Company or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of The O M Scott & Sons Company.

ProTurf Division  
The O M Scott & Sons Company  
14111 Scottslawn Road  
Marysville, Ohio 43041

FRONT

(S-1205T) - HIGH END OF RANGE  
PROTURF FERTILIZER PLUS TURF  
WEEDGRASS CONTROL XXXXX

10/03/85 JF 01/31/90 VS  
07/07/93 VS/cr  
01/28/94 VS/cr

(Scotts(R))  
ProTurf(R)  
BRAND

---

High Density  
30-3-9  
Fertilizer Plus  
Turf Weedgrass  
Control

---

- o preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, and Poa annua
  - o provides controlled-release nitrogen feeding
- 

Net Weight 21 3/16 lbs (9.60 kg)

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. In case of spills, either collect for use or dispose of properly.

ACTIVE INGREDIENT:

pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl  
-2,6-dinitrobenzenamine\*

1.79%

INERT INGREDIENTS:

98.21%

Total 100.00%

EPA Reg. No. 538-196

EPA Est 538-OH-1

Fertilizer Plus Turf Weedgrass Control

30-3-9

XXX-XXXX

Guaranteed Analysis

Total nitrogen (N) . . . . . 30%

0.6% ammoniacal nitrogen

22.9% urea, methylene ureas nitrogen

6.5% water insoluble nitrogen

Available phosphoric acid (P<sub>2</sub>O<sub>5</sub>) . . . . . 3%

Soluble potash (K<sub>2</sub>O) . . . . . 9%

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA

U.S. Patent No's 3,705,794; 3,989,470

\*Licensed under one or more of the following

U.S. Patents: 3,320,742; 4,066,441

ProTurf Division, The O M Scott & Sons Company

14111 Scottslawn Road

Marysville, Ohio 43041

(c) 1993. The O M Scott & Sons Company. All rights reserved.

Recommended for Use by Professional Turfgrass Managers

83816

30-3-9  
Fertilizer Plus  
Turf Weedgrass  
Control

GUESSETS:

RIGHT GUESSET

83816

(Scotts(R))  
ProTurf(R)

30-3-9  
Fertilizer Plus  
Turf Weedgrass  
Control

LEFT GUESSET

(Scotts(R))  
ProTurf(R)  
BRAND

30-3-9  
Fertilizer Plus  
Turf Weedgrass  
Control

83816

BACK:

10512

30-3-9  
Fertilizer Plus  
Turf Weedgrass Control

(Scotts(R))  
ProTurf(R)  
BRAND

30-3-9  
Fertilizer Plus  
Turf Weedgrass Control

---

### Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for reasearch purposes.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass and Poa annua (except putting greens and tees), bermudagrass, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass, Bentgrass and Poa annua Precautions".

Not for use on bentgrass and Poa annua putting greens and tees, or on dichondra.

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

#### Grassy Weeds

crabgrass  
foxtail  
goosegrass\*  
Poa annua  
barnyardgrass  
fall panicum

#### Dicot Weeds

chickweed  
oxalis  
prostrate spurge\*  
knotweed  
cudweed  
hop clover  
eveningprimrose  
henbit  
purslane

To prevent crabgrass, foxtail, barnyardgrass, fall panicum, Poa annua, cudweed, eveningprimrose, chickweed, oxalis, spurge, henbit and knotweed, apply product to dry foliage at the recommended rate range in the spring or fall before they sprout.

\*To prevent goosegrass and heavy infestations of spurge, repeat application at the suggested rate after 6-8 weeks.

To prevent Poa annua, henbit and fall germinating chickweed, apply product to dry foliage at the suggested rate in late summer-early fall before germination. A repeat application after Poa annua treatment should not be made until the following spring.



**BACK CONTINUED:**

Apply product uniformly to dry foliage at the suggested rate with a properly calibrated spreader. For best results, water after application (1/4-1/2 inch) to remove granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying subsequent to application. Allow at least a four month span between any two applications except when repeat treatments are necessary after 6-8 weeks.

Do not mow treated area until product has been watered in and turf foliage has dried.

**Precautions**

Do not use on bentgrass and Poa annua putting greens and tees where it is the desired species, or on dichondra.

Do not apply this product to cool season turfgrasses when temperatures are regularly in the 80's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, scalping, and excessive soil moisture).

**Bermudagrass, 1/2 Inch Height or Taller Bentgrass and Poa annua Precautions**

Do not apply to dormant bentgrass and Poa annua (turf that is not actively growing or has experienced frost).

For dormant bermudagrass, an application can be made if it is maintained at a high cut (1/2 inch or more) on medium-fine textured soils such as clays and clay loams.

Where turf thinning or damage has occurred due to winter injury, wait until greenup, fill-in and good rooting before making an application.

Do not apply to bentgrass or Poa annua after late summer.

Application of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseeding.

**Other Precautions**

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly seeded areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Do not apply more than a cumulative total of 226 lbs. of product/acre per year on cool season grasses and 339 lbs of product/acre per year on warm season grasses.

BACK CONTINUED:

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water. If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 MPH) due to its staining potential.

Where turf has been severely thinned due to winter damage, insects, disease, scalping, etc., an application should be withheld until filling-in and rooting of stolons/rhizomes is complete and desired turf density is achieved.

<u>Turfgrass Species</u>	<u>Weeds</u>	<u>Rates of Fertilizer Plus Turf Weedgrass Control</u>	<u>Comments</u>
Cool Season Grasses Kentucky bluegrass Perennial ryegrass Fine fescue Tall fescue	crabgrass foxtail <u>Poa annua</u> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	83-96 lbs./acre as the initial application prior to weed germination.	Apply a repeat application of 70-83 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected.
	goosegrass	83-96 lbs./acre in the spring prior to weed germination. Apply a repeat application after 6-8 weeks at 83-96 lbs./acre	
Bentgrass and established <u>Poa annua</u> (1/2 inch height or taller. Not for use on greens and tees)	crabgrass foxtail <u>Poa annua</u> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed	83 lbs./acre in the spring prior to weed germination.	Apply a repeat application of 70-83 lbs./acre after 6-8 weeks where heavy infestations are expected.
	goosegrass	83 lbs./acre in the spring prior to weed germination with a repeat application of 83 lbs/acre after 6-8 weeks.	

<u>Turfgrass Species</u>	<u>Weeds</u>	<u>Rates of Turf Fertilizer Plus Preemergent Weed Control</u>	<u>Comments</u>
Warm Season Grasses Bermudagrass St. Augustinegrass Bahiagrass Zoysiagrass Centipedegrass Tall fescue	crabgrass foxtail <u>Poa annua</u> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	83-133 lbs./acre prior to weed germination.	Apply a repeat application of 70-83 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected.
	goosegrass cudweed hop clover evening-primrose henbit	83-96 lbs./acre prior to weed germination.	Apply a repeat application of 83-96 lbs./acre after 6-8 weeks.
Bermudagrass Greens	crabgrass foxtail <u>Poa annua</u> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	83 lbs./acre prior to weed germination.	Apply a repeat application of 83 lbs./acre after after 6-8 weeks is necessary.
	goosegrass cudweed hop clover evening-primrose henbit	83 lbs./acre prior to weed germination with a repeat application of 83 lbs./acre after 6-8 weeks.	

**Storage and disposal**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

**Suggested Spreader Settings**

To provide proper distribution calibrate spreader before application. Do not adjust rates in order to meet fertilizer requirements of your soil.

21 3/16 lbs treats 11,000 ft<sup>2</sup> (1/4 acre) at the 83 lbs product/acre rate (equivalent to 1.5 lb active ingredient/acre and 0.6 lb N/1,000 sq ft)

<u>SPREADER</u>	<u>GROUND OR PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>RECOMMENDED RATE RANGE</u>			
			<u>Lbs product/acre</u>			
			<u>70</u>	<u>83</u>	<u>96</u>	<u>113</u>
Scotts Drop Type	3 mph	overlap wheels	X	X	X	X
Gandy - 30"	3 mph	overlap wheels	X	X	X	X
Scotts Rotaries*						
R-7, R-7X	3 mph	12 feet	X	X	X	X
R-8, (cone 8)	3 mph	11 feet	X	X	X	X
Lely Models	4.5 mph	30 feet	X	X	X	X

\*Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide 1/4 closed.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.  
Metric Coverage: 9.60 kg treats 1022 m<sup>2</sup> at the 83 lbs product/acre rate

**LIMITATION OF LIABILITY**

It must be understood that this product has not been evaluated on all bentgrass varieties and Poa annua under all possible growing conditions. The user should exercise judgement and caution when using this product on a given variety until familiar with the performance under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT ON BENTGRASS OR POA ANNUA WHEN NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES. The exclusive remedy of the user or Buyer, and the limit of liability of The O M Scott & Sons Company or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of The O M Scott & Sons Company.

ProTurf Division  
14111 Scottslawn Road  
The O M Scott & Sons Company  
Marysville, Ohio 43041