



FEB 12 1996

Vincent Snyder
The Scotts Company
14111 Scottslawn Road
Marysville, OH 43041

Dear Mr. Snyder:

Subject: Label Amendments for:

Southern Weedgrass Control
EPA Registration Number 538-188
Your Application Dated 2/1/96, 1/19/96

Turf Fertilizer Plus Preemergent Weed Control
EPA Registration Number 538-213
Your Application Dated 1/19/96

Proturf Weedgrass Control 60 WDG
EPA Registration Number 538-195
Your Application Dated 1/19/96

Turf Weedgrass Control 8637
EPA Registration Number 538-193
Your Application Dated 1/19/96

Proturf Fertilizer Plus Southern Turf Weedgrass Control
EPA Registration Number 538-206
Your Application Dated 1/19/96

Proturf Fertilizer Plus Turf Weedgrass Control
EPA Registration Number 538-196
Your Application Dated 1/22/96

Fertilizer Plus Preemergent Weed Control
EPA Registration Number 538-214
Your Application Dated 1/22/96

The labeling referred to above submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable subject to the following comment.

1. A Statement of Practical Treatment section is added to each product label. Please see the attached sample format for guidance.


BEST AVAILABLE COPY



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber

Your amended labels have been stamped accepted with comment and the copies are enclosed. You must incorporate the comment above on your labels. The new labels must be on products released for shipment within 1 year or by the next label printing, whichever occurs first.

Sincerely,



Robert J. Taylor
Product Manager 25
Fungicide-Herbicide Branch
Registration Division (H7505C)

Enclosures

BEST AVAILABLE COPY

3 of 19

Figure 3. Sample Format For Unrestricted Use Product Label.

8	<p>PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS CAUTION</p> <p>_____</p> <p>_____</p> <p>_____</p>	<p>P R O D U C T</p> <p>W A R N E</p>	SITE (CROP)
8A			SITE (CROP)
8B	<p>ENVIRONMENTAL HAZARDS</p> <p>_____</p> <p>_____</p>		SITE (CROP)
8C	<p>PHYSICAL OR CHEMICAL HAZARDS</p> <p>_____</p> <p>_____</p>	<p>ACTIVE INGREDIENT: X < 6A 6C</p> <p>INERT INGREDIENTS: X < 6B 6E</p> <p>TOTAL: 100.00 %</p>	SITE (CROP)
9B	<p>DIRECTIONS FOR USE</p> <p>It is a violation of Federal law to use this product in a manner inconsistent with its labeling.</p>	<p>THIS PRODUCT CONTAINS LBS OF PER GALLON < 6B</p>	SITE (CROP)
10A	<p>RE-ENTRY STATEMENT (if Applicable)</p> <p>_____</p> <p>_____</p>	<p>KEEP OUT OF REACH OF CHILDREN < 7A</p>	SITE (CROP)
10B	<p>STORAGE AND DISPOSAL</p> <p>STORAGE _____</p> <p>DISPOSAL _____</p>	<p>CAUTION</p> <p>STATEMENT OF PRACTICAL TREATMENT < 7D</p> <p>IF SWALLOWED _____</p> <p>IF INHALED _____</p> <p>IF ON SKIN _____</p> <p>IF IN EYES _____</p>	SITE (CROP)
10C	<p>SITE (CROP) _____</p> <p>_____</p>	<p>SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS < 7E</p> <p>MFG. BY _____ < 2</p> <p>CITY, STATE _____ < 4</p> <p>EPA REGISTRATION NO. _____ < 3</p> <p>ESTABLISHMENT NO. _____ < 5</p> <p>NET CONTENTS _____ < 3</p>	WARRAN

538-191p
(Scotts®)
ProTurf®
BRAND

High Density
30-3-9
Fertilizer Plus
Turf Weedgrass
Control

- preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, and *Poa annua*
 - provides controlled-release nitrogen feeding
-

Net Weight 75 $\frac{3}{4}$ lbs (34.36 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment washwaters. In case of spills, either collect for use or dispose of properly.

ACTIVE INGREDIENT:

pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl
-2,6-dinitrobenzenamine* 0.50%

INERT INGREDIENTS: 99.50%

Total 100.00%

EPA Reg. No. 538-196

EPA Est 538-OH-1

5 7 19

FRONT CONTINUED

Fertilizer Plus Turf Weedgrass Control 30-3-9

XXX-XXXX

Guaranteed Analysis

Total nitrogen (N).....30%
 0.6% ammoniacal nitrogen
 22.9% urea, methylene ureas nitrogen
 6.5% water insoluble nitrogen
 Available phosphate (P₂O₅).....3%
 Soluble potash (K₂O).....9%

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA

U.S. Patent No's 3,705,794; 3,989,470

*Licensed under one or more of the following U.S. Patents: 3,920,742; 4,066,441

The Scotts Company

14111 Scottslawn Road

Marysville, Ohio 43041

© 1996. The Scotts Company. World rights reserved.

Recommended for Use by Professional Turfgrass Managers

83816 30-3-9
Fertilizer Plus
Turf Weedgrass
Control

GUSSETS:

RIGHT GUSSET

83816	(Scotts®) ProTurf®	30-3-9 Fertilizer Plus Turf Weedgrass Control
-------	-----------------------	--

LEFT GUSSET

(Scotts®) ProTurf® BRAND	30-3-9 Fertilizer Plus Turf Weedgrass Control	83816
--------------------------------	--	-------

BACK:

30-3-9
Fertilizer Plus
Turf Weedgrass Control

(Scotts®)
ProTurf®
BRAND

30-3-9
Fertilizer Plus
Turf Weedgrass
Control

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass and *Poa annua* (except putting greens and tees), bermudagrass, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass, Bentgrass and *Poa annua* Precautions".

Not for use on bentgrass and *Poa annua* putting greens and tees, or on dichondra.

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

Grassy Weeds

- barnyardgrass
- crabgrass
- fall panicum
- foxtail
- goosegrass*
- Poa annua*

Dicot Weeds

- chickweed
- cudweed
- eveningprimrose
- henbit
- hop clover
- knotweed
- oxalis
- prostrate spurge*
- purslane

To prevent crabgrass, foxtail, barnyardgrass, fall panicum, *Poa annua*, cudweed, eveningprimrose, chickweed, oxalis, spurge, henbit and knotweed, apply product to dry foliage at the recommended rate range in the spring or fall before they sprout.

*To prevent goosegrass and heavy infestations of spurge, repeat application at the suggested rate after 6-8 weeks.

To prevent *Poa annua*, henbit and fall germinating chickweed, apply product to dry foliage at the suggested rate in late summer-early fall before germination. A repeat application after *Poa annua* treatment should not be made until the following spring.

Apply product uniformly to dry foliage at the suggested rate with a properly calibrated spreader. For best results, water after application (1/4 - 1/2 inch) to remove granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying subsequent to application. Allow at least a four month span between any two applications except when repeat treatments are necessary after 6-8 weeks.

Do not mow treated area until product has been watered in and turf foliage has dried

BACK CONTINUED:

Precautions

Do not use on bentgrass and *Poa annua* putting greens and tees where it is the desired species, or on dichondra.

Do not apply this product to cool season turfgrasses when temperatures are regularly in the 80's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, scalping, and excessive soil moisture).

Bermudagrass, 1/2 Inch Height or Taller Bentgrass and *Poa annua* Precautions

Do not apply to dormant bentgrass and *Poa annua* (turf that is not actively growing or has experienced frost).

When applied to dormant bermudagrass, green-up may be delayed 2 to 3 weeks in the spring. The delay in green-up may be more extensive on low-cut bermudagrass and on sandy soils.

Where turf thinning or damage has occurred due to winter injury, wait until greenup, fill-in and good rooting before making an application.

Do not apply to bentgrass or *Poa annua* after late summer.

Application of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseeding.

Other Precautions

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly seeded areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Do not apply more than a cumulative total of 601 lbs. of product/acre (3.0 lbs. active ingredient/acre) per year on bentgrass and *Poa annua*; 1203 lbs. of product/acre (6.0 lbs. active ingredient/acre) per year on cool season grasses; and 1804 lbs. of product/acre (9.0 lbs. active ingredient/acre) per year on warm season grasses.

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water. If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 MPH) due to its staining potential.

Where turf has been severely thinned due to winter damage, insects, disease, scalping, etc., an application should be withheld until filling-in and rooting of stolons/rhizomes is complete and desired turf density is achieved.

BACK CONTINUED

<u>Turfgrass Species</u>	<u>Weeds</u>	<u>Rates of Fertilizer Plus Turf Weedgrass Control</u>	<u>Comments</u>
Cool Season Grasses Kentucky bluegrass Perennial ryegrass Fine fescue Tall fescue	crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	301-401 lbs./acre as the initial application prior to weed germination.	Apply a repeat application of 248-301 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected.
	goosegrass	301-401 lbs./acre in the spring prior to weed germination. Apply a repeat application after 6-8 weeks at 301-348 lbs./acre	
Bentgrass and established <i>Poa annua</i> (1/2 inch height or taller. Not for use on greens and tees)	crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed	301 lbs./acre in the spring prior to weed germination.	Apply a repeat application of 248-301 lbs./acre after 6-8 weeks where heavy infestations are expected.
	goosegrass	301 lbs./acre in the spring prior to weed germination with a repeat application of 301 lbs/acre after 6-8 weeks.	

BACK CONTINUED

Turfgrass Species	Weeds	Rates of Turf Fertilizer Plus Preemergent Weed Control	Comments
Warm Season Grasses Bermudagrass St. Augustinegrass Bahiagrass Zoysiagrass Centipedegrass Tall fescue	crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	301-401 lbs./acre prior to weed germination.	Apply a repeat application of 248-301 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected.
	goosegrass cudweed hop clover evening-primrose henbit	301-401 lbs./acre prior to weed germination.	Apply a repeat application of 301-348 lbs./acre after 6-8 weeks.
Bermudagrass Greens	crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	301 lbs./acre prior to weed germination.	Apply a repeat application of 301 lbs./acre after after 6-8 weeks is necessary.
	goosegrass cudweed hop clover evening-primrose henbit	301 lbs./acre prior to weed germination with a repeat application of 301 lbs./acre after 6-8 weeks.	

10 of 19

BACK CONTINUED

Storage and disposal

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application. Do not adjust rates in order to meet fertilizer requirements of your soil.

75 3/4 lbs treats 11,000 ft² (1/4 acre) at the 301 lbs product/acre rate (equivalent to 1.5 lb active ingredient/acre and 2.0 lb N/1,000 sq ft)

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	RECOMMENDED RATE RANGE			
			Lbs product/acre			
			248	301	348	401
Scotts Drop Type	3 mph	overlap wheels	X	X	X	X
Gandy - 30"	3 mph	overlap wheels	X	X	X	X
Scotts Rotaries*						
R-7, R-7X	3 mph	12 feet	X	X	X	X
R-8, (cone 8)	3 mph	11 feet	X	X	X	X
Lely Models	4-5 mph	30 feet	X	X	X	X

*Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide 1/4 closed.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.
Metric Coverage: 34.56 kg treats 1022 m² at the 301 lbs product/acre rate

11 7 19

BACK CONTINUED

LIMITATION OF LIABILITY

It must be understood that this product has not been evaluated on all bentgrass varieties and *Poa annua* under all possible growing conditions. The user should exercise judgement and caution when using this product on a given variety until familiar with the performance under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT ON BENTGRASS OR *POA ANNUA* WHEN NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES. The exclusive remedy of the user or Buyer, and the limit of liability of The Scotts Company or any of its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of The Scotts Company.

The Scotts Company
14111 Scottslawn Road
Marysville, Ohio 43041

12 9 19

FRONT

(S-1205T) - HIGH END OF RANGE
PROTURF FERTILIZER PLUS TURF
WEEDGRASS CONTROL XXXXX

10/03/85 JF 01/31/90 VS 04/06/93 VS/cr
05/07/93 VS/cr 01/22/96 VS/kl

(Scotts®)
ProTurf®
BRAND

High Density
30-3-9
Fertilizer Plus
Turf Weedgrass
Control

- preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, and *Poa annua*
- provides controlled-release nitrogen feeding

Net Weight 21^{3/16} lbs (9.60 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment washwaters. In case of spills, either collect for use or dispose of properly.

ACTIVE INGREDIENT:

pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl
-2,6-dinitrobenzenamine* 1.79%

INERT INGREDIENTS: 98.21%
Total 100.00%

EPA Reg. No. 538-196

EPA Est 538-OH-1

FRONT CONTINUED

Fertilizer Plus Turf Weedgrass Control 30-3-9 XXX-XXXX

Guaranteed Analysis

Total nitrogen (N).....30%
0.6% ammoniacal nitrogen
22.9% urea, methylene ureas nitrogen
6.5% water insoluble nitrogen
Available phosphate (P2O5).....3%
Soluble potash (K2O).....9%

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA
U S. Patent No's 3,705,794; 3,989,470
*Licensed under one or more of the following
U.S. Patents: 3,920,742; 4,066,441

The Scotts Company
14111 Scottslawn Road
Marysville, Ohio 43041
©1996. The Scotts Company. World rights reserved.

Recommended for Use by Professional Turfgrass Managers

83816 30-3-9
Fertilizer Plus
Turf Weedgrass
Control

GUSSETS:

RIGHT GUSSET

83816 (Scotts®) ProTurf® 30-3-9
Fertilizer Plus
Turf Weedgrass
Control

LEFT GUSSET

(Scotts®) ProTurf® BRAND 30-3-9
Fertilizer Plus
Turf Weedgrass
Control 83816

BACK:

30-3-9
Fertilizer Plus
Turf Weedgrass Control

(Scotts®)
ProTurf®
BRAND

30-3-9
Fertilizer Plus
Turf Weedgrass
Control

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass and *Poa annua* (except putting greens and tees), bermudagrass, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass, Bentgrass and *Poa annua* Precautions".

Not for use on bentgrass and *Poa annua* putting greens and tees, or on dichondra.

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

Grassy Weeds

barnyardgrass
crabgrass
fall panicum
foxtail
goosegrass*
Poa annua

Dicot Weeds

chickweed knotweed
cudweed oxalis
eveningprimrose prostrate spurge*
henbit purslane
hop clover

To prevent crabgrass, foxtail, barnyardgrass, fall panicum, *Poa annua*, cudweed, eveningprimrose, chickweed, oxalis, spurge, henbit and knotweed, apply product to dry foliage at the recommended rate range in the spring or fall before they sprout.

*To prevent goosegrass and heavy infestations of spurge, repeat application at the suggested rate after 6-8 weeks.

To prevent *Poa annua*, henbit and fall germinating chickweed, apply product to dry foliage at the suggested rate in late summer-early fall before germination. A repeat application after *Poa annua* treatment should not be made until the following spring.

Apply product uniformly to dry foliage at the suggested rate with a properly calibrated spreader. For best results, water after application (¼- ½ inch) to remove granules from the foliage and to move the product to the soil surface

BACK CONTINUED:

for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying subsequent to application. Allow at least a four month span between any two applications except when repeat treatments are necessary after 6-8 weeks.

Do not mow treated area until product has been watered in and turf foliage has dried.

Precautions

Do not use on bentgrass and *Poa annua* putting greens and tees where it is the desired species, or on dichondra.

Do not apply this product to cool season turfgrasses when temperatures are regularly in the 80's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, scalping, and excessive soil moisture).

Bermudagrass, 1/2 Inch Height or Taller Bentgrass and *Poa annua* Precautions

Do not apply to dormant bentgrass and *Poa annua* (turf that is not actively growing or has experienced frost).

When applied to dormant bermudagrass, green-up may be delayed 2 to 3 weeks in the spring. The delay in green-up may be more extensive on low-cut bermudagrass and on sandy soils.

Where turf thinning or damage has occurred due to winter injury, wait until greenup, fill-in and good rooting before making an application.

Do not apply to bentgrass or *Poa annua* after late summer.

Application of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseeding.

Other Precautions

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly seeded areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Do not apply more than a cumulative total of 169 lbs. of product/acre (3.0 lbs. active ingredient/acre) per year on bentgrass and *Poa annua*; 339 lbs. of product/acre (6.0 lbs. active ingredient/acre) per year on cool season grasses; and 505 lbs. of product/acre (9.0 lbs. active ingredient/acre) per year on warm season grasses.

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water. If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 MPH) due to its staining potential.

Where turf has been severely thinned due to winter damage, insects, disease, scalping, etc., an application should be withheld until filling-in and rooting of stolons/rhizomes is complete and desired turf density is achieved.

BACK CONTINUED

Turfgrass Species	Weeds	Rates of Fertilizer Plus Turf Weedgrass Control	Comments
Cool Season Grasses Kentucky bluegrass Perennial ryegrass Fine fescue Tall fescue	crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	83-113 lbs./acre as the initial application prior to weed germination.	Apply a repeat application of 70-83 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected.
	goosegrass	83-113 lbs./acre in the spring prior to weed germination. Apply a repeat application after 6-8 weeks at 83-96 lbs./acre	
Bentgrass and established <i>Poa annua</i> (1/2 inch height or taller. Not for use on greens and tees)	crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed	83 lbs./acre in the spring prior to weed germination.	Apply a repeat application of 70-83 lbs./acre after 6-8 weeks where heavy infestations are expected.
	goosegrass	83 lbs./acre in the spring prior to weed germination with a repeat application of 83 lbs/acre after 6-8 weeks.	

BACK CONTINUED

<u>Turfgrass Species</u>	<u>Weeds</u>	<u>Rates of Turf Fertilizer Plus Preemergent Weed Control</u>	<u>Comments</u>
Warm Season Grasses Bermudagrass St. Augustinegrass Bahia grass Zoysiagrass Centipedegrass Tall fescue	crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	83-113 lbs./acre prior to weed germination.	Apply a repeat application of 70-83 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected.
	goosegrass cudweed hop clover evening- primrose henbit	83-113 lbs./acre prior to weed germination.	Apply a repeat application of 83-96 lbs./acre after 6-8 weeks.
Bermudagrass Greens	crabgrass foxtail <i>Poa annua</i> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	83 lbs./acre prior to weed germination.	Apply a repeat application of 83 lbs./acre after after 6-8 weeks is necessary.
	goosegrass cudweed hop clover evening- primrose henbit	83 lbs./acre prior to weed germination with a repeat application of 83 lbs./acre after 6-8 weeks.	

BACK CONTINUED

Storage and disposal

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application. Do not adjust rates in order to meet fertilizer requirements of your soil.

21 3/16 lbs treats 11,000 ft² (1/4 acre) at the 83 lbs product/acre rate (equivalent to 1.5 lb active ingredient/acre and 0.6 lb N/1,000 sq ft)

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	RECOMMENDED RATE RANGE			
			Lbs product/acre			
			70	83	96	113
Scotts Drop Type	3 mph	overlap wheels	X	X	X	X
Gandy - 30"	3 mph	overlap wheels	X	X	X	X
Scotts Rotaries*						
R-7, R-7X	3 mph	12 feet	X	X	X	X
R-8. (cone 8)	3 mph	11 feet	X	X	X	X
Lely Models	4.5 mph	30 feet	X	X	X	X

*Use discharge hole cover #13030 (already attached to the PT-10 and PT-II models). For the R-7X model use with adjustable pattern slide 1/4 closed

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.
Metric Coverage: 9.60 kg treats 1022 m² at the 83 lbs product/acre rate

BACK CONTINUED

LIMITATION OF LIABILITY

It must be understood that this product has not been evaluated on all bentgrass varieties and *Poa annua* under all possible growing conditions. The user should exercise judgement and caution when using this product on a given variety until familiar with the performance under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT ON BENTGRASS OR *POA ANNUA* WHEN NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES. The exclusive remedy of the user or Buyer, and the limit of liability of The Scotts Company or any of its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of The Scotts Company.

The Scotts Company
14111 Scottslawn Road
Marysville, Ohio 43041