

FLUID FUNGICIDE II #####
03/15/85 MC 05/22/85 JF
04/14/85 JF 01/17/86 JF
04/18/85 JF

FRONT:

(Scotts (R))
ProTurf (R)
BRAND #####

FLUID FUNGICIDE II

o Prevents and controls turfgrass diseases.

Net Contents 1/2 gallon (1.89 liters)

KEEP OUT OF THE REACH OF CHILDREN
DANGER
PRECAUTIONARY STATEMENTS

ACCEPTED
APR 2 1986
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticides
registered under
EPA Reg. No. 538-203

Hazards to Humans and Domestic Animals: DANGER, corrosive, causes eye damage and skin irritation. Harmful if swallowed, inhaled, or absorbed through the skin. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Avoid breathing vapors or spray mist. ✓ Remove and wash contaminated clothing before reuse. Do not graze treated areas. Do not feed clippings to livestock.

FIRST AID: In case of eye contact, flush with plenty of water for 15 minutes and get medical attention. In case of skin contact, wash thoroughly with soap and water. If swallowed, give water or milk if conscious and get medical attention; do not induce vomiting. If inhaled, move to fresh air, if not breathing, give artificial respiration and get medical attention.

Environmental Hazards: Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply only as specified on this label.

ACTIVE INGREDIENTS:

Triadimefon, 1-(4-Chlorophenoxy) -3,3-dimethyl-1-	
(1H-1,2,4-triazol-1-yl)	
-2-butanone.....	16.00%
Metalaxyl, N-(2,6-Dimethylphenyl)-N-(methoxyacetyl)	
alanine, methyl ester.....	16.00%

INERT INGREDIENTS:	68.00%
Total	100.00%

EPA Reg No 538-XXX
Product of USA

EPA Est

Sold by: ProTurf Division, The O.M. Scott & Sons Company
Marysville, Ohio 43041

(c) 1985. The O M Scott & Sons Company. All rights reserved.

Recommended for Use by Professional Turfgrass Managers

#####

BACK:

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For use on turf consisting of bentgrass, Kentucky bluegrass, annual bluegrass, perennial ryegrass, bermudagrass or mixtures where these grasses predominate.

Prevents and controls the following diseases on turf:

- Pythium blight
- Pythium damping-off
- Brown patch
- Dollar spot (including benzimidazole tolerant strains)

MIXING INSTRUCTIONS

Add ProTurf Fluid Fungicide II to the spray tank during filling, while maintaining some agitation.

Thoroughly rinse container and pour remaining contents into spray tank.

Do not combine ProTurf Fluid Fungicide II in the spray tank with surfactants or other adjuvants.

APPLICATION DIRECTIONS

Apply with standard pressurized spray equipment. Works well with normal spray heads.

Use the rates recommended with 4-7 gallons of water per 1,000 ft² to achieve satisfactory coverage.

To Prevent Disease - Use the NORMAL RATE when weather conditions are favorable for the development of disease. Repeat on a 10 day schedule for Pythium and 7-10 days for brown patch and dollar spot control.

To Control Moderate to Severe Disease Infestations - Use the DOUBLE RATE. Repeat on a 7-14 day schedule until disease is brought under control, then treat at the NORMAL RATE on a preventive schedule.

For Newly Seeded Areas -

Use the NORMAL RATE immediately after seeding. Irrigate with 1/4 inch of water. Repeat at 5 - 7 day intervals if conditions remain favorable for disease. Under severe disease conditions, apply the DOUBLE RATE and repeat in 10-14 days.

BACK CONTINUED:

Coverage Chart for the Labeled Rates
of ProTurf Fluid Fungicide II

Rate (fl. oz./M)	1 Pint (16 fl oz) (473 ml) <u>Will cover 1/</u>	1 Quart (32 fl oz) (946 ml) <u>Will cover 1/</u>	1/2 Gallon (64 fl oz) (1892 ml) <u>Will cover 1/</u>
---------------------	--	---	---

NORMAL

(0.7 fl. oz.)	22,500 ft ²	45,000 ft ²	90,000 ft ²
---------------	------------------------	------------------------	------------------------

DOUBLE

(1.4 fl. oz.)	11,250 ft ²	22,500 ft ²	45,000 ft ²
---------------	------------------------	------------------------	------------------------

1/Use with 4 - 7 gallons of water per 1,000 ft². Use of lower rates of water may result in turf injury, especially under hot and humid conditions.

LIMITATIONS: Not a control for turfgrass diseases such as leaf spot or red leaf spot.

ProTurf Fluid Fungicide II contains metalaxyl, a systemic fungicide that has a specific mode of action for the control of Pythium. Metalaxyl could be subject to the development of resistant strains of Pythium. Development of resistance cannot be predicted. Consult with your ProTurf Tech Rep or state agricultural experiment station or extension service specialist for guidance and ways to control any metalaxyl-resistant strain of Pythium which may occur. Consideration should be given to use of other suitable ProTurf fungicides.

NOTE TO USER

Do not allow spray mixture to stand in the tank for longer than 12 hours.

Spray mixture should be agitated continuously during application.

Use a properly calibrated sprayer.

For best results, do not water or mow treated areas within 24 hours of application.

Except in cases of severe infestation, allow one week between applications of this and other control products.

STORAGE AND DISPOSAL

STORAGE: Store in a cool, clean, dry place. Do not store at temperatures exceeding 120° F nor at temperatures below 32° F.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

BACK CONTINUED:

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke. Do not reuse empty container.

ProTurf Division
The O. M. Scott & Sons Company
Marysville, Ohio 43041

06490

