


538-204

01/16/1987

1/2



ACCEPTED

JAN 16 1987

538-204

Scott's
Lawn Insect Control
plus Fertilizer
 New Western



New Western
Lawn Insect Control plus Fertilizer

New Western
Lawn Insect Control
plus Fertilizer
Scott's

For use on grass or dichondra lawns

Controls sod webworms, European crane fly larvae (leatherjackets), chinch bugs, flea beetles and vegetable weevils (in dichondra), white grubs- and other insects listed on other side.

Feeds your lawn with Scott's exclusive slow-release nitrogen fertilizer.

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS.



NET WEIGHT 8 1/6 LBS (3.65 kg)

28-6-4

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals: CAUTION: Harmful if swallowed. May be absorbed through the skin. Do not get in eyes, on skin, or on clothing. Avoid inhalation. **First Aid:** remove contaminated clothing and wash affected areas with soap and water. **Eyes:** Flush eyes with plenty of water. Call a physician if irritation persists. Wash thoroughly after handling and before eating or drinking. Do not consume any food or foodstuffs. Wash treated area after application and allow to dry before permitting use.

How to Pesticide: Diazinon is a cholinesterase inhibitor. Atropine is antidotal if symptoms of cholinesterase inhibition are present.

Environmental Hazards: This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

STORAGE: Store in a clean, dry place. Retain original bag by folding top down and securing.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

ACTIVE INGREDIENT

Diazinon (0,0-Dimethyl 0-2-isopropyl-0-methyl-0-pyridinyl) phosphorothioate 1.00%
 INERT INGREDIENTS 99.00%
 Total 100.00%

EPA Reg No 438-704 EPA Est 338-OH-1
 Product of USA
 New Western Lawn Insect Control plus Fertilizer 28-6-4 641-1121

Guaranteed Analysis

Total Nitrogen (N)	28%
(15% ammoniacal nitrogen)	
(11.5% urea, methylene urea, nitrogen)	
0.5% water insoluble nitrogen	
Available phosphoric acid (P ₂ O ₅)	6%
Soluble potash (K ₂ O)	4%
Derived from monoammonium phosphate urea methylene urea and potassium sulfate	
U.S. Pat No 1,705,764	Other Pat Pend

The O.M. Scott & Sons Company
 Marysville, Ohio 43041

35584

35584

Easy Open



Treats 2,500 sq ft (232 m²)



Scotts Lawn Insect Control plus Fertilizer

Controls insects and feeds the lawn.
Controls sod webworms (lawn moth larvae), European crane fly larvae (leatherjackets), chinch bugs, flea beetles and vegetable weevils (in dichondra).

Also controls the following insects in grass or dichondra lawns:

ants	chiggers	earwigs	white grubs
army worms	elder weevils	leafhoppers	(Japanese beetle, European chafar,
Bermuda mites	crickets	moor crickets	Masked chafar, Phyllophaga-Via,
billbugs	cutworms	ticks	or June beetle)

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO apply when insect activity or damage (chewed or browned leaves) is first observed or suspected in the lawn.

NOTE:

For sod webworm, leatherjackets, army worms, March to mid-April: Apply when insect activity is first observed in the spring; usually mid-March to mid-April.

For chinch bugs: Start treatments when daily high temperatures are 85° or above and repeat at 6 week intervals.

For grub control: The most likely period for grubs in lawns through September but in some cases grubs are active in the spring. An application for spring grub activity requires a repeat summer-early fall application for effective control.

For mole crickets: The best time to apply is in May to July.

DO apply when lawn is dry. For white grubs and mole crickets, water thoroughly, let lawn dry and then apply product.

DO water lawn after application to wash granules from the lawn foliage. If treating for grubs or mole crickets, water thoroughly so that active ingredients will be carried into the rootzone.

DO make a repeat application of a Scotts insect control product (with or without fertilizers) if there are signs of renewed insect activity. (Normal application gives effective control for approximately 4 weeks.)

DO apply at:

Applicator Rate	Scotts Drop Spreader	Scotts EasyGreen™ Rotary
treats 2,500 sq ft	6	26

DO NOT treat just the browned areas of lawn. To prevent further damage, treat entire lawn.

DO NOT use any other lawn control product for at least one week after application.

DO NOT allow traffic on lawn after application until granules are washed off lawn foliage and lawn has dried.

DO NOT use on fruit or vegetable plantings.

DO NOT apply with a rotary-type spreader except Scotts EasyGreen™ Rotary, which is especially designed for safe, even application of control products.

A SCOTTS Lawn Pro PRODUCT

Available only at authorized Lawn Pro stores

Scotts Policy
No-Quibble Guarantee

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

the lawn people
Marysville, Ohio 43041

6/17/86



New Western
Scotts Lawn Insect Control plus Fertilizer

35584