


FRONT:

PROTURF FERTILIZER PLUS
DICOT WEED CONTROL V XXXXX

02/04/92 VS/cr
03/27/92 Vs/cr

(Scotts(R))
ProTurf(R)
BRAND



XX-X-X
Fertilizer Plus
Dicot Weed
Control V

- . provides postemergence dicot weed control
- . provides controlled release nitrogen feeding

Net Weight XX X/X lbs (XX.X kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION. Causes eye injury. Avoid contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. When handling this product, wear chemically resistant gloves. Wash non-disposal gloves thoroughly with soap and water before removing.

Environmental Hazards: Do not apply when weather conditions favor drift from target areas. Apply this product only as specified on the label. Drift or runoff may adversely affect non-target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

This product can reach groundwater from improper handling.

When cleaning equipment, do not pour the washwaters on the ground; spray or drain over a large area from wells and other water sources. If spill occurs, follow disposal instructions on this label.

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic acid	0.680 to	1.220%
Dicamba: 3,6-dichloro-o-anisic acid	0.027 to	0.081%
INERT INGREDIENTS:	99.293 to	98.699%
	Total	100.000%

EPA Reg No 538-209

EPA Est 538-OH-1

ProTurf Fertilizer Plus Dicot Weed Control V

XX-X-X

XXX-XXXX

Guaranteed Analysis

Total nitrogen (N)	XX%
X.X% ammoniacal nitrogen	
XX.X% urea, methylene ureas nitrogen	
XX.X% water insoluble nitrogen	
Available phosphoric acid (P ₂ O ₅)	X%
Soluble potash (K ₂ O)	X%
Derived from monoammonium phosphate, urea, methylene ureas and potassium sulfate.	✓

U S Pat No 3,989,470

Product of USA

ProTurf Division, The O M Scott & Sons Company
 Marysville, Ohio 43041
 (c) 1992. The O M Scott & Sons Company. All rights reserved.

Recommended For Use By Professional Turfgrass Managers

XXXXX

XX-X-X
 Fertilizer Plus
 Dicot Weed
 Control V

GUSSETS:

RIGHT GUSSET

XXXXX

(Scotts(R))
ProTurf(R)
BRAND

XX-X-X
Fertilizer Plus
Dicot Weed
Control V

LEFT GUSSET

Scotts(R)
ProTurf(R)
BRAND

XX-X-X
Fertilizer Plus
Dicot Weed
Control V

XXXXX

BACK:

XX-X-X
Fertilizer Plus
Dicot Weed
Control V

XXXXX

(Scotts(R))
ProTurf(R)
BRAND

XX-X-X
Fertilizer Plus
Dicot Weed
Control V

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on golf course putting greens. Fertilizes grass and controls these rosette and vining type weeds:

aster	English daisy	pigweed
bittercress	eveningprimrose	plantain
black medic	filaree	prickly lettuce
bristly oxtongue	fleabane	purslane
buckhorn	ground ivy	pussytoes
bull thistle	hawkweed	ragweed
burclover	healall	scarlet pimpernel
buttonweed	henbit	shadflower (whitlowort)
carpetweed	horseweed	shepherdspurse
catsear (coast dandelion)	knotweed (seedling)	smartweed
chickweed	lambquarter	sowthistle
chicory	lippia (matchweed)	stitchwort
cinquefoil	mallow	threeseeded mercury
clover (white)	moneywort	thymeleaf sandwort
cudweed	morningglory	vetch
curly dock	mustard	yellow rocket
dandelion	pennywort	yellow toadflax
dichondra	pepperweed	

For use on ornamental Turfgrass areas other than putting greens.

Apply to moist foliage at the RECOMMENDED RATE or for better distribution when using a rotary spreader, apply at the HALF RATE with a reduced width of coverage (see spreader settings). Apply when weeds are growing actively. Grass should not be so tall it touches the bottom of the spreader.

Rain, sprinkling, mowing, or heavy traffic within 24 hours after application can knock granules off weeds and may reduce effectiveness. Apply under non-windy conditions, thereby reducing drift to non-target areas and optimizing particle distribution for good weed control. Retreating may be necessary, but wait at least 3 weeks to observe full results of first application.

BACK CONTINUED:

Precautions

Avoid application when daytime temperatures are consistently 90°F or above.

Do not over apply, especially under the branch spread of trees and shrubs. ✓

On new grass plantings delay application until after the fourth mowing.

Delay planting grass seed until after a heavy rain or watering and not sooner than 1 month after application.

Not for use on creeping bentgrass, St. Augustinegrass, carpetgrass, dichondra, ground covers, garden plants, flowers and shrubs.

Do not apply to centipedegrass turf in the Spring before March 15 (or before the danger of frost has passed), or after September. Do not apply to bahiagrass before March 15 or after November 1.

Storage and disposal

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

OK PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. ✓

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ProTurf Division, The O M Scott & Sons Company
Marysville, Ohio 43041

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application.

This bag treats 22,000 ft² (1/2 acre) at RECOMMENDED RATE.
 This bag treats 22,000 ft² (1/2 acre) at HALF RATE and HALF WIDTH.**

<u>SPREADER</u>	<u>GROUND SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>SPREADER SETTINGS</u>	
			<u>RECOMMENDED RATE</u>	<u>HALF RATE/ HALF WIDTH**</u>
Scotts Drop Type Gandy	3 mph	overlap wheels	X	Not recommended ✓
Scotts Rotaries	3 mph	overlap wheels	X	Not recommended
R-7*, R-7X*, PT-10*, PT-11*	3 mph	7 feet**	X	X
R-8 (cone 8)	3 mph	9 feet	X	X
Lely models (see below)	4.5 mph	18 feet**	X	X

* Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide 1/4 closed.

**At the HALF RATE setting, apply at coverage width of 3 1/2 feet for Scotts rotaries and 9 feet for the Lely rotaries to achieve the overall RECOMMENDED RATE.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.

Metric coverage: 45.4 kg treats 2044m² at the RECOMMENDED RATE.

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY

This product has been researched to provide necessary data to support its uses listed on the label. The user should always follow the label directions and exercise his judgement and caution when using the product under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS, OR IMPLIED, CONCERNING THE RESULTS TO BE OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. The exclusive remedy of the user or Buyer, and the limit of liability of The O M Scott & Sons Company or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The O M Scott & Sons Company.

ProTurf Division
 The O M Scott & Sons Company
 Marysville, Ohio 43041