


PREEMERGENT WEED CONTROL XXXXX

09/11/85 JF 11/14/85 JF
09/19/85 JF 06/04/86 JF
10/02/85 JF 06/10/86 CR
11/01/85 JF 04/04/88 SC
05/23/88 SC 06/14/88 SC
01/30/90 CR



(Scotts(R))
ProTurf(R)
BRAND

30-3-9
Turf Fertilizer
Plus
Preemergent
Weed Control

- preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed and Poa annua
- provides quick green-up plus extended feeding with controlled release SREF nitrogen (slow release encapsulated fertilizer)

Net Weight 75 3/4 lbs (34.36 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In case of spills, either collect for use or dispose of properly.

ACTIVE INGREDIENT:

pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl
-2,6-dinitrobenzenamine*

0.50%

INERT INGREDIENTS:

99.50%

Total 100.00%


EPA Reg. No. 538-213

EPA Est 538-OH-1

Turf Fertilizer Plus Preemergent Weed Control

30-3-9

643-XXXX



Guaranteed Analysis

Total nitrogen (N) 30%
0.6% ammoniacal nitrogen
22.9% urea, methylene ureas nitrogen
6.5% water insoluble nitrogen
Available phosphoric acid (P₂O₅) 3%
Soluble potash (K₂O) 9%

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA

U.S. Patent No's 3,705,794; 3,989,470

*Licensed under one or more of the following U.S. Patents: 3,920,742;
4,066,441

ProTurf Division

The O M Scott & Sons Company

Marysville, Ohio 43041


(c) 1990 The O M Scott & Sons Company. All rights reserved.

Recommended for Use by Professional Turfgrass Managers

XXXXX

30-3-9
Turf Fertilizer
Plus Preemergent
Weed Control

The O M Scott & Sons Company
Marysville, Ohio 43041



RIGHT GUESSET

XOOOX

(Scotts(R))
ProTurf(R)


30-3-9
Turf Fertilizer
Plus Preemergent
Weed Control

LEFT GUESSET

XOOOX

(Scotts(R))
ProTurf(R)

30-3-9
Turf Fertilizer
Plus Preemergent
Weed Control



Directions for use

LEFT COLUMN

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass and Poa annua (except putting greens and tees), bermudagrass, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass, Bentgrass and Poa annua Precautions" and "Limitation of Liability" statements.

Not for use on bentgrass, Poa annua and bermudagrass putting greens and tees, or on dichondra. Where bentgrass, bentgrass-containing turf and Poa annua are the desired species, a light rate application only is recommended.

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

Grassy Weeds

- crabgrass
- foxtail
- goosegrass
- Poa annua
- barnyardgrass
- fall panicum

Dicot Weeds


- chickweed
- oxalis
- knotweed
- prostrate spurge
- purslane
- cudweed
- hop clover
- henbit
- eveningprimrose

To prevent Poa annua and fall germinating chickweed, apply product to dry foliage in late summer/early fall (before germination). Repeat application yearly for continued control. A repeat application made after Poa annua treatment should not be sooner than 4 months.

Where lower infestations are expected, the lighter rates may be used.

Apply product uniformly to dry foliage at the suggested rate with a properly calibrated spreader. For best results, water after application (1/4-1/2 inch) to remove granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying as a segment to application. Allow at least a four month span between the applications except when repeat applications are necessary after 6-8 weeks.

Do not mow treated area until product has been watered in and turf foliage has dried.



Precautions

Do not use on bentgrass, Poa annua and bermudagrass putting greens and tees, or on dichondra. Where bentgrass and bentgrass-containing turf and Poa annua are the desired species, a light rate application only is recommended.

On cool season turfgrasses avoid applications when temperatures are regularly in the 80's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, scalping, and excessive soil moisture).

RIGHT COLUMN

Bermudagrass, Bentgrass and Poa annua Precautions

Do not apply to dormant bentgrass and Poa annua (turf that is not actively growing or has experienced frost).

For dormant bermudagrass, an application can be made if it is maintained at a high cut (1/2 inch or more) on medium-fine textured soils such as clays and clay loams.

Where turf thinning or damage has occurred due to winter injury of bermudagrass, wait until greenup, fill-in and good rooting before making an application.

Application of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseeding.

Do not apply to bentgrass or Poa annua after late summer.

Other Precautions


Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.


Annually, do not apply more than a cumulative total of 802 lbs. of product/A (4 lbs. a/A) on cool season turfgrasses and 1203 lbs of product/A (6 lbs. a/A) on warm season grasses.

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water.

If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 MPH) due to its staining potential.



<u>Turfgrass Species</u>	<u>Weeds</u>	<u>Rates of Turf Fertilizer Plus Preemergent Weed Control</u>	<u>Comments</u>
o Cool Season Grasses Kentucky bluegrass Perennial ryegrass Fine fescue Tall fescue	crabgrass foxtail <u>Poa annua</u> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	301-348 lbs./acre as the initial application prior to weed germination.	Apply a repeat application of 248-301 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected.
	goosegrass	301-348 lbs./acre in the spring prior to weed germination. Apply a repeat application after 6-8 weeks at 301-348 lbs./acre	
o Bentgrass and established <u>Poa annua</u> (1/2 inch height or taller. Not for use on greens and tees)	crabgrass foxtail <u>Poa annua</u> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	301 lbs./acre in the spring prior to weed germination.	Apply a repeat application of 248-301 lbs./acre after 6-8 weeks where heavy infestations are expected.
	goosegrass	301 lbs./acre in the spring prior to weed germination with a repeat application of 301 lbs/acre after 6-8 weeks.	




<u>Turfgrass Species</u>	<u>Weeds</u>	<u>Rates of Turf Fertilizer Plus Preemergent Weed Control</u>	<u>Comments</u>
o Warm Season Grasses Bermudagrass St. Augustinegrass Bahagrass Zoysiagrass Centipedegrass Tall fescue	crabgrass foxtail <u>Poa annua</u> barnyardgrass fall panicum oxalis prostrate spurge purslane knotweed chickweed	301-401 lbs./acre prior to weed germination.	Apply a repeat application of 248-301 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected.
	goosegrass cudweed hop clover evening-primrose henbit	301-348 lbs./acre prior to weed germination with a repeat application of 301-348 lbs./acre after 6-8 weeks.	

Storage and disposal

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



LIMITATION OF LIABILITY

It must be understood that this product has not been evaluated on all bentgrass varieties and Poa annua under all possible growing conditions. The user should exercise judgment and caution when using this product on a given variety until familiar with the performance under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT ON BENTGRASS AND POA ANNUA IF NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES. The exclusive remedy of the user or Buyer, and the limit of liability of The O M Scott & Sons Company or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of The O M Scott & Sons Company.

The O M Scott & Sons Company
Marysville, Ohio 43041

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application.
Do not adjust rates in order to meet fertilizer requirements of your soil.

75 3/4 lbs treats 11,000 ft² (1/4 acre) at the 301 lbs product/acre rate
(equivalent to 1.5 lb active ingredient/acre and 2.0 lb N/1,000 sq ft)

<u>SPREADER</u>	<u>GROUND OR PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>RECOMMENDED RATE RANGE</u>			
			<u>Lbs product/acre</u>			
			<u>248</u>	<u>301</u>	<u>348</u>	<u>401</u>
Gandy - 30"	3 mph	overlap wheels	X	X	X	X
Scotts Rotaries*						
R-7, R-7X	3 mph	XX feet	X	X	X	X
R-8, R-8A (cone 8)	3 mph	XX feet	X	X	X	X
Lely Models	4.5 mph	XX feet	X	X	X	X
Vicon	4.5 mph	XX feet	X	X	X	X

*Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide 1/4 closed.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.
Metric coverage: 34.36 kg treats 1022 m² at the 301 lbs product/acre rate

