

538-213

02/18/2000

117

Please read instructions on reverse before completing form.

Form Approved OMB No. 2070-0060. Approval expires 05-31-98

EPA	United States Environmental Protection Agency Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number
------------	---	--	-----------------------

Application for Pesticide - Section I

1. Company/Product Number 538-213	2. EPA Product Manager Jim Thompkins	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Turf Fertilizer Plus Preemergent Weed Control	PM# PM-25	
5. Name and Address of Applicant (Include ZIP Code) The Scotts Company 14111 Scottslawn Road Marysville, OH 43041 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(I), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section II

NOTIFICATION

<input type="checkbox"/> Amendment - Explain below. <input type="checkbox"/> Resubmission in response to Agency letter dated XX-XX-XX <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated XX-XX-XX <input type="checkbox"/> "Me Too" Application <input type="checkbox"/> Other - Explain below.
---	---

FEB 18 2000

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notifications per PR Notice 98-10.

Section III

1. Material This Product Will Be Packaged In:			
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Unit Packaging wgt. No. per Container	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No If "Yes" Package wgt. No. per Container	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify)
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container	5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			

Section IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Sheila R. Kendrick	Title Senior Regulatory Specialist	Telephone No. (Include Area Code) (937) 644-7214

Certification


I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.

6. Date Application Received
(Stamped)

2. Signature The Scotts Company	3. Title Senior Regulatory Specialist	RECEIVED (Stamped)
BY:		
4. Typed Name Sheila R. Kendrick	5. Date January 17, 2000	



NOTIFICATION
FEB 18 2000



11-3-11 Turf Fertilizer Plus Preemergent Weed Control

ProTurf
11-3-11
Turf Fertilizer
Plus Preemergent
Weed Control

POLY-S[®]
TECHNOLOGY

- preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed and *Poa annua*
- provides a one-to-one feeding of nitrogen and potassium

Net Weight 50 lb (22.68 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION: causes eye irritation. Avoid contact with eyes, skin, or clothing. May be harmful if swallowed. Avoid inhalation. Do not contaminate food or feedstuffs. Do not graze treated areas. Do not feed offspring to livestock.

First Aid:

Eye Irritation: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting. **Swallowing:** Flush with plenty of water. Call a physician if irritation persists. **Inhalation:** Wash with plenty of soap and water. Get medical attention if irritation persists. **Skin Contact:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: Long-sleeved shirt and long pants, water-proof gloves, shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washing, use detergent and hot water. Keep and wash personal protective equipment separately from other laundry.

How to Store/Dispose of Containers

Use as directed.

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if possible gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: This product is toxic to fish. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In case of spill, either collect for use or dispose of properly.

ACTIVE INGREDIENT:

Pendimethalin	0.98%
OTHER INGREDIENTS:	99.02%
	Total 100.00%

EPA Reg. No. 528-213 EPA Est. No. E38-04-1 (S)
(CAS # 49487-43-1) E378-06-1 (R)

Subscript used in first letter of run code on end of container.

Turf Fertilizer Plus Preemergent Weed Control 11-3-11 P643

Guaranteed Analysis

Total nitrogen (N)	11%
0.6% ammoniacal nitrogen	
10.4% urea nitrogen*	
Available phosphate (P ₂ O ₅)	3%
Soluble potash (K ₂ O)	11%
Sulfur (S)	2%
2% Free sulfur (S)	
Derived from: urea, potassium chloride, polymer-coated urea and ammoniacal phosphate.	
Chlorine (Cl), not more than	11%

*A portion of the urea has been coated to provide a "slow-release" effect.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.scottscompany.com>

This product and/or its use is covered by one or more of the following U.S. patent and foreign counterparts thereof: U.S. Pat. Nos. 5,218,466, 5,300,178 and 5,405,426.

Product of U.S.A.

Distributed by:
Professional Business Group
The Scotts Company
14111 Scottswalk Rd.
Marysville, OH 43041
www.scottscompany.com

©1998, The Scotts Company. World rights reserved.



66498
070199

8633

Recommended For Use By Professional Turfgrass Managers



11-3-11 Turf Fertilizer Plus Preemergent Weed Control

Prevents spreading of the following grassy and broad weeds:

Creary Weeds crabgrass foxtail goosegrass Poa annua bromus fall panicum	Claw Weeds chickweed cress henbit prostrate spurge purslane quackgrass sheep sorrel henbit eveningprimrose
---	---

Directions for Use

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 155. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, communication, notification and emergency procedures. It also contains specific instructions and education concerning the safe use of this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow workers entry into treated areas during the restricted-entry interval (REI) of 14 hours.

REI required for entry only to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, for example, susceptible gloves, shoes, plant tools.

For use on established turfgrasses including bermudagrass, buffalograss, and Poa annua (except putting greens and tees), Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, St. Augustinegrass, bahiagrass, centropogon, cynodactylon, and blends of these grasses. See "Limitation of Liability" statement.

Do not use on bermudagrass, Poa annua and bermudagrass putting greens and tees, or on diamonds. Where bermudagrass, bermudagrass containing turf or Poa annua are the desired species, a high-rate application only is recommended.

How/When to Use

For control of broadleaf weeds and broadleaf weeds such as crabgrass, goosegrass, foxtail, bromus, and fall panicum, Poa annua, chickweed, henbit, purslane, and prostrate spurge, apply in the spring before they appear.

To prevent Poa annua and fall germinating annuals, apply product to dry foliage in late summer-early fall (before germination). Repeat application yearly for continued control. A repeat application made after Poa annua treatment should not be sooner than 4 months.

Where lower infestations are expected, the lighter rates may be used.

Apply product uniformly to dry foliage at the suggested rate with a properly calibrated spreader. For best results, under other applications (1 1/2 - 1/2 inch) to remove the granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or covering as a sign of application. Allow at least 4 hours to pass between the applications except when repeat applications are necessary after 4-6 weeks.

Do not enter treated area until product has been washed to and turf foliage has dried.

WEEDS	HOW TO APPLY	HOW TO APPLY	HOW TO APPLY
Bermudagrass Crabgrass Foxtail Goosegrass Poa annua Bromus Fall panicum	Apply to dry foliage in late summer-early fall (before germination). Repeat application yearly for continued control. A repeat application made after Poa annua treatment should not be sooner than 4 months.	Apply to dry foliage in late summer-early fall (before germination). Repeat application yearly for continued control. A repeat application made after Poa annua treatment should not be sooner than 4 months.	Apply to dry foliage in late summer-early fall (before germination). Repeat application yearly for continued control. A repeat application made after Poa annua treatment should not be sooner than 4 months.
Chickweed Cress Henbit Prostrate spurge Purslane Quackgrass Sheep sorrel Henbit Eveningprimrose	Apply to dry foliage in late summer-early fall (before germination). Repeat application yearly for continued control. A repeat application made after Poa annua treatment should not be sooner than 4 months.	Apply to dry foliage in late summer-early fall (before germination). Repeat application yearly for continued control. A repeat application made after Poa annua treatment should not be sooner than 4 months.	Apply to dry foliage in late summer-early fall (before germination). Repeat application yearly for continued control. A repeat application made after Poa annua treatment should not be sooner than 4 months.
Chickweed Cress Henbit Prostrate spurge Purslane Quackgrass Sheep sorrel Henbit Eveningprimrose	Apply to dry foliage in late summer-early fall (before germination). Repeat application yearly for continued control. A repeat application made after Poa annua treatment should not be sooner than 4 months.	Apply to dry foliage in late summer-early fall (before germination). Repeat application yearly for continued control. A repeat application made after Poa annua treatment should not be sooner than 4 months.	Apply to dry foliage in late summer-early fall (before germination). Repeat application yearly for continued control. A repeat application made after Poa annua treatment should not be sooner than 4 months.

Precautions

Do not use on bermudagrass, Poa annua and bermudagrass putting greens and tees, or on diamonds. Where bermudagrass, bermudagrass-containing turf or Poa annua are the desired species, a high-rate application only is recommended.

Do not apply to dormant bermudagrass and Poa annua turf that is not actively growing or has experienced frost.

When any preemergent herbicide is applied to dormant bermudagrass, green-up may be delayed 2 to 3 weeks in the spring. The delay in green-up may be more extensive in low-cut bermudagrass and on sandy soils.

As with most other preemergent herbicides, application of this product to bermudagrass overseeded with cool season turfgrasses may injure and kill the overseedings.

Do not exceed turfgrasses avoid applications when temperatures are regularly in the 60's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, seeping, and excessive soil moisture).

Do not apply to bermudagrass and Poa annua after late summer.

Avoid application to cement, stone, stoneware or others. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water.

If metallic surfaces (ball course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-15 mph) due to its staining potential.

Properties of All Dinitroaniline (DNA) Products

May cause temporary staining of turf 18-in and stolon-licensae rooting where turf has been thinned.

Requires a waiting period before seeding or sprigging.

Seeding or Sprigging Intervals after Treatment

Only seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to allow any remaining chemical residues.

Treatment Interval after Seeding or Sprigging

Do not reseed treated areas until the grass has been mowed at least 4 times and has dried to before making an application.

Notes

Do not apply more than a cumulative total of 340 lbs. of product/acre (24.0 lbs. active ingredient/acre) per year on bermudagrass and Poa annua, 690 lbs. of product/acre (48.0 lbs. active ingredient/acre) per year on cool season turfgrasses, and 1940 lbs. of product/acre (136.0 lbs. active ingredient/acre) per year on warm season turfgrasses.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Re-seal opened bag by folding top down and sealing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

LIMITATION OF LIABILITY

It must be understood that this product has not been evaluated under all possible growing conditions. The user should exercise judgment and caution when using this product on a given variety used familiar with the performance under the growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EITHER BY IMPLICATION OR OTHERWISE, THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT WHEN NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES. The exclusive remedy of the user or Buyer, and the limit of liability of The Scotts Company or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have assented to the terms of this notice, which may be varied only by agreement in writing signed by a duly authorized representative of The Scotts Company.



Professional Business Group
The Scotts Company
14111 Southtown Rd.
Marysville, OH 43041
www.scottscompany.com

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application.
50 lb (22.68 kg) treats 10,729 sq. ft. at the 203 lb product/acre rate
(equivalent to 1.75 lb active ingredient/acre) and 0.5 lb Nitrogen and Potassium/1,000 sq. ft.)

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	RECOMMENDED RATE RANGE				
			145 lb Product/A (1.25 lb w/A) (0.37 lb N/A)	175 lb Product/A (1.5 lb w/A) (0.44 lb N/A)	203 lb Product/A (1.75 lb w/A) (0.51 lb N/A)	340 lb Product/A (2.0 lb w/A) (0.88 lb N/A)	
Scott's Drop	3 mph	overlap wheels	6 1/2	6 1/2	6	4 1/2	7 1/2
Scott's Rotator							
AccuPro 2000 SH 2000 (1000)	3 mph	11 feet	J 1/2	K	K 1/2	L	N
R-R, R-RA, SR-1 (1000)	3 mph	11 feet	I 1/2	J	J 1/2	K	N
R-TX (1000)	3 mph	12 feet	F	F 1/2	G 1/2	H	J 1/2
Lady Mowlin NTR 400, 400 and 1200 PPE of 1000	4.5 mph	36 feet	4 1/2	5	5 1/2	6 1/2	6 1/2



NOTIFICATION 417

FEB 10 2000



13-3-7 NPK Fertilizer Plus Preemergent Weed Control



13-3-7
NPK
Fertilizer Plus
Preemergent
Weed Control



1299

8613-1

- preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed and *Poa annua*
- provides quick green-up
- controlled release Poly-S* nitrogen provides uniform growth with extended feeding
- * Scott's, slow-release polymer-encapsulated fertilizer

Net Weight 50 lb (22.68 kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

First Aid

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.
If in eyes: Flush with plenty of water. Call a physician if irritation persists.
If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.
If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

Personal Protective Equipment (PPE):

Applicators and other handlers must wear: Long sleeved shirt (and long pants), waterproof gloves, shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash personal protective equipment separately from other laundry.

User Safety Recommendations:

- Users should:
- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 - Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 - Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water, or to areas where surface water is present or to adjacent areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. In case of spill, advise control authority in diagram of property.

ACTIVE INGREDIENT:

Pendimethalin	0.86%
OTHER INGREDIENTS	99.14%
Total 100.00%	
1 Cas #40487-42-1	EPA Est. No. 8378-BN-1 (K)
EPA Reg. No. 536-213	536-011-1 (S)

Subscript listed in first letter of nsn code on end of containers

NPK Fertilizer Plus Preemergent Weed Control 13-3-7 F 643

Guaranteed Analysis

Total nitrogen (N)	13%
0.8% ammoniacal nitrogen	
12.4% urea nitrogen*	
Available phosphate (P ₂ O ₅)	3%
Soluble potash (K ₂ O)	7%
Sulfur (S)	1.0%
1.0% free sulfur (S)	

*Distilled from urea, non-mercaptan polymer, primer and chitosan and polymer-encapsulated sulfur-coated urea

Chlorine, not more than

*A portion of the urea has been coated to provide 3.5% coated slow release nitrogen.

Information regarding the analysis and levels of metals in this product is available on the Internet at <http://www.rip.fda.gov>

U.S. Pat. No. 5,219,465; 5,200,125; 5,405,476 and 5,466,274

Product of U.S.A.

Distributed by:
Professional Business Group
The Scotts Company
14111 Scindden Rd.
Marysville, OH 43041

©1999, The Scotts Company.
All rights reserved.

Recommended For Use By Professional Turfgrass Manager



22-0-6 Turf Fertilizer Plus Preemergent Weed Control

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Not for use on turf being grown for sale or other commercial use as seed, or for commercial seed production, or for research purposes.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass and *Poa annua* (except putting greens and tees), bermudagrass, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass, Bentgrass and *Poa annua* Precautions" and "Limitation of Liability" statements.

Not for use on bentgrass, *Poa annua* and bermudagrass putting greens and tees, or on dishdrains. Where bentgrass, bermudagrass containing turf and *Poa annua* are the desired species, a light rate application only is recommended.

Prevents sprouting of the following grassy and seed weeds and provides a controlled-release nitrogen feeding:

GRASSY WEEDS	SEED WEEDS
crabgrass	chickweed
foxtail	goats
sparganium	henbit
<i>Poa annua</i>	preclude spurge
bermudagrass	purple
leaf panicle	stinkweed
	leaf clover
	henbit
	eveningprimrose

To prevent *Poa annua* and leaf germinating chickweed, apply product to dry foliage in late summer/early fall (before germination).

Repeat application yearly for continued control. A repeat application made after *Poa annua* treatment should not be closer than 6 months.

Where lower infestations are expected, the lighter rates may be used.

Apply product uniformly to dry foliage of the suggested rate with a properly calibrated spreader. For best results, water after application (1/4-1/2 inch) to remove granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or mowing or a treatment to application. Allow at least a four week gap between the applications except when repeat applications are necessary after 8-8 weeks.

Do not mow treated area until product has been watered in and turf foliage has dried.

Precautions

Do not use on bentgrass, *Poa annua* and bermudagrass putting greens and tees, or on dishdrains. Where bentgrass and bermudagrass-containing turf and *Poa annua* are the desired species, a light rate application only is recommended.

On cool season turfgrasses avoid applications when temperatures are regularly in the 60's.

This product may cause desiccation and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40 F, scalping, and excessive salt buildup).

Bermudagrass, Bentgrass, and *Poa annua* Precautions

Do not apply to dormant bermudagrass or *Poa annua* (turf that is not actively growing or has experienced loss).

When applied to dormant bermudagrass, green-up may be delayed 2 to 3 weeks in the spring; the delay in green-up may be more extensive in low-cut bermudagrass on sandy soils.

Where turf thinning or damage has occurred due to winter injury of bermudagrass, wait until green-up, 10-in and good rooting before making an application.

Applications of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseed.

Do not apply to bentgrass or *Poa annua* after late summer.

Other Precautions

Delay seeding by 4 months and springing by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Do not apply more than a cumulative total of 200 lbs. of product/acre (2.0 lbs. active ingredient/acre) per year on bentgrass and *Poa annua* (100 lbs. of product/acre (1.0 lbs. active ingredient/acre) per year on cool season grasses; and 1100 lbs. product/acre (2.0 lbs. active ingredient/acre) per year on warm season turfgrasses.

Avoid application to cement, stone, clothing, or shoes. If contact occurs, rinse off with water to prevent staining. If necessary, rinse with water. If salt-like material (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (3-10 MPH) due to its staining potential.

Grass Species	Rate	Apply to turf	Frequency
Cool Season Grasses	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Kentucky bluegrass	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Perennial ryegrass	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
<i>Poa annua</i>	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Tall fescue	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Bermudagrass	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
St. Augustinegrass	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Bahiagrass	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Zoysiagrass	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Centipedegrass	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Warm Season Grasses	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Bermudagrass	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
St. Augustinegrass	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Bahiagrass	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Zoysiagrass	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Centipedegrass	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Putting Greens and Tees	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Putting Greens	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.
Tees	100-200 lbs./acre	Apply to turf	Apply a repeat application of 100-200 lbs./acre after 8-8 weeks for extended control or where heavy weed infestations are expected.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reclose opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

LIMITATION OF LIABILITY

It must be understood that this product has not been evaluated on all grasses varieties and *Poa annua* under all possible growing conditions. The user should exercise judgment and caution when using this product on a given variety until familiar with the performance under his growing conditions. **NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT ON BENTGRASS AND *POA ANNUA* IF NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES.** The exclusive remedy of the user or buyer, and the limit of liability of The Scotts Company or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of The Scotts Company.

Distributed by:
Professional Business Group
The Scotts Company
14111 Scottsville Rd.
Marysville, OH 43041
www.scottscompany.com

Suggested Spreader Settings
To provide proper distribution calibrate spreader before application.
Do not adjust rates in order to meet fertilizer requirements of your soil.
50 lbs. treats 11,000 sq. ft. (1/4 acre) at the 198 lbs. product/acre rate (4.54 lb. product/1000 sq. ft.)
(equivalent to 1.5 lbs. active ingredient/acre and 1.0 lb. N/1000 sq. ft.)

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	RECOMMENDED RATE RANGE			
			2.0 lbs. product/1000 sq. ft.	4.54 lbs. product/1000 sq. ft.	5.3 lbs. product/1000 sq. ft.	6.1 lbs. product/1000 sq. ft.
Scotts Broadcast	1 mph	overlap 1/4 inch	6	6 1/2	6 1/2	6 1/2
Scotts Broadcast	1 mph	11 ft	L	M 1/2	O	P
H.S. M.A. (11 ft)	1 mph	11 ft	L	M 1/2	O	P 1/2
K.T. # 77	1 mph	11 ft	H	F	F	K
Club	1 mph	overlap 1/4 inch	5	5	5	5
Club Models (11 ft)	1 mph	11 ft	5 1/2	5 1/2	5 1/2	5 1/2