

RESTRICTIONS: DO NOT USE TREATED SEED FOR FOOD,  
FEED, OR OIL PURPOSES.

pg. 30 of 31

**NOTICE -- READ CAREFULLY  
LIMITED WARRANTY STATEMENT**

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, except for the purposes of curative control of existing disease or insect pests, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson, Inc., no claims are made to guarantee germination of carry-over seed.

Gustafson warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label. Gustafson neither makes, nor authorizes any agent or representative to make any other warranty of FITNESS or MERCHANTABILITY, guarantee or representation, expressed or implied, concerning this material. In the event of any breach of this limited warranty, Gustafson shall not be liable for consequential damages.


A quality product manufactured by:  
Gustafson, Inc.  
1400 Preston Road, Suite 400  
Plano, Texas 75093  
(214) 985-8877

Gustafson (logo) is a Reg. TM of Gustafson, Inc.  
Terraclor SuperX is a Reg. TM of Uniroyal Chemical Company,  
Inc.

54(S)-096170

**BEST AVAILABLE COPY**

(S-1206T)  
 30-3-9 FERTILIZER PLUS  
 PRE-EMERGENT WEED CONTROL (RANGE) XXXX  
 09/11/85 JF 11/14/85 JF  
 09/19/85 JF 06/04/86 JF  
 10/02/85 JF 06/10/86 CR  
 11/01/85 JF



FRONT:

Plus  
 538-214  
 P. 10-14

30-3-9  
 Fertilizer  
 Plus  
 Pre-Emergent  
 Weed Control

- o preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed and Poa annua
- o provides quick green-up from soluble nitrogen

Net Weight 75 3/4 lbs (34.36 kg)

KEEP OUT OF REACH OF CHILDREN  
 CAUTION  
 PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In case of spills, either collect for use or dispose of properly.

ACTIVE INGREDIENT:  
 pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine\* . . . . . 0.50%

INERT INGREDIENTS: . . . . . 99.50%

Total 100.00%

EPA Reg. No. 538-214 EPA Est

Fertilizer Plus Pre-Emergent Weed Control 30-3-9 643-XXXX

FRONT CONTINUED:

08/20/14

**Guaranteed Analysis**

Total nitrogen (N) . . . . .	30%
0.6% ammoniacal nitrogen	
22.9% urea, methylene ureas nitrogen	
6.5% water insoluble nitrogen	
Available phosphoric acid (P <sub>2</sub> O <sub>5</sub> ) . . . . .	3%
Soluble potash (K <sub>2</sub> O) . . . . .	9%

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA

U.S. Pat nos. 3,705,794; 3,989,470

\*Licensed under one or more of the following U.S. Patents: 3,920,742; 4,066,441

*Formulated For:* The O M Scott & Sons Company  
Marysville, Ohio 43041

(c) 1986 The O M Scott & Sons Company. All rights reserved.

---

Recommended for Use by Professional Turfgrass Managers

---

XXXXX

30-3-9  
Fertilizer Plus  
Pre-Emergent  
Weed Control

GUSSETS:

Pr 2/14

RIGHT GUSSET

XXXXX

30-3-9  
Fertilizer Plus  
Pre-Emergent  
Weed Control

LEFT GUSSET

30-3-9  
Fertilizer Plus  
Pre-Emergent  
Weed Control

XXXXX

**BACK:**

30-3-9  
Fertilizer Plus  
Pre-Emergent  
Weed Control

30-3-9  
Fertilizer Plus  
Pre-Emergent  
Weed Control

---

**Directions for use**

**LEFT COLUMN**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass and Poa annua (except putting greens and tees), bermudagrass (non-winter overseeded), St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass, Bentgrass and Poa annua Precautions" and "Limitation of Liability" statements.

Not for use on bentgrass and Poa annua putting greens and tees, or on dichondra. Where bentgrass, bentgrass-containing turf and Poa annua are the desired species, a light rate application only is recommended.

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

Grassy Weeds

- crabgrass
- foxtail
- goosegrass
- Poa annua
- barnyardgrass
- fall panicum

Dicot Weeds

- chickweed
- oxalis
- prostrate spurge
- knotweed

To prevent goosegrass, crabgrass, foxtail, barnyardgrass, fall panicum, oxalis, knotweed and prostrate spurge, apply in the spring before they sprout.

To prevent Poa annua and fall germinating chickweed, apply product to dry foliage in late summer/early fall (before germination). Repeat application yearly for continued control.

Northern (Cool Season) Grasses

Apply 5.7 to 8.0 lbs. of product per 1,000 sq. ft. (1.25 to 1.75 lbs. of active ingredient per acre) prior to germination of weeds. See "Suggested Spreader Settings".

BACK CONTINUED:

pg. 50, 141

Southern (Warm Season) Grasses

Apply 6.9 to 9.2 lbs. of product per 1,000 sq. ft. (1.5 to 2.0 lbs. of active ingredient per acre) prior to germination of weeds. See "Suggested Spreader Settings".

Where lower weed infestations are expected, the lighter rates may be used.

For best results, water after all applications (1/4 - 1/2 inch) to remove the granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying after applying.

RIGHT COLUMN

Precautions:

Do not use on bentgrass and Poa annua putting greens and tees, or on dichondra. Where bentgrass and bentgrass-containing turf and Poa annua are the desired species, a light rate application only is recommended.

On cool season turfgrasses avoid applications when temperatures are regularly in the 80's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature, scalping - below 40°F, and excessive soil moisture).

Bermudagrass, Bentgrass and Poa annua Precautions

Do not apply to dormant bermudagrass, bentgrass or Poa annua (turf that is not actively growing or has experienced frost). However, for dormant bermudagrass, an application can be made if it is maintained at a high cut (1/2 inch or more) on medium-fine textured soils such as clays and clay loams.

Where turf thinning or damage has occurred due to winter injury of bermudagrass, wait until greenup, fill-in and good rooting before making an application.

Applications to overseeded bermudagrass may injure and thin the cool season variety. Do not apply to bentgrass or Poa annua after late summer.

BACK CONTINUED:

#### Other Precautions

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Applications should not exceed twice yearly.

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water.

#### Storage and disposal

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#### LIMITATION OF LIABILITY

It must be understood that this product has not been evaluated on all bentgrass varieties and Poa annua under all possible growing conditions. The user should exercise judgment and caution when using this product on a given variety until familiar with the performance under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT ON BENTGRASS AND POA ANNUA IF NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES. The exclusive remedy of the user or Buyer, and the limit of liability of O. M. Scott & Sons Company or any other Seller, for any and all losses, injured or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of O. M. Scott & Sons Company.

The O M Scott & Sons Company  
Marysville, Ohio 43041

**Suggested Spreader Settings**

To provide proper distribution calibrate spreader before application.  
Do not adjust rates in order to meet fertilizer requirements of your soil.

75 3/4 lbs treats 11,000 ft<sup>2</sup> (1/4 acre) at the 6.9 lbs product/1,000 sq ft rate (equivalent to 1.5 lb active ingredient/acre and 2.0 lb N/1,000 sq ft)

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	RECOMMENDED RATE RANGE				
			Lbs. Product/ 1,000 Sq. Ft.	5.7	6.9	8.0	9.2
Scotts Drop Type	3 mph	overlap wheels		X	X	X	X
Gandy - 30"	3 mph	overlap wheels		X	X	X	X
Scotts Rotaries*							
R-7, R-7X	3 mph	XX feet		X	X	X	X
R-8 (cone 8)	3 mph	XX feet		X	X	X	X
Lely Models	4.5 mph	XX feet		X	X	X	X
Vicon	4.5 mph	XX feet		X	X	X	X

\*Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide 1/4 closed.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.  
Metric coverage: 22.67 kg treats 1022 m<sup>2</sup> at the 6.9 lbs. product/1,000 sq. ft. rate

(S-1205T) 98-114

30-3-9 FERTILIZER PLUS  
PRE-EMERGENT WEED CONTROL (RANGE) XXXX  
09/11/85 JF 11/14/85 JF  
09/19/85 JF 06/04/86 JF  
10/02/85 JF 06/10/86 CR  
11/01/85 JF

FRONT:

---

30-3-9  
Fertilizer  
Plus  
Pre-Emergent  
Weed Control

---

- o preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed and Poa annua
  - o provides quick green-up from soluble nitrogen
- 

Net Weight 21 3/16 lbs (9.60 kg)

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In case of spills, either collect for use or dispose of properly.

ACTIVE INGREDIENT:  
    pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl  
    -2,6-dinitrobenzenamine\* . . . . . 1.79%

INERT INGREDIENTS: . . . . . 98.21%

Total 100.00%

EPA Reg. No. 538-214

EPA Est

Fertilizer Plus Pre-Emergent Weed Control      30-3-9      643-XXXF

FRONT CONTINUED:

Page 4

Guaranteed Analysis

Total nitrogen (N) . . . . .	30%
0.6% ammoniacal nitrogen	
22.9% urea, methylene ureas nitrogen	
6.5% water insoluble nitrogen	
Available phosphoric acid (P <sub>2</sub> O <sub>5</sub> ) . . . . .	3%
Soluble potash (K <sub>2</sub> O) . . . . .	9%

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA

U.S. Pat Nos 3,705,794; 3,989,470

\*Licensed under one or more of the following U.S. Patents, 3,920,742; 4,066,441

*Formulated For:* The O M Scott & Sons Company  
Marysville, Ohio 43041

(c) 1986 The O M Scott & Sons Company. All rights reserved.

---

Recommended for Use by Professional Turfgrass Managers

---

XXXXX

30-3-9  
Fertilizer Plus  
Pre-Emergent  
Weed Control

GUSSETS:

2. 100, 14

RIGHT GUSSET

XXXXX

30-3-9  
Fertilizer Plus  
Pre-Emergent  
Weed Control

LEFT GUSSET

30-3-9  
Fertilizer Plus  
Pre-Emergent  
Weed Control

XXXXX

**BACK:**

P. No. 14

30-3-9  
Fertilizer Plus  
Pre-Emergent  
Weed Control

30-3-9  
Fertilizer Plus  
Pre-Emergent  
Weed Control

**Directions for use**

**LEFT COLUMN**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass and Poa annua (except putting greens and tees), bermudagrass (non-winter overseeded), St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass, Bentgrass and Poa annua Precautions" and "Limitation of Liability" statements.

Not for use on bentgrass and Poa annua putting greens and tees, or on dichondra. Where bentgrass, bentgrass-containing turf and Poa annua are the desired species, a light rate application only is recommended.

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

Grassy Weeds

Dicot Weeds

crabgrass  
foxtail  
goosegrass  
Poa annua  
barnyardgrass  
fall panicum

chickweed  
oxalis  
prostrate spurge  
knotweed

To prevent goosegrass, crabgrass, foxtail, barnyardgrass, fall panicum, oxalis, knotweed and prostrate spurge, apply in the spring before they sprout.

To prevent Poa annua and fall germinating chickweed, apply product to dry foliage in late summer/early fall (before germination). Repeat application yearly for continued control.

Northern (Cool Season) Grasses

Apply 1.6 to 2.2 lbs. of product per 1,000 sq. ft. (1.25 to 1.75 lbs. of active ingredient per acre) prior to germination of weeds. See "Suggested Spreader Settings".

BACK CONTINUED:

P 17 01 14

Southern (Warm Season) Grasses

Apply 1.9 to 2.6 lbs. of product per 1,000 sq. ft. (1.5 to 2.0 lbs. of active ingredient per acre) prior to germination of weeds. See "Suggested Spreader Settings".

Where lower weed infestations are expected, the lighter rates may be used.

For best results, water after all applications (1/4 - 1/2 inch) to remove the granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying after applying.

RIGHT COLUMN

Precautions

Do not use on bentgrass and Poa annua putting greens and tees, or on dichondra. Where bentgrass and bentgrass-containing turf and Poa annua are the desired species, a light rate application only is recommended.

On cool season turfgrasses avoid applications when temperatures are regularly in the 80's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature, scalping - below 40°F, and excessive soil moisture).

Bermudagrass, Bentgrass and Poa annua Precautions

Do not apply to dormant bermudagrass, bentgrass or Poa annua (turf that is not actively growing or has experienced frost). However, for dormant bermudagrass, an application can be made if it is maintained at a high cut (1/2 inch or more) on medium-fine textured soils such as clays and clay loams.

Where turf thinning or damage has occurred due to winter injury of bermudagrass, wait until greenup, fill-in and good rooting before making an application.

Applications to overseeded bermudagrass may injure and thin the cool season variety. Do not apply to bentgrass or Poa annua after late summer.

BACK CONTINUED:

p. 13 of 14

#### Other Precautions

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Applications should not exceed twice yearly.

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water.

#### Storage and disposal

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

#### LIMITATION OF LIABILITY

It must be understood that this product has not been evaluated on all bentgrass varieties and Poa annua under all possible growing conditions. The user should exercise judgment and caution when using this product on a given variety until familiar with the performance under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT ON BENTGRASS AND POA ANNUA IF NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES. The exclusive remedy of the user or Buyer, and the limit of liability of O. M. Scott & Sons Company or any other Seller, for any and all losses, injured or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of O. M. Scott & Sons Company.

The O M Scott & Sons Company  
Marysville, Ohio 43041

**Suggested Spreader Settings**

To provide proper distribution calibrate spreader before application.  
Do not adjust rates in order to meet fertilizer requirements of your soil.

21 3/16 lbs treats 11,000 ft<sup>2</sup> (1/4 acre) at the 1.9 lbs product/1,000 sq ft rate (equivalent to 1.5 lb active ingredient/acre and 0.6 lb N/1,000 sq ft)

<u>SPREADER</u>	<u>GROUND OR PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>RECOMMENDED RATE RANGE</u>				
			<u>Lbs. Product/1,000 sq. ft.</u>	<u>1.6</u>	<u>1.9</u>	<u>2.2</u>	<u>2.6</u>
Scotts Drop Type	3 mph	overlap wheels		X	X	X	X
Gandy - 30"	3 mph	overlap wheels		X	X	X	X
Scotts Rotaries*							
R-7, R-7X	3 mph	XX feet		X	X	X	X
R-8 (cone 8)	3 mph	XX feet		X	X	X	X
Lely Models	4.5 mph	XX feet		X	X	X	X
Vicon	4.5 mph	XX feet		X	X	X	X

\*Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide 1/4 closed.


Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.  
Metric coverage: 9.60 kg treats 1022 m<sup>2</sup> at the 1.9 lbs. product/1,000 sq. ft. rate

FRONT

(S-1205T) — LOW END OF RANGE  
30-3-1 FERTILIZER PLUS  
PREEMERGENT WEED CONTROL

(RANGE)

09/11/85 JF 11/14/85 JF  
09/19/85 JF 06/04/86 JF  
10/02/85 JF 06/10/86 CR  
11/01/85 JF 06/20/90 MK/CR



P425  
538-214  
p. 1 of 14

(Scotts(R))  
ProTurf(R)  
BRAND

30-3-9  
Fertilizer  
Plus  
Preemergent  
Weed Control

- o preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed and Poa annua
- o provides quick green-up plus extended feeding with controlled release SREF nitrogen (slow release encapsulated fertilizer)

Net Weight 21 3/16 lbs (9.60 kg)

**BEST AVAILABLE COPY**

KEEP OUT OF REACH OF CHILDREN  
CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In case of spills, either collect for use or dispose of properly.

ACTIVE INGREDIENT:

pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl  
-2,6-dinitrobenzenamine\* . . . . . 1.79%

INERT INGREDIENTS: . . . . . 98.21%  
Total 100.00%

EPA Reg. No. 538-214

EPA Est 538-OH-1

Fertilizer Plus Preemergent Weed Control

30-3-9

643-XXXX

10/1/81

Guaranteed Analysis

Total nitrogen (N) . . . . .	30%
0.6% ammoniacal nitrogen	
22.9% urea, methylene ureas nitrogen	
6.5% water insoluble nitrogen	
Available phosphoric acid (P <sub>2</sub> O <sub>5</sub> ) . . . . .	3%
Soluble potash (K <sub>2</sub> O) . . . . .	9%

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA

U.S. Patent No's 3,705,794; 3,989,470

\*Licensed under one or more of the following U.S. Patents: 3,920,742;

4,066,441

ProTurf Division

The O M Scott & Sons Company

Marysville, Ohio 43041

(c) 1990 The O M Scott & Sons Company. All rights reserved.

Recommended for Use by Professional Turfgrass Managers

XXXXX

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

BEST AVAILABLE COPY

RIGHT GUESSET

XXXXX

(Scotts(R))  
ProTurf(R)

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

LEFT GUESSET

XXXXX

(Scotts(R))  
ProTurf(R)

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

(Scotts(R))  
ProTurf(R)

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

### Directions for use

#### LEFT COLUMN

✓ It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass and Poa annua (except putting greens and tees), bermudagrass, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass, Bentgrass and Poa annua Precautions" and "Limitation of Liability" statements.

✓ Not for use on bentgrass and Poa annua greens and tees, or on dichondra. Where bentgrass, bentgrass-containing turf and Poa annua are the desired species, a light rate application only is recommended.

✓ Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

#### Grassy Weeds

crabgrass  
foxtail  
goosegrass  
Poa annua  
barnyardgrass  
fall panicum

#### Dicot Weeds

chickweed  
oxalis  
prostrate spurge  
knotweed

✓ To prevent goosegrass, crabgrass, foxtail, barnyardgrass, fall panicum, oxalis, knotweed and prostrate spurge, apply in the spring before they sprout.

✓ To prevent Poa annua and fall germinating chickweed, apply product to dry foliage in late summer/early fall (before germination). Repeat application yearly for continued control.

#### Northern (Cool Season) Grasses

✓ Apply 1.3 to 2.2 lbs. of product per 1,000 sq. ft. (1.0 to 1.75 lbs. of active ingredient per acre) prior to germination of weeds. (See "Suggested Spreader Settings").

**Southern (Warm Season) Grasses**

Apply 1.3 to 2.6 lbs. of product per 1,000 sq. ft. (1.0 to 2.0 lbs. of active ingredient per acre) prior to germination of weeds. (See "Suggested Spreader Settings".

Where lower weed infestations are expected, the higher rates may be used. For extended weed control, a repeat application is recommended.

For best results, water after all applications (1/4 - 1/2 inch) to remove the granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying after applying.

**RIGHT COLUMN****Precautions**

Do not use on bentgrass and Poa annua putting greens and tees, or on dichondra. Where bentgrass and bentgrass-containing turf and Poa annua are the desired species, a light rate application only is recommended.

On cool season turfgrasses avoid applications when temperatures are regularly in the 80's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, scalping, and excessive soil moisture).

**Bermudagrass, Bentgrass and Poa annua Precautions**

When applied to dormant bermudagrass, greenup may be delayed 2-3 weeks in the spring. The delay in greenup may be more extensive on low-cut bermudagrass and on sandy soils.

Where turf thinning or damage has occurred due to winter injury of bermudagrass, wait until greenup, fill-in and good rooting before making an application.

Applications of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseeding.

Do not apply to bentgrass or Poa annua after late summer.

**Other Precautions**

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Applications should not exceed twice yearly.

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water. If metallic surfaces (golf course equipment, automobiles, ect.) are present at the application site, do not use this product under windy conditions (5-10 MPH) due to its staining potential.

**Storage and disposal**

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

LIMITATION OF LIABILITY

It must be understood that this product has not been evaluated on all bentgrass varieties and Poa annua under all possible growing conditions. The user should exercise judgment and caution when using this product on a given variety until familiar with the performance under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT ON BENTGRASS AND POA ANNUA IF NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES. The exclusive remedy of the user or Buyer, and the limit of liability of The O M Scott & Sons Company or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of The O M Scott & Sons Company.

The O M Scott & Sons Company  
Marysville, Ohio 43041

**BEST AVAILABLE COPY**

**Suggested Spreader Settings**

To provide proper distribution calibrate spreader before application.  
Do not adjust rates in order to meet fertilizer requirements of your soil.

21 3/16 lbs treats 11,000 ft<sup>2</sup> (1/4 acre) at the 1.9 lbs product/1,000 sq ft rate  
(equivalent to 1.5 lb active ingredient/acre and 0.6 lb N/1,000 sq ft)

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	RECOMMENDED RATE RANGE				
			Lbs product/1,000 sq ft				
			1.3	1.6	1.9	2.2	2.6
Scotts Drop Type	3 mph	overlap wheels	X	X	X	X	X
Gandy - 30"	3 mph	overlap wheels	X	X	X	X	X
Scotts Rotaries*							
R-7, R-7X	3 mph	XX feet	X	X	X	X	X
R-8, R-8A (cone X)	3 mph	XX feet	X	X	X	X	X
Lely Models	4.5 mph	XX feet	X	X	X	X	X
Vicon	4.5 mph	XX feet	X	X	X	X	X


\*Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide 1/4 closed.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.  
Metric coverage: 9.60 kg treats 1022 m<sup>2</sup> at the 1.9 lbs  
product/1,000 sq ft rate

FRONT

(S-1206T) --- LOW END OF RANGE  
30-3-9 FERTILIZER PLUS  
PREEMERGENT WEED CONTROL (RANGE)

09/11/85 JF 11/14/85 JF  
09/19/85 JF 06/04/86 JF  
10/02/85 JF 06/10/86 CR  
11/01/85 JF 06/20/90 MK/CR



(Scotts(R))  
ProTurf(R)  
BRAND

30-3-9  
Fertilizer  
Plus  
Preemergent  
Weed Control

- o preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed and Poa annua
- o provides quick green-up plus extended feeding with controlled release SREF nitrogen (slow release encapsulated fertilizer)

Net Weight 75 3/4 lbs (34.36 kg)

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
PRECAUTIONARY STATEMENTS



Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In case of spills, either collect for use or dispose of properly.

ACTIVE INGREDIENT:

pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl  
-2,6-dinitrobenzenamine\*

INERT INGREDIENTS:	C. 50%
	95.50%
	Total 100.00%

EPA Reg. No. 538-214

EPA Est 538-OH-1

Fertilizer Plus Preemergent Weed Control

30-3-9

643-XXXX

Guaranteed Analysis

Total nitrogen (N) . . . . .	30%
0.6% ammoniacal nitrogen	
22.9% urea, methylene ureas nitrogen	
6.5% water insoluble nitrogen	
Available phosphoric acid (P <sub>2</sub> O <sub>5</sub> ) . . . . .	3%
Soluble potash (K <sub>2</sub> O) . . . . .	9%

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA

U.S. Patent No's 3,705,794; 3,989,470

\*Licensed under one or more of the following U.S. Patents: 3,920,742; 4,066,441

ProTurf Division

The O M Scott & Sons Company

Marysville, Ohio 43041

(c) 1990 The O M Scott & Sons Company. All rights reserved.

Recommended for Use by Professional Turfgrass Managers

XXXXX

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

RIGHT GUESSET

XVXXX

(Scotts(R))  
ProTurf(R)

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

LEFT GUESSET

XXXXX

(Scotts(R))  
ProTurf(R)

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

✓

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

(Scotts(R))  
ProTurf(R)

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

Directions for use

LEFT COLUMN

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass and Poa annua (except putting greens and tees), bermudagrass, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass, Bentgrass and Poa annua Precautions" and "Limitation of Liability" statements.

Not for use on bentgrass and Poa annua greens and tees, or on dichondra. Where bentgrass, bentgrass-containing turf and Poa annua are the desired species, a light rate application only is recommended.

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

Grassy Weeds

Dicot Weeds

- crabgrass
- foxtail
- goosegrass
- Poa annua
- barnyardgrass
- fall panicum

- chickweed
- oxalis
- prostrate spurge
- knotweed

To prevent goosegrass, crabgrass, foxtail, barnyardgrass, fall panicum, oxalis, knotweed and prostrate spurge, apply in the spring before they sprout.

To prevent Poa annua and fall germinating chickweed, apply product to dry foliage in late summer/early fall (before germination). Repeat application yearly for continued control.

Northern (Cool Season) Grasses

Apply 4.6 to 8.0 lbs. of product per 1,000 sq. ft. (1.0 to 1.75 lbs. of active ingredient per acre) prior to germination of weeds. (See "Suggested Spreader Settings".

Southern (Warm Season) Grasses

Apply 4.6 to 9.2 lbs. of product per 1,000 sq. ft. (1.0 to 2.0 lbs. of active ingredient per acre) prior to germination of weeds. (See "Suggested Spreader Settings").

Where lower weed infestations are expected, the higher rates may be used. For extended weed control, a repeat application is recommended.

For best results, water after all applications (1/4 - 1/2 inch) to remove the granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying after applying.

## RIGHT COLUMN

## Precautions

Do not use on bentgrass and Poa annua putting greens and tees, or on dichondra. Where bentgrass and bentgrass-containing turf and Poa annua are the desired species, a light rate application only is recommended.

On cool season turfgrasses avoid applications when temperatures are regularly in the 80's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, scalping, and excessive soil moisture).

Bermudagrass, Bentgrass and Poa annua Precautions

When applied to dormant bermudagrass, greenup may be delayed 2-3 weeks in the spring. The delay in greenup may be more extensive on low-cut bermudagrass and on sandy soils.

Where turf thinning or damage has occurred due to winter injury of bermudagrass, wait until greenup, fill-in and good rooting before making an application.

Applications of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseeding.

Do not apply to bentgrass or Poa annua after late summer.

BACK CONTINUED:

### Other Precautions

- ✓ Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.
- ✓ On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.
- ✓ Applications should not exceed twice yearly.
- ✓ Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water. If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 MPH) due to its staining potential.

### Storage and disposal

- ✓ STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.
- ✓ PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
- ✓ CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### LIMITATION OF LIABILITY

It must be understood that this product has not been evaluated on all bentgrass varieties and Poa annua under all possible growing conditions. The user should exercise judgment and caution when using this product on a given variety until familiar with the performance under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT ON BENTGRASS AND POA ANNUA IF NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES. The exclusive remedy of the user or Buyer, and the limit of liability of The O M Scott & Sons Company or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of The O M Scott & Sons Company.

The O M Scott & Sons Company  
Marysville, Ohio 43041