



PM 2 ) 538-214 )

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

1/17

FEB 12 1996

Vincent Snyder  
The Scotts Company  
14111 Scottslawn Road  
Marysville, OH 43041

Dear Mr. Snyder:

Subject: Label Amendments for:

Southern Weedgrass Control  
EPA Registration Number 538-188  
Your Application Dated 2/1/96, 1/19/96

Turf Fertilizer Plus Preemergent Weed Control  
EPA Registration Number 538-213  
Your Application Dated 1/19/96

Proturf Weedgrass Control 60 WDG  
EPA Registration Number 538-195  
Your Application Dated 1/19/96

Turf Weedgrass Control 8637  
EPA Registration Number 538-193  
Your Application Dated 1/19/96

Proturf Fertilizer Plus Southern Turf Weedgrass Control  
EPA Registration Number 538-206  
Your Application Dated 1/19/96

Proturf Fertilizer Plus Turf Weedgrass Control  
EPA Registration Number 538-196  
Your Application Dated 1/22/96

Fertilizer Plus Preemergent Weed Control  
EPA Registration Number 538-214  
Your Application Dated 1/22/96

7505C  
Fault

The labeling referred to above submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable subject to the following comment.

1. A Statement of Practical Treatment section is added to each product label. Please see the attached sample format for guidance.



Recycled/Recyclable  
Printed with Soy/Candla Ink on paper that contains at least 50% recycled fiber

2/17

Your amended labels have been stamped accepted with comment and the copies are enclosed. You must incorporate the comment above on your labels. The new labels must be on products released for shipment within 1 year or by the next label printing, whichever occurs first.

Sincerely,



Robert J. Taylor  
Product Manager 25  
Fungicide-Herbicide Branch  
Registration Division (H7505C)

) Enclosures



Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 538-214

06/10/88 CR 06/20/90 MK/CR 12/12/90 CR  
12/20/90 CR 06/15/93 VS/cr 01/23/96 VS/kl

(Scotts®)  
ProTurf®  
BRAND

**ACCEPTED  
with COMMENTS  
In EPA Letter Dated**

30-3-9  
Fertilizer  
Plus  
Preemergent  
Weed Control

FEB 12 1998  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.  
538-214

- preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed and *Poa annua*
- provides quick green-up plus extended feeding with controlled release nitrogen

Net Weight 75 ¼ lbs (34.36 kg)

**KEEP OUT OF REACH OF CHILDREN  
CAUTION  
PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals:** CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

**Personal Protective Equipment (PPE)**

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

5/17

FRONT CONTINUED

**User Safety Recommendations:**  
 Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards:** This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In case of spills, either collect for use or dispose of properly.

**ACTIVE INGREDIENT:**

pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl

-2,6-dinitrobenzenamine\* ..... 0.50%

INERT INGREDIENTS: ..... 99.50%

Total 100.00%

EPA Reg. No. 538-214

EPA Est 538-OH-1

Fertilizer Plus Preemergent Weed Control 30-3-9

643-XXXX

**Guaranteed Analysis**

Total nitrogen (N)..... 30%

0.6% ammoniacal nitrogen

22.9% urea, methylene ureas nitrogen

6.5% water insoluble nitrogen

Available phosphate (P<sub>2</sub>O<sub>5</sub>) ..... 3%

Soluble potash (K<sub>2</sub>O)..... 9%

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA

U.S. Patent No's 3,705,794; 3,989,470

\*Licensed under one or more of the following U.S. Patents: 3,920,742; 4,066,441

The Scotts Company

Marysville, Ohio 43041

(c) 1996. The Scotts Company. World rights reserved.

Recommended for Use by Professional Turfgrass Managers

**OR (Optional)**

For sale to and use by Professional Lawn Care (or Service) Applicators only

XXXXX

30-3-9

Fertilizer Plus

Preemergent

Weed Control

6/17

RIGHT GUSSET

XXXXX

(Scotts®)  
ProTurf®

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

LEFT GUSSET

XXXXX

(Scotts®)  
ProTurf®

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

7/17

**BACK:**

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

(Scotts®)  
ProTurf®

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

**Directions for use**

**LEFT COLUMN**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Workers Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the worker protection standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, waterproof gloves, shoes plus socks.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass and *Poa annua* (except putting greens and tees), bermudagrass, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass, Bentgrass and *Poa annua* Precautions" and "Limitation of Liability" statements.

Not for use on bentgrass and *Poa annua* greens and tees, or on dichondra. Where bentgrass, bentgrass-containing turf and *Poa annua* are the desired species, a light rate application only is recommended.

**BACK CONTINUED:**

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

Grassy Weeds

Dicot Weeds

- crabgrass
- foxtail
- goosegrass
- Poa annua*
- barnyardgrass
- fall panicum

- chickweed
- oxalis
- prostrate spurge
- knotweed

To prevent goosegrass, crabgrass, foxtail, barnyardgrass, fall panicum, oxalis, knotweed and prostrate spurge, apply in the spring before they sprout.

To prevent *Poa annua* and fall germinating chickweed, apply product to dry foliage in late summer/early fall (before germination). Repeat application yearly for continued control.

Northern (Cool Season) and Southern (Warm Season) Grasses

Apply 4.6 to 9.2 lbs. of product per 1,000 sq. ft. (1.0 to 2.0 lbs. of active ingredient per acre) prior to germination of weeds. (See "Suggested Spreader Settings.")

Where lower weed infestations are expected, the lighter rates may be used. For extended weed control, a repeat application is recommended.

For best results, water after all applications (¼ - ½ inch) to remove the granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying after applying.

**RIGHT COLUMN**

**Precautions**

Do not use on bentgrass and *Poa annua* putting greens and tees, or on dichondra. Where bentgrass and bentgrass-containing turf and *Poa annua* are the desired species, a light rate application only is recommended.

On cool season turfgrasses avoid applications when temperatures are regularly in the 80's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, scalping, and excessive soil moisture).



9/17

**BACK CONTINUED:**

**Bermudagrass, Bentgrass and *Poa annua* Precautions**

When applied to dormant bermudagrass, greenup may be delayed 2-3 weeks in the spring. The delay in greenup may be more extensive on low-cut bermudagrass and on sandy soils.

Where turf thinning or damage has occurred due to winter injury of bermudagrass, wait until greenup, fill-in and good rooting before making an application.

Applications of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseeding.

Do not apply to bentgrass or *Poa annua* after late summer.

**Other Precautions**

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Applications should not exceed twice yearly.

**NOTE TO EPA -- EITHER ONE OF THE FOLLOWING OPTIONS WILL BE USED**

OPTION A (CURRENT STATEMENT)

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water. If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 MPH) due to its staining potential.

OPTION B

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water. Exercise caution when metallic surfaces (automobiles, etc.) are in the area of application as drift under windy conditions may cause staining.

**Storage and disposal**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

19/17

**BACK CONTINUED:**

LIMITATION OF LIABILITY

It must be understood that this product has not been evaluated on all bentgrass varieties and *Poa annua* under all possible growing conditions. The user should exercise judgment and caution when using this product on a given variety until familiar with the performance under his growing conditions. **NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT ON BENTGRASS AND *POA ANNUA* IF NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES.** The exclusive *remedy* of the user or Buyer, and the limit of liability of The Scotts Company, or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of The Scotts Company.

The Scotts Company  
 Marysville, Ohio 43041

**Suggested Spreader Settings**

To provide proper distribution calibrate spreader before application.  
 Do not adjust rates in order to meet fertilizer requirements of your soil.

75 3/4 lbs treats 11,000 ft<sup>2</sup> (1/4 acre) at the 6.9 lbs product/1,000 sq ft rate  
 (equivalent to 1.5 lb active ingredient/acre and 2.0 lb N/1,000 sq ft)

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	<u>RECOMMENDED RATE RANGE</u> Lbs product/1,000 sq ft				
			4.6	5.7	6.9	8.0	9.2
Scotts Drop Type	3 mph	overlap wheels	X	X	X	X	X
Gandy - 30"	3 mph	overlap wheels	X	X	X	X	X
Scotts Rotaries*							
R-7, R-7X	3 mph	XX feet	X	X	X	X	X
R-8, R-8A							
(cone X)	3 mph	XX feet	X	X	X	X	X
Lely Models	4.5 mph	XX feet	X	X	X	X	X
Vicon	4.5 mph	XX feet	X	X	X	X	X

\*Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide 1/4 closed.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.  
 Metric coverage: 22.67 kg treats 1022 m<sup>2</sup> at the 6.9 lbs  
 product/1,000 sq ft rate

11/17

FRONT

(S-1205T) -- HIGH END C/F RANGE  
30-3-9 FERTILIZER PLUS  
PREEMERGENT WEED CONTROL

09/11/85 JF	09/19/85 JF	10/02/85 JF
11/01/85 JF	11/14/85 JF	06/04/86 JF
06/10/86 CR	06/20/90 MK/cr	12/12/90 CR
12/20/90 CR	06/15/93 VS/cr	01/23/96 VS/kl

(Scotts®)  
ProTurf®  
BRAND

30-3-9  
Fertilizer  
Plus  
Preemergent  
Weed Control

- preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, knotweed and *Poa annua*
- provides quick green-up plus extended feeding with controlled release nitrogen

Net Weight 21<sup>3/16</sup> lbs (9.60 kg)

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals:** CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

**Personal Protective Equipment (PPE)**

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

12/17

**FRONT CONTINUED**

**User Safety Recommendations:**  
 Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards:** This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In case of spills, either collect for use or dispose of properly.

**ACTIVE INGREDIENT:**

pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl  
 -2,6-dinitrobenzenamine\* ..... 1.79%

**INERT INGREDIENTS:** ..... 98.21%  
 Total ..... 100.00%

EPA Reg. No. 538-214 ..... EPA Est 538-OH-1

Fertilizer Plus Preemergent Weed Control 30-3-9 643-XXXX

**Guaranteed Analysis**

Total nitrogen (N) ..... 30%  
 0.6% ammoniacal nitrogen  
 22.9% urea, methylene ureas nitrogen  
 6.5% water insoluble nitrogen  
 Available phosphate (P<sub>2</sub>O<sub>5</sub>) ..... 3%  
 Soluble potash (K<sub>2</sub>O) ..... 9%

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA

U.S. Patent No's 3,705,794; 3,989,470

\*Licensed under one or more of the following U.S. Patents: 3,920,742; 4,066,441

The Scotts Company

Marysville, Ohio 43041

(c) 1996. The Scotts Company. World rights reserved.

Recommended for Use by Professional Turfgrass Managers

**OR (Optional)**

For sale to and use by Professional Lawn Care (or Service) Applicators only

XXXXX                      ✓    30-3-9  
    Fertilizer Plus  
    Preemergent  
    Weed Control

**RIGHT GUSSET**

XXXXX

(Scotts®)  
ProTurf®

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

**LEFT GUSSET**

XXXXX

(Scotts®)  
ProTurf®

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

18/17

**BACK:**

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

(Scotts®)  
ProTurf®

30-3-9  
Fertilizer Plus  
Preemergent  
Weed Control

**Directions for use**

**LEFT COLUMN**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application.

Use this product only in accordance with its labeling and with the Workers Protection Standard, 40CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label (in this labeling) about personal protective equipment and restricted-entry interval.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the worker protection standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, waterproof gloves, shoes plus socks.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass and *Poa annua* (except putting greens and tees), bermudagrass, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass, Bentgrass and *Poa annua* Precautions" and "Limitation of Liability" statements.

Not for use on bentgrass and *Poa annua* greens and tees, or on dichondra. Where bentgrass, bentgrass-containing turf and *Poa annua* are the desired species, a light rate application only is recommended.

15/17

**BACK CONTINUED:**

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

Grassy Weeds

Dicot Weeds

- crabgrass
- foxtail
- goosegrass
- Poa annua*
- barnyardgrass
- fall panicum

- chickweed
- oxalis
- prostrate spurge
- knotweed

To prevent goosegrass, crabgrass, foxtail, barnyardgrass, fall panicum, oxalis, knotweed and prostrate spurge, apply in the spring before they sprout.

To prevent *Poa annua* and fall germinating chickweed, apply product to dry foliage in late summer/early fall (before germination). Repeat application yearly for continued control.

Northern (Cool Season) and Southern (Warm Season) Grasses

Apply 1.3 to 2.6 lbs. of product per 1,000 sq. ft. (1.0 to 2.0 lbs. of active ingredient per acre) prior to germination of weeds. (See "Suggested Spreader Settings.")

Where lower weed infestations are expected, the lighter rates may be used. For extended weed control, a repeat application is recommended.

For best results, water after all applications (1/4 - 1/2 inch) to remove the granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying after applying.

**RIGHT COLUMN**

**Precautions**

Do not use on bentgrass and *Poa annua* putting greens and tees, or on dichondra. Where bentgrass and bentgrass-containing turf and *Poa annua* are the desired species, a light rate application only is recommended.

On cool season turfgrasses avoid applications when temperatures are regularly in the 80's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, scalping, and excessive soil moisture).

10/17

**BACK CONTINUED:**

**Bermudagrass, Bentgrass and *Poa annua* Precautions**

When applied to dormant bermudagrass, greenup may be delayed 2-3 weeks in the spring. The delay in greenup may be more extensive on low-cut bermudagrass and on sandy soils.

Where turf thinning or damage has occurred due to winter injury of bermudagrass, wait until greenup, fill-in and good rooting before making an application.

Applications of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseeding.

Do not apply to bentgrass or *Poa annua* after late summer.

**Other Precautions**

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Applications should not exceed twice yearly.

**NOTE TO EPA -- EITHER ONE OF THE FOLLOWING OPTIONS WILL BE USED**

**OPTION A (CURRENT STATEMENT)**

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water. If metallic surfaces (golf course equipment, automobiles, etc.) are present at the application site, do not use this product under windy conditions (5-10 MPH) due to its staining potential.

**OPTION B**

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water. Exercise caution when metallic surfaces (automobiles, etc.) are in the area of application as drift under windy conditions may cause staining.

**Storage and disposal**

Do not contaminate water, food or feed by storage or disposal.

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



17/17

**BACK CONTINUED:**

LIMITATION OF LIABILITY

It must be understood that this product has not been evaluated on all bentgrass varieties and *Poa annua* under all possible growing conditions. The user should exercise judgment and caution when using this product on a given variety until familiar with the performance under his growing conditions. **NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT ON BENTGRASS AND *POA ANNUA* IF NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES.** The exclusive remedy of the user or Buyer, and the limit of liability of The Scotts Company, or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of The Scotts Company.

The Scotts Company  
14111 Scottslawn Rd.  
Marysville, Ohio 43041

**Suggested Spreader Settings**

To provide proper distribution calibrate spreader before application.  
Do not adjust rates in order to meet fertilizer requirements of your soil.

21<sup>3/16</sup> lbs treats 11,000 ft<sup>2</sup> (1/4 acre) at the 1.9 lbs product/1,000 sq ft rate  
(equivalent to 1.5 lb active ingredient/acre and 0.6 lb N/1,000 sq ft)

<u>SPREADER</u>	<u>GROUND OR PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>RECOMMENDED RATE RANGE</u>				
			<u>Lbs product/1,000 sq ft</u>				
			<u>1.3</u>	<u>1.6</u>	<u>1.9</u>	<u>2.2</u>	<u>2.6</u>
Scotts Drop Type	3 mph	overlap wheels	X	X	X	X	X
Gandy - 30"	3 mph	overlap wheels	X	X	X	X	X
Scotts Rotaries*							
R-7, R-7X	3 mph	XX feet	X	X	X	X	X
R-8, R-8A							
(cone X)	3 mph	XX feet	X	X	X	X	X
Lely Models	4.5 mph	XX feet	X	X	X	X	X
Vicon	4.5 mph	XX feet	X	X	X	X	X

\*Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide 1/2 closed.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm  
Metric coverage: 9.60 kg treats 1022 m<sup>2</sup> at the 1.9 lbs product/1,000 sq ft rate