

(C

N

FRONT CONTINUED:

Net Weight 13 1/4 lbs (6.00 kg)

33-3-3

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS

<u>Hazards to Humans and Domestic Animals</u>: CAUTION. Harmful if swallowed. Avoid inhalation and contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Avoid contamination of feed or foodstuffs. Do not contaminate water used for inrigational or domestic purposes.

<u>Environmental Hazards</u>: Do not apply directly to water. Do not apply when weather conditions favor drift from target area. Do not contaminate water used for irrigation or domestic purposes.


STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

ACTIVE INGREDIENTS: 2-(2-Methyl-4-chlorophenoxy) propionic acid . . . . . . . . . . . . . 1.18% 97.64% Total 100.00% EPA Reg No 538-XXX EPA Est 538-0H-1 Lawn Pro Weed and Feed 33-3-3 Guaranteed Analysis 0.6% ammoniacal nitrogen 25.1% urea, methylene ureas nitrogen 7.3% water insoluble nitrogen 3% 3% Derived from: monoammonium phosphate, urea, methylene ureas, and potassium sulfate. U S Pat Nos. 3,705,794 and 3,989,470 Product of USA The O M Scott & Sons Company Marysville, Ohio 43041 Allowed States I. C. Market 

RIGHT GUSSET:



<u>F</u>

BACK:

PULL TAB and remove for pour spout (ILLUS.) Easy Open PULL HÉRE

Treats 5,000 sq ft (465 m<sup>2</sup>)

(

(Scotts (R)) (Lawn Pro ) (R)

Weed and Feed weed control plus fertilizer

LPWC+LF 24612

5M

PURCHASE

(Scotts)

PROOF

OF

## Advantages

. Lightweight and concentrated--made by Scotts patented Polyform process

. Safe feeding--won't burn when applied as directed

. Quick-greening--usually in 3 to 5 days

. Long-lasting--feeds up to 2 months

. Dependable--traditional Scotts quality

. Controls more than 50 weeds...

aster bittercress black medic bristly oxtongue buckhorn bull thistle burclover buttonweed carpetweed centella chicory chickweed cinquefoil cudweed curly dock dandelion dichondra dollarweed	English daisy eveningprimrose filaree fleabane ground ivy healall henbit horseweed knotweed (seedling) lambsquarter lippia mallow moneywort rorningglory mustard pennywort pepperweed	pigweed plantain prickly lettuce puncturevine purslane pusley, Florida ragweed scarlet pimpernel shadflower shepherdspurse smartweed sowthistle stitchwort thymeleaf sandwort vetch white clover yellow rocket			
It is a violation of federal law to use this product in a manner inconsistent with its labeling.					
APPLICATION	-		,	•	••••

. Apply any time weeds are actively growing (when daytime temperatures are consistently above 60°F).

. Apply on a calm day.

E.

BACK CONTINUED:

. Apply to moist grass. To help particles stick to weed leaves the grass should be thoroughly watered or wet from dew before applying Lawn Pro Weed and Feed. 5.4

. Do not use this product on flowers, shrubs, fruit or vegetable plants or on creeping bentgrass, St. Augustine, lippia and dichondra lawns. On bermudagrass lawns, under certain conditions, some temporary discoloration may result.

#### WATERING OR RAIN

Do not water your lawn for 24 hours after application. Water will wash the weed control off the weed's leaves and reduce the effectiveness of this product.

This product should not be applied if rain is expected within the next 24 hours.

### SPREADER

Be careful when applying this product near shrubs, flower beds or vegetable gardens as contact with weed control may damage these desirable plants. The Scotts PF Drop Spreaders provide excellent controlled application to protect these areas. When using the EasyGreen<sup>TM</sup> Rotary Spreader take special care mear desirable plants.

Spreader Settings	Scotts Drop Spreaders	Scotts EasyGreenTH Rotary
(treats 5,000 sq ft)	X	xx
IMPORTANT	-	-

WHAT TO EXPECT

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control.

This product will not prevent weed seeds from producing new plants.

EXTREME TEMPERATURES

Avoid applying when temperatures are consistently above 90 degrees.

SEEDING/SODDING

Wait until the new grass has been mowed at least four times before applyin Lawn Pro Weed and Feed weed control.

If you are planning to seed, you should wait at least one month after you happlied Lawn Pro Weed and Feed.

BACK CONTINUED:

a start

TIPS ON SPREADING WITH SCOTTS SPREADERS

Scotts Drop:

- First apply 2 header strips
- Operate spreader the longest way of the lawn (ILLUS.)
- Overlap wheels to insure uniform feeding
- Shut off spreader when you have reached the header str:ps or stopped
  (ILLUS.)
- Mow lawn first if grass is tall enough to touch the bottom of spreader

# EasyGreen<sup>TM</sup> Rotary:

08250

- Slide shut-off lever to the right. Application automatically starts and stops when you do
- Avoid misses by making each pass about 30 inches from the previous one so some overlapping occurs

Your Do-It-Yourself Lawn Care Headquarters

Scotts Lawn Pro retailers can provide you with the do-it-yourself expertise and products to have a beautiful Scotts lawn simply and easily...for less money than hiring a lawn service. That's why we call them Lawn Pros!

## 24612

Ω

.....

(UPC)

32247 24612

Scotts Policy No-Quibble Guarantee

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly. Scotts...the lawn people Marysville, Ohio 43041

(c) 1986. The O M Scott & Sons Company. All-rights reserved.

77.7.0