

(S-1...)
07/10/86 AR 08/20/86 AR
08/14/86 AR 07/21/88 AR
06/16/89 VS 10/05/89 VS

24687

(Scotts (R))
(Lawn Pro)
(R)

Weed and Feed
Weed Control Plus Lawn Fertilizer

(NOTE: The description above will be inverted.)

24687

Treats 15,000 sq ft (1395 m²)

(Scotts (R))
(Lawn Pro)
(R)

SCOTTS POLICY
No-Quibble
Guarantee
SEE BACK FOR DETAILS

WEED AND FEED
weed control plus lawn fertilizer



Controls dandelions
and over 50 other broadleaf weeds

Thicker, greener turf
--satisfaction guaranteed

SCOTTS(R) Lawn Pro(R) PRODUCT

Available only at authorized Lawn Pro stores

ACCEPTED
with COMMENTS
in EPA Letter Dated

MAY 30 1990

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

538-215

Net Weight 39 3/4 lbs (18.00 kg)

33-3-3

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION. Harmful if swallowed. Avoid inhalation and contact with skin, eyes, or clothing. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash thoroughly with soap and water after handling. Avoid contamination of feed or foodstuffs. Do not contaminate water used for irrigational or domestic purposes. Wear long pants and a long-sleeved shirt while loading spreader and applying product.

Environmental Hazards: Do not apply when weather conditions favor drift from target area. Do not contaminate water used for irrigation or domestic purposes. Drift or runoff may adversely affect non-target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate washwaters.

Phenoxy herbicides such as MCPP generally will not reach groundwater from use as specified on product labels. To prevent contamination of water supplies, caution should be exercised when handling MCPP pesticides at mixing, loading, and disposal sites.

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic acid	1.18%
2-(2-Methyl-4-chlorophenoxy) propionic acid	1.18%
INERT INGREDIENTS:	97.64%
Total	100.00%

EPA Reg No 538-215
Lawn Pro Weed and Feed

643-1166

33-3-3

EPA Est 538-OH-1

Guaranteed Analysis

Total nitrogen(N)	33%
0.6% ammoniacal nitrogen	
25.1% urea, methylene ureas nitrogen	
7.3% water insoluble nitrogen	
Available phosphoric acid (P ₂ O ₅)	3%
Soluble potash (K ₂ O)	3%
Derived from: monoammonium phosphate, urea, methylene ureas, and potassium sulfate.	

U S Pat Nos. 3,705,794 and 3,989,470

Product of USA

The O M Scott & Sons Company
Marysville, Ohio 43041

RIGHT GUSSET:

24687	A SCOTT'S Lawn Pro PRODUCT	(Scotts (R)) (Lawn Pro) (R)	Weed and Feed Weed Control Plus Fertilizer	SCOTT'S POLICY No-Quibble Guarantee SEE BACK FOR DETAILS
-------	----------------------------------	------------------------------------	--	---

LEFT GUSSET:

(Scotts (R)) (Lawn Pro) (R)	Weed and Feed Weed Control Plus Fertilizer	SCOTT'S POLICY No-Quibble Guarantee SEE BACK FOR DETAILS	A SCOTT'S Lawn Pro PRODUCT	24687
------------------------------------	--	---	----------------------------------	-------

PULL TAB
and remove for
pour spout
(ILLUS.)
Easy
Open PULL HERE

LPWC+LF 24687
15M
PROOF
OF
PURCHASE
(Scotts)

Treats 15,000 sq ft (1395 m²)

(Scotts (R))
(Lawn Pro)
(R)

Weed and Feed
Weed Control Plus Fertilizer

Advantages

- . Lightweight and concentrated-- made by Scotts patented Polyform process
- . Safe feeding --won't burn when applied as directed
- . Quick--greening-- usually in 3 to 5 days
- . Long--lasting --feeds up to 2 months
- . Dependable-- traditional Scotts quality
- . Controls more than 50 weeds...

aster	English daisy	pigweed
bittercress	eveningprimrose	plantain
black medic	filaree	prickly lettuce
bristly oxtongue	fleabane	puncturevine
buckhorn	ground ivy	purslane
bull thistle	healall	pusley, Florida
burclover	henbit	ragweed
buttonweed	horsweed	scarlet
carpetweed	knotweed	pimpernel
centella	(seedling)	shadflower
chicory	lambsquarter	shepherdspurse
chickweed	lippia	smartweed
cinquefoil	mallow	sowthistle
cudweed	moneywort	stitchwort
curly dock	morninglory	thymeleaf
dandelion	mustard	sandwort
dichondra	pennywort	vetch
dollarweed	pepperweed	white clover
		yellow rocket

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

APPLICATION

- . Apply any time weeds are actively growing (when daytime temperatures are consistently above 60°F).
- . Apply on a calm day.
- . Apply to moist grass. To help particles stick to weed leaves the grass should be thoroughly watered or wet from dew before applying Lawn Pro Weed and Feed.
- . Do not use this product on flowers, shrubs, fruit or vegetable plants or on creeping bentgrass, St. Augustine, lippia and dichondra lawns. On bermudagrass lawns, under certain conditions, some temporary discoloration may result.

WATERING OR RAIN

Do not water your lawn for 24 hours after application. Water will wash the weed control off the weed's leaves and reduce the effectiveness of this product.

This product should not be applied if rain is expected within the next 24 hours.

SPREADER

Be careful when applying this product near shrubs, flower beds or vegetable gardens as contact with weed control may damage these desirable plants. The Scotts PF Drop Spreaders provide excellent controlled application to protect these areas. When using the EasyGreen™ Rotary Spreader take special care near desirable plants.

Spreader Settings

	Scotts Drop Spreaders	Scotts EasyGreen™ Rotary	Scotts Speedy Green™
(treats 15,000 sq ft)	4 1/2	25	4

RIGHT COLUMN

IMPORTANT

WHAT TO EXPECT

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control.

This product will not prevent weed seeds from producing new plants.

EXTREME TEMPERATURES

Avoid applying when temperatures are consistently above 90 degrees.

SEEDING/SODDING

Wait until the new grass has been mowed at least four times before applying Lawn Pro Weed and Feed weed control.

If you are planning to seed, you should wait at least one month after you have applied Lawn Pro Weed and Feed.

TIPS ON SPREADING WITH SCOTTS SPREADERS

Scotts Drop:

- First apply 2 header strips
- Operate spreader the longest way of the lawn (ILLUS.)
- Overlap wheels to insure uniform feeding
- Shut off spreader when you have reached the header strips or stopped (ILLUS.)
- Mow lawn first if grass is tall enough to touch the bottom of spreader

EasyGreen™ Rotary:

- Slide shut-off lever to the right. Application automatically starts and stops when you do
- Avoid misses by making each pass about 30 inches from the previous one so some overlapping occurs

Speedy Green™ Rotary:

- Shut off spreader anytime you stop
- Avoid misses by making each pass about 5 feet from the previous one so some overlapping occurs

Additional Spreader Settings (Non-Scotts spreader settings are approximate. You may need to adjust for proper coverage.)

Cyclone Rotary (Model B)	3 1/2
Central Drop (#205)	6
Sears Drop (#19810)	8
Ortho Drop (#5000)	6
Greenview Drop (#105)	7
Precision Drop	5 1/2
(Red Devil)	

Your Do-It-Yourself Lawn Care Headquarters

Scotts Lawn Pro retailers can provide you with the do-it-yourself expertise and products to have a beautiful Scotts lawn simply and easily...for less money than hiring a lawn service. That's why we call them Lawn Pros!

**Scotts Policy
No-Quibble Guarantee**

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

Scotts...the lawn people
Marysville, Ohio 43041

0 (UPC) 0

32247 24687

(c) 1989. The O M Scott & Sons Company. All rights reserved.

(NOTE: The description below will be inverted.)

24687

(Scotts (R))
(Lawn Pro)
(R)

Weed and Feed
Weed Control Plus Lawn Fertilizer