FRONT:

awm Pro Weed & Feed 24612

(S-1858)

07/16/86 AR 08/20/86 AR 08/14/86 AR 07/21/88 AR

06/16/89 VS

04/04/90 VS/CR 💍

05/14/90 VS/CR

24612

(Scotts (R))

(Lawn Pro)

Weed and Feed

Weed Control Plus Lawn Fertilizer

(NOTE: The description above will be inverted.)

(R)

Treats $5,000 \text{ sd} \text{ ft } (465 \text{ m}^2)$

> (Scotts (R)) (Lawn Pro)

(R)

24612

SCOTTS POLICY No-Quibble

Guarantee

SEE BACK FOR DETAILS

WEED AND FEED weed control plus lawn fertilizer

weeds (photo)

grass (photo) ACCEPTED

JUL 1 2 1990

Under the Federal Insecticide, Fungiciae, and Rodenticide Act, as amended, for the pesticide ternstered under 539-215 EPA Reg. No.

Controls dandelions and over 50 other broadleaf weeds Thicker, greener turf -- :atisfaction quaranteed

SCOTTS(R) Lawn Pro(R) PRODUCT

Available only at authorized Lawn Pro stores

Net Weight 14.59 lbs

27-3-6

KEEP OUT OF REACH OF CHILDREN CAUTION PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION. Harmful if swallowed. Avoid inhalation and contact with skin, eyes, or clothing. In care of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Avoid contamination of feed or foodstuffs. Do not '''' contaminate water used for irrigational or domestic purposes. pents: long-sleeved shirt and shoes when applying this product. **

menter or permit others to reenter treated areas until dusts have

Environmental Hazards: Do not apply when weather conditions favor drift from target area. Do not contaminate water used for irrigation or domestic purposes, Prift or runoff may adversely affect non-target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

STORAGE: Store in a clean, dry place. Reseal opened bag by fulding top down and securing.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: "Do not reuse bag. Discard bag in trash.

ACTIVE INGRELIENTS:

2,4—Dichlorophenoxyacetic acid		. 1.18%
2-(2-Methyl-4-chlorophenoxy) propionic acid		. 1.18%
INERT INGREDIENTS:		
	Total	100 00%

EPA Reg No 538-215 6
Lawn Pro Weed and Feed

643-1178

EPA Est 538-OH-1

27-3-6

Guaranteed Analysis

Total nitrogen(N)	
0.8% water insoluble nitrogen Available phosphoric acid (P ₂ O ₅)	
Derived from: monoammonium phosphate, urea, methylene ureas, and muriate of potash.	:
Chlorine, not more than	

* Contains 6.0% slowly available methylenediurea and dimethylenetriurea nitrogen.

U S Pat No 4,378,238

Product of USA

The O M Scott & Sons Company Marysville, Ohio 43041

Ground white see

FRONT CONTINUED

U S Pat No. 4,378,238

Product of USA

The O. M. Scott & Sons Company Marysville, Ohio 43041

RIGHT GUSSET:

A SCOTTS_R 24612-1 (Scotts (R)) Weed and Feed Lawn (Lawn Pro) Weed Control Plus Prop PRODUĈT (R) Fertilizer

SCOTT'S POLICY No-Quibble Guarantee SEE BACK FOR DETAILS

LEFT GUSSET:

(Scotts (R)) Weed and Feed (Lawn Pro) Weed Control Plus (R) Fertilizer Treats 5,000 Sq.Ft.

SCOTT'S POLICY No-Quibble Guarantee SEE BACK FOR DETAILS

Treats 5,000 Sq.Ft.

A SCOTTS_R 24612-1 Pro_R PRODUCT

PULL TAB

and remove for pour spout

(ILLUS.)

Easy

Open PULL HERE Treats 5,000 sq ft

5M **PROOF OF PURCHASE** (Scotts)

LPNC+LF-24612-1

(Scotts (R)) (Lawn Pro) (R)

Weed and Feed Weed Control Plus Fertilizer

Advantages

- . Safe feeding-won't burn when applied as directed
- . Quick-greening--usually in 4 to 6 days
- . Long-lasting—feeds up to 2 months
- . Dependable—traditional Scotts quality
- . Controls more than 50 weeds...

aster English daisy pigweed bittercress eveningprimrose plantain black medic filaree prickly lettuce bristly oxtongue fleabane puncturevine buckhorn ground ivy purs lane bull thistle healall pusley, Florida burclover henbit ragweed buttonweed horseweed scarlet carpetweed knotweed pimperrel centella (seedling) shadflower chicory lambsquarter shepherdspurse chickweed lippia smartweed cinquefoil mallow sowthistle cudweed moneywort stitchwort curly dock morningglory thymeleaf dandelion mustard sandwort dichondra pennywort vetch dollarweed pepperweed white clover yellow rocket

Directions for Use

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner incons with its labeling.

Po. bc.

APPLICATION

- . Apply any time weeds are actively growing (when daytime temperatures are consistently above $60^{\circ}F$).
- . Apply on a calm day.
- . Apply to moist grass. To help particles stick to weed leaves the grass should be thoroughly watered or wet from dew before applying Lawn Pro Weed and Feed.
- . Do not use this product on flowers, shrubs, fruit or vegetable plants or on creeping bentgrass, St. Augustine, lippia and dichendra lawns. On bermudagrass lawns, under certain conditions, some temporary discoloration may result.

WATERING OR RAIN

Do not water your lawn for 24 hours after application. Water will wash the weed control off the weed's leaves and reduce the effectiveness of this product.

This product should not be applied if rain is expected within the next 24 hours.

SPREADER

Be careful when applying this product near shrubs, flower beds or vegetable gardens as contact with weed control may damage these desirable plants. The Scotts PF Drop Spreaders provide excellent controlled application to protect these areas. When using the EasyGreenTM or SpeedyGreenTM Rotary Spreader take special care near desirable plants.

Spreader Settings

	Scotts	Scotts	Scotts
	Drop	EasyGreen ^R	Speedy
	Spreaders	Rotary	Green TM
(treats 5,000 sq ft)	5	26	4

Additional Spreader Settings (Non-Scotts spreader settings are approximate. You may need to adjust for proper coverage.)

Cyclone Rotary (Model B) 3 1/4	
Central Drop (#205) 5 1/2	
Sears Drop (#19810) 3	•
Ortho Drop (#5000) 6	
Greenview Drop (#105) 3	
Prict ion Drop 7 1/2	
(Red Devil)	

. . . .

RIGHT COLUMN

IMPORTANT

WHAT TO EXPECT

Weed stems and leaves will begin to twist and curl in a few days, but a month or more may be required for complete control.

This product will not prevent weed seeds from producing new plants.

EXTREME TEMPERATURES

Avoid applying when temperatures are consistently above 90 degrees.

SEEDING/SODDING

Wait until the new grass has been mowed at least four times before applying Lawn Pro Weed and Feed weed control.

If you are planning to seed, sod, or sprig you should wait at least one month after you have applied Lawn Pro Weed and Feed.

TIPS ON SPREADING WITH SCOTTS SPREADERS

Scotts Drop:

- First apply 2 header strips
- Operate spreader the longest way of the lawn

(ILLUS.)

- Overlap wheels to insure uniform feeding
- Shut off spreader when you have reached the header strips or stopped

(ILLUS.)

 Mow lawn first if grass is tall enough to touch the bottom of spreader

EasyGreenTM Rotary:

- Slide shut-off lever to the right. Application automatically starts and stops when you do
- For complete coverage make each pass about 30 inches from the previous one so

Speedy GreenTM Rotary:

- Shut off spreader anytime you stop
- For complete coverage make each pass about 5 feet from the previous one so
- Avoid applying product in flower beds adjacent to lawns.

Parkey!

Scotts Policy No-Quibble Guarantee

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

Scotts...the lawn people Marysville, Ohio 43041

0 (UPC) (

(c) 1989. The O M Scott & Sons Company. All rights reserved.

(NOTE: The description below will be inverted.)

24612

(Scotts (R)) (Lawn Pro)

Weed and Feed

(R)

Weed Control Plus Lawn Fertilizer

1262R/15-21

· . :

,