	DM51	532-217	Pg.105
(08/05/86 AR	TILIZER #36111 09/16/86 AR 09/18/86 AR	ن ن کو او
(Scotts _(R))		ntrola Fertilizer	
(NOTE: The description above will	be inverted.)		
Treats up to 5,000 sq ft ((See directions for coverage reco	465m ²) mmendations)	36111 ₄	
(photo) "			
(Scotts _(R))	No Gu	TS POLICY D-Quibble Jarantee CK FOR DETAILS	

DISEASE CONTROL & plus Lawn Fertilizer §

ì

Corrects most lawn fungus problems

r

÷

. . . .

•

FRONT:

ACCEPTED

SEP 1 5 1987

Under the Federal Insecticity, Fungicide, and Refentickly Act, as aniended, for the perticide togistered under EPA Reg. No. 538-217

••••	
•	• • • •
	•••••
• ••	
• • • • • • • •	•
۰ ۱	• • •
1.1	

FRONT CONTINUED:

Net Weight 7 5/8 lbs (3.45 kg)

(

KEEP OUT OF REACH GF CHILDREN CAUTION PRECAUTIONARY STATEMENTS

<u>Hazards to Humans and Domestic Animals</u>: CAUTION, causes eye irritation. Do not get in eyes, on skin, or on clothing. In case of contact immediately wash eyes or skin with plenty of water. Get medical attention if irritation persists. May be harmful if swallowed or absorbed through skin. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

(

Pg. 2015

<u>Environmental Hazards</u>: Do not apply directly to water. Do not contaminate, water by cleaning of equipment or disposal of wastes.

STORAGE: Store in clean, dry place. Reseal opened bag by folding top down and securing. *

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard in trash. ,

(iminocarbonotl Iprodione, 3-(: -2,4-dioxo-l-in	thyl, f Dimethyl (1,2-phenylene)bis hioyl)bis(carbamate)]	0.87% <u>97.38%</u>	
		Total 100.00%	
EPA Reg No 538-217 Disease Control p	7 EPA Est 538-OH-1 lus lawn fertilizer ? 23-3-3 Guaranteed Analysis	643XXXX	
0.6% ammoniaca 14.7% urea, meth 7.7% water inso Available phosphon Soluble potash (K Derived from monoa of potash.	hylene ureas nitrogen oluble nitrogen ric acid (P ₂ O ₅)		,
Chlorine, not more Product of USA	e than	2.60%	
The O M Scott & Sc Marysville, Ohio 4 (c) 198<u>8</u>) The O M		/ d.	•••••
•	25." 213 we also we are the first of the first of the	1 1 1 1	

Sease Control

Harmaran Salaran

BEST AVAILABLE COP

N 36111 A

	<u>GUSSETS:</u>	C	C.	
	RIGHT GUSSET		• • •	
	36111	(Scotts)	Disease,Control plus Lawn Fertilizer	,
	LEFT GUSSET		:	
÷	(Scotts)	Disvase Control plus(Lawn)Fertilizer		<u>3610178</u>
		•		
				•
				د د د ۲ د د ۲
				• • • • • • • • • • • • • • • • • • • •
· .		BECY	AVAN ABLE COPY	τιι ι ι ι ι

BESY AVAILABLE COPY

...

... ...

...

•••

;				(Pg. 4ofS
<u>B</u>	<u>ACK</u> :	C		(
E	PULL TAB and remove for pour spout (ILLUS.) asy pen PULL HERE		Treats 5,000 sq ft (465m ²))	DC+LF 36111 § 5M PROOF OF PURCHASE (Scotts)
			(Scotts _(R))	Disease Contro plus Lawn Fert	ol g cilizer }

<u>Prevents or suppresses most lawn-damaging fungus problems</u>, especially those that plague lawns during warm, humid weather. Scotts scientists originally developed this formulation for golf courses by combining a systemic and a contact fungicide to control a variety of disease problems. This product has proven so successful that we are now making it available for home lawns.

Encourages lawn recovery with a light feeding of slow-release fertilizer.

Easy to Apply

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

- Apply to any type of lawn
- Apply 1/2" of water after application to activate the fungicide. Continue to supply 1" of water per week during hot, dry weather.
 - To control <u>leaf spot</u>, <u>melting-out</u>, <u>dollar spot</u>, <u>pink patch (red thread)</u>, <u>brown patch</u>, (and <u>Fusarium patch</u> in California, Oregon, and Washington). Apply at the REGULAR RATE (5,000 sq ft coverage). You may reapply at two to three week intervals if weather conditions DO NOT change and disease activity continues. To control severe disease infestations, apply at the DOUBLE RATE and repeat monthly.
 - To prevent and suppress <u>summer patch diseases</u> (also known as <u>Fusarium</u> <u>blight</u>, <u>Fusarium roseum</u>, <u>frog-eye</u>, and <u>necrotic ring spot</u>). Apply at the DOUBLE RATE (2,500 sq ft coverage). You may reapply at monthly intervals if weather conditions DO NOT change and disease activity continues.

Summer patch symptoms return annually to the same lawn areas when hot, dry weather promotes stress. For best results begin preventive applications in late spring-early summer. Scotts Play(R) Lawn Seed will offer some resistance to these diseases if lawn renovation becomes necessary.

Allow one week between the application of this and any other control

Pc, Sof S

••••

BEST AVAILABLE COPY

C

BACK CONTINUED:

.

Ċ

Application Rate	Scotts Drop Spreaders	Scotts EasyGreenTM Rotary
Regular Rate	4 3/4	25
(covers 5,000 sq ft) Double Rate (covers 2,500 sq ft)	6 1/4	26
TIPS ON SPREADING WITH SCOTTS SPR	EADERS	
Scotts Drop:		
- First apply two header strips		
- Operate spreader the longest way	y of the lawn	
- Overlap wheels to insure complet	te disease control	
 Shut off spreader when you have strips or anytime you stop 	reached the header	
- Mow lawn first if grass is tall bottom of spreader	enough to touch the	
EasyGreen TM Rotary:		
- Slide shut-off lever to the rig stops when you do	ht. Application auto	matically starts and
 Avoid misses by making each pass some overlapping occurs 	s about 30 inches fro	m the previous one so
(ILLUS.)		.US.)
If you have questions you may tal Counselors. Simply dial this to b	kudinectiyawithwone. Térée filmber == 1=800 (In ohio 1=800	f=ScottsuLawn -543-TURE- 1/762-4010)
0.44.0.11	3611	1
Scotts Policy No-Quibble Guara		
	entitled to get your nce of purchase and w	e 0 (UPC) e <u>32247#36</u> ניו
(NOTE: The descr	iption below will be	inverted.)

Disease Incident Lier



08370