

FRONT:

PROTURF FERTILIZER PLUS
DICOT WEED CONTROL IV XXXXX
(F-1)
12/22/86 CR
01/05/87 CR

04/08/87 CR 538-201

(Scotts(R))
ProTurf(R)
BRAND

36-0-0
Fertilizer Plus
Dicot Weed
Control IV

- . provides postemergence dicot weed control
- . provides controlled release nitrogen feeding

Net Weight 25 lbs (11.34 kg)

KEEP OUT OF REACH OF CHILDREN
WARNING
PRECAUTIONARY STATEMENTS

ACCEPTED

MAY 13 1987
Under the Federal Insecticide, Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under
EPA Reg. No. 538 281

accepted range
0.06 - 1.37%
each ai

Hazards to Humans and Domestic Animals: Causes eye irritation. Do not get in eyes, on skin, or on clothing. In case of contact immediately wash eyes or skin with plenty of water. Get medical attention if irritation persists. Harmful if swallowed. Do not contaminate feed or foodstuffs.

Environmental Hazards: Do not apply directly to any body of water. Do not apply when weather conditions favor drift from treated areas.

ACTIVE INGREDIENTS:

2-Methyl-4-chlorophenoxyacetic acid	1.37%	}
2-(2-Methyl-4-chlorophenoxy) propionic acid	1.37%	

INERT INGREDIENTS:	97.26%
Total	100.00%

EPA Reg No 538-XXX

EPA Est 538-OH-1

ProTurf Fertilizer Plus Dicot Weed Control IV

36-0-0

643-XXXX

Guaranteed Analysis

Total nitrogen (N)	36%
24.0% urea, methylene ureas nitrogen	
12.0% water insoluble nitrogen	
Available phosphoric acid (P ₂ O ₅)	0%
Soluble potash (K ₂ O)	0%

Derived from: urea and methylene ureas

Product of USA

ProTurf Division, The O M Scott & Sons Company
 Marysville, Ohio 43041
 (c) 1987. The O M Scott & Sons Company. All rights reserved.

Recommended For Use By Professional Turfgrass Managers

XXXXX

36-0-0
 Fertilizer Plus
 Dicot Weed
 Control IV

GUSSETS:

30F12

RIGHT GUSSET

XXXXX

(Scotts(R))
ProTurf(R)
BRAND

36-0-0
Fertilizer Plus
Dicot Weed
Control IV

LEFT GUSSET

Scotts(R)
ProTurf(R)
BRAND

36-0-0
Fertilizer Plus
Dicot Weed
Control IV

XXXXX

BACK:

4 of 12

36-0-0
Fertilizer Plus
Dicot Weed
Control IV

XXXXXX

(Scotts(R))
ProTurf(R)
BRAND

36-0-0
Fertilizer Plus
Dicot Weed
Control IV

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on golf course putting greens. Fertilizes grass and controls these rosette and vining type weeds:

bindweed	English daisy	plantain
black medic	eveningprimrose	ragweed
buckhorn	fleabane	red sorrel
coast dandelion (catsear)	ground ivy	scarlet pimpernel
chickweed	healall	shepherdspurse
cliquefoil	henbit	threeseeded mercury
cudweed	mallow	white clover
dandelion	moneywort	wild carrot
	pepperweed	yellow rocket

For use on ornamental Turfgrass areas other than putting greens.

Apply to moist foliage at the RECOMMENDED RATE or for better distribution when using a rotary spreader, apply at the HALF RATE with a reduced width of coverage (see spreader settings). Apply when weeds are growing actively. Grass should not be so tall it touches the bottom of the spreader.

Rain, sprinkling, mowing, or heavy traffic within 24 hours after application can knock granules off weeds and may reduce effectiveness. Apply under non-windy conditions, thereby reducing drift to non-target areas, and optimizing particle distribution for good weed control. Retreating may be necessary, but wait at least 3 weeks to observe full results of first application.

Precautions

Avoid application when daytime temperatures are consistently 90°F or above.

On new grass plantings delay application until after the fourth mowing.

Delay planting grass seed until after a heavy rain or watering and not sooner than 1 month after application.

Not for use on creeping bentgrass; St. Augustinegrass, carpetgrass, dichondra, ground covers, garden plants, flowers and shrubs.

Do not apply to centipedegrass turf in the Spring before March 15 (or before the danger of frost has passed), or after September. Do not apply to bahiagrass before March 15 or after November 1.

Storage and disposal

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ProTurf Division, The O M Scott & Sons Company
Marysville, Ohio 43041

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application.

This bag treats 6,000 ft² (X/X acre) at RECOMMENDED RATE.
This bag treats 6,000 ft² (X/X acre) at HALF RATE and HALF WIDTH.**

SPREADER	GROUND SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS	
			RECOMMENDED RATE	HALF RATE/ HALF WIDTH**
Scotts Drop Type	3 mph	overlap wheels		Not recommended
Gandy	3 mph	overlap wheels		Not recommended
Scotts Rotaries				
R-7*, R-7X*, PT-10*, PT-11*	3 mph	7 feet**		
R-8 (cone 8)	3 mph	9 feet		
Lely models (see below)	4.5 mph	18 feet**		

* Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide 1/4 closed.

**At the HALF RATE setting, apply at coverage width of 3 1/2 ft. for the R-7, R-7X, PT-10, PT-11 and 4 1/2 ft. for the R-8, and 9 ft. for the Lely rotaries to achieve the overall RECOMMENDED RATE.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.

Metric coverage: 15.73 kg treats 557m² at the RECOMMENDED RATE.

FRONT:

PROTURF FERTILIZER PLUS
DICOT WEED CONTROL IV XXXXX
(F-3)
12/22/86 CR 04/08/87 CR
01/05/87 CR
03/02/87 CR

(Scotts(R))
ProTurf(R)
BRAND

24-3-3
Fertilizer Plus
Dicot Weed
Control IV

ACCEPTED
MAY 1 1987
Under the Federal Insecticide
Fungicide and Rodenticide Act,
as amended by the Docket
registered under EPA Reg. No. 538-221

- . provides postemergence dicot weed control
- . provides controlled release nitrogen feeding

accepted range
0.66 - 1.37%
sachai

Net Weight 26 lbs (11.79 kg)

KEEP OUT OF REACH OF CHILDREN
WARNING
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: Causes eye irritation. Do not get in eyes, on skin, or on clothing. In case of contact immediately wash eyes or skin with plenty of water. Get medical attention if irritation persists. Harmful if swallowed. Do not contaminate feed or foodstuffs. ✓

Environmental Hazards: Do not apply directly to any body of water. Do not apply when weather conditions favor drift from treated areas. ✓

ACTIVE INGREDIENTS:

2-Methyl-4-chlorophenoxyacetic acid	0.66%	✓
2-(2-Methyl-4-chlorophenoxy) propionic acid	0.66%	✓
INERT INGREDIENTS:	96.68%	
Total	100.00%	

8 of 10

FRONT CONTINUED:

EPA Reg No 538-XXX

EPA Est 538-OH-1

ProTurf Fertilizer Plus Dicot Weed Control IV

24-3-3

643-XXXX

Guaranteed Analysis

Total nitrogen (N)	24%
0.6% ammoniacal nitrogen	
15.7% urea, methylene ureas nitrogen	
7.7% water insoluble nitrogen	
Available phosphoric acid (P ₂ O ₅)	3%
Soluble potash (K ₂ O)	3%
Chlorine, not more than	2.59%

Derived from: monoammonium phosphate, urea, methylene ureas and muriate of potash.

Product of USA

ProTurf Division, The O M Scott & Sons Company

Marysville, Ohio 43041

(c) 1987. The O M Scott & Sons Company. All rights reserved.

Recommended For Use By Professional Turfgrass Managers

XXXXX

24-3-3
Fertilizer Plus
Dicot Weed
Control IV

GUSSETS:

9 of 12

RIGHT GUSSET

XXXXX

(Scotts(R))
ProTurf(R)
BRAND

24-3-3
Fertilizer Plus
Dicot Weed
Control IV

LEFT GUSSET

Scotts(R)
ProTurf(R)
BRAND

24-3-3
Fertilizer Plus
Dicot Weed
Control IV

XXXXX

BACK:

24-3-3
Fertilizer Plus
Dicot Weed
Control IV

XXXXX

(Scotts(R))
ProTurf(R)
BRAND

24-3-3
Fertilizer Plus
Dicot Weed
Control IV

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on golf course putting greens. Fertilizes grass and controls these rosette and vining type weeds:

- | | | |
|---------------------------|-----------------|---------------------|
| bindweed | English daisy | plantain |
| black medic | eveningprimrose | ragweed |
| buckhorn | fleabane | red sorrel |
| coast dandelion (catsear) | ground ivy | scarlet pimpernel |
| chickweed | healall | shepherdspurse |
| ciquefoil | henbit | threeseeded mercury |
| cudweed | mallow | white clover |
| dandelion | moneywort | wild carrot |
| | pepperweed | yellow rocket |

For use on ornamental Turfgrass areas other than putting greens.

Apply to moist foliage at the RECOMMENDED RATE or for better distribution when using a rotary spreader, apply at the HALF RATE with a reduced width of coverage (see spreader settings). Apply when weeds are growing actively. Grass should not be so tall it touches the bottom of the spreader.

Rain, sprinkling, mowing, or heavy traffic within 24 hours after application can knock granules off weeds and may reduce effectiveness. Apply under non-windy conditions, thereby reducing drift to non-target areas, and optimizing particle distribution for good weed control. Retreating may be necessary, but wait at least 3 weeks to observe full results of first application.

BACK CONTINUED:

11/2/82

Precautions

Avoid application when daytime temperatures are consistently 90°F or above.

On new grass plantings delay application until after the fourth mowing.

Delay planting grass seed until after a heavy rain or watering and not sooner than 1 month after application. ✓

Not for use on creeping bentgrass, St. Augustinegrass, carpetgrass, dichondra, ground covers, garden plants, flowers and shrubs.

Do not apply to centipedegrass turf in the Spring before March 15 (or before the danger of frost has passed), or after September. Do not apply to bahiagrass before March 15 or after November 1.

Storage and disposal

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing. ✓

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. ✓

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke. ✓

ProTurf Division, The O M Scott & Sons Company
Marysville, Ohio 43041

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application.

This bag treats 5,000 ft² (465 m²) at the RECOMMENDED RATE.
 This bag treats 5,000 ft² (465 m²) at the HALF RATE and HALF WIDTH.**

<u>SPREADER</u>	<u>GROUND SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>SPREADER SETTINGS</u>	
			<u>RECOMMENDED RATE</u>	<u>HALF RATE/ HALF WIDTH**</u>
Scotts Drop Type	3 mph	overlap wheels		Not recommended
Gandy	3 mph	overlap wheels		Not recommended
Scotts Rotaries				
R-7*, R-7X*, PT-10*, PT-11*	3 mph	7 feet**		
R-8 (cone 8)	3 mph	9 feet		
Lely models (see below)	4.5 mph	18 feet**		

* Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide 1/4 closed.

**At the HALF RATE setting, apply at coverage width of 3 1/2 ft. for the R-7, R-7X, PT-10, PT-11 and 4 1/2 ft. for the R-8, and 9 ft. for the Lely rotaries to achieve the overall RECOMMENDED RATE.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.

Metric coverage: 11.79 kg treats 465m² at the RECOMMENDED RATE.