

25

538-227

4/28/88

(S-1647R - 1.5 lb rate)

PROTURF FERTILIZER PLUS

WEED LOSS CONTROL XXXXX

07/16/85 JF 08/27/85 JF

08/09/85 JF 09/11/85 JF

P7/f12

**ACCEPTED**

APR 28 1988

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 538-227

FRONT:

(Scotts(R))  
ProTurf(R)  
BRAND

High Density  
26-3-8  
Fertilizer Plus Weedgrass Control

- o preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge and Poa annua
- o provides controlled-release nitrogen feeding

Net Weight 9 3/16 lbs (4.16 kg)

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. If in eyes: Flush with plenty of water. Get medical attention if irritation persists. Wash thoroughly with soap and water after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In case of spills, either collect for use or dispose of properly.

ACTIVE INGREDIENT:

pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine*	4.13%
INERT INGREDIENTS:	95.87%
	Total 100.00%

EPA Reg. No. 538-227

EPA Est 538-OH-1

Fertilizer Plus Weedgrass Control 26-3-8

Guaranteed Analysis

Total nitrogen (N) . . . . .	) . . . . .	26%	2/12
0.6% ammoniacal nitrogen			
19.7% urea, methylene ureas nitrogen			
5.7% water insoluble nitrogen			
Available phosphoric acid (P <sub>2</sub> O <sub>5</sub> ) . . . . .		3%	
Soluble potash (K <sub>2</sub> O) . . . . .		8%	

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA

U.S. Pat Nos 3,705,794; 3,989,470

\*Licensed under one or more of the following

U.S. Patents: 3,920,742; 4,066,441

ProTurf Division, The O M Scott & Sons Company

Marysville, Ohio 43041

(c) 1985 The O M Scott & Sons Company. All rights reserved.

---

Recommended for Use by Professional Turfgrass Managers

---

XXXXX

26-3-8  
Fertilizer Plus  
Weedgrass  
Control

GUSSETS:

3/12

RIGHT GUSSET

XXXXX

(Scotts(R))  
Proturf(R)  
BRAND

26-3-8  
Fertilizer Plus  
Weedgrass  
Control

LEFT GUSSET

Scotts(R)  
ProTurf(R)  
BRAND

26-3-8  
Fertilizer Plus  
Weedgrass  
Control

XXXXX

BACK:

26-3-8  
Fertilizer Plus  
Weedgrass Control

(Scotts(R))  
ProTurf(R)  
BRAND

26-3-8  
Fertilizer Plus  
Weedgrass Control

4/12

---

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on bentgrass or Poa annua where it is the desired species or on dichondra.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bermudagrass (non-winter overseeded), St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass Precautions."

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

Grassy Weeds

crabgrass  
foxtail  
goosegrass\*  
Poa annua  
barnyardgrass  
fall panicum

Dicot Weeds

chickweed  
oxalis  
prostrate spurge\*

To prevent crabgrass, foxtail, barnyardgrass, fall panicum, chickweed, oxalis and spurge, apply product to dry foliage at the RECOMMENDED RATE in the spring before they sprout.

\*To prevent goosegrass and heavy infestations of spurge, repeat application at the RECOMMENDED RATE after 6-8 weeks. Apply to established, dense turf.

To prevent Poa annua and fall germinating chickweed, apply product to dry foliage in late summer-early fall (before germination) at the RECOMMENDED RATE. Repeat application yearly for continued control.

For best results, water after all applications (1/4-1/2 inch) to remove the granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying after application.

### Precautions

Do not use on bentgrass and Poa annua where it is the desired species or on dichondra.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, and excessive soil moisture).

### Bermudagrass Precautions

Do not apply to dormant bermudagrass (turf that is not actively growing or has experienced frost). However, an application can be made if the bermudagrass is maintained at a high cut (1/2 inch or more) on medium-fine textured soils such as clays and clay loams.

Where turf thinning or damage has occurred due to winter injury of bermudagrass, wait until greenup, fill-in and good rooting before making an application.

Application to overseeded bermudagrass may injure and thin the cool season variety.

### Other Precautions

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Applications should not exceed twice yearly.

Avoid applications to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water.

### STORAGE AND DISPOSAL

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Suggested Spreader Settings

to provide proper distribution calibrate spreader before application

9 3/16 lbs (4.16 kg) treats 11,000 ft<sup>2</sup> (1/4 acre/1022 m<sup>2</sup>) at recommended rate.

<u>SPREADER</u>	<u>GROUND OR PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>RECOMMENDED RATE</u>	<u>HALF RATE with 1/2 SWATH WIDTH**</u>
-----------------	----------------------------	--------------------------	-------------------------	---

Scott Drop Type  
 Gandy (30")  
 Scott Rotaries\*  
     R-7, R-7X  
     R-8 (cone X)  
 Lely Models

\*Set pattern slide to 1/4 closed position.

\*\*At the HALF RATE setting, apply at a swath width of XXXXX ft. for the Scott Rotary and X ft. for the Lely to achieve the overall RECOMMENDED RATE.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.

06790/1529X

(S-1525TR - 1.5 lb rate)  
PROTURF FERTILIZER PLUS  
WEED CONTROL XXXXX  
07/16/85 JF 08/28/85 JF  
08/08/85 JF 09/11/85 JF

7/12

ACCEPTED  
APR 28 1988  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 538-227

FRONT:

(Scotts(R))  
ProTurf(R)  
BRAND

High Density  
30-3-10  
Fertilizer Plus Weedgrass Control

- o preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge and Poa annua
- o provides controlled-release nitrogen feeding

Net Weight 66 lbs (29.94 kg)

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. If in eyes: Flush with plenty of water. Get medical attention if irritation persists. Wash thoroughly with soap and water after handling. May be harmful if swallowed. Avoid inhalation. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In case of spills, either collect for use or dispose of properly.

ACTIVE INGREDIENT:  
pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl  
-2,6-dinitrobenzenamine\* . . . . . 0.57%

INERT INGREDIENTS: . . . . . 99.43%  
Total 100.00%

EPA Reg. No. 538-227 EPA Est 538-OH-1

Fertilizer Plus Weedgrass Control 30-3-10 643-XXXX

Guaranteed Analysis

8/12

Total nitrogen (N) . . . . . 30%  
 0.6% ammoniacal nitrogen  
 23.0% urea, methylene ureas nitrogen  
 6.4% water insoluble nitrogen  
 Available phosphoric acid (P<sub>2</sub>O<sub>5</sub>) . . . . . 3%  
 Soluble potash (K<sub>2</sub>O) . . . . . 10%

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA

U.S. Pat Nos 3,705,794; 3,989,470

\*Licensed under one or more of the following

U.S. Patents: 3,920,742; 4,066,441

ProTurf Division, The O M Scott & Sons Company

Marysville, Ohio 43041

(c) 1985 The O M Scott & Sons Company. All rights reserved.

---

Recommended for Use by Professional Turfgrass Managers

---

XXXXX

30-3-10  
Fertilizer Plus  
Weedgrass  
Control



GUSSETS:

9/12

RIGHT GUSSET

XXXXX

(Scotts(R))  
Proturf(R)  
BRAND

30-3-10  
Fertilizer Plus  
Weedgrass  
Control

LEFT GUSSET

Scotts(R)  
ProTurf(R)  
BRAND

30-3-10  
Fertilizer Plus  
Weedgrass  
Control

XXXXX

BACK:

30-3-10  
Fertilizer Plus  
Weedgrass Control

(Scotts(R))  
ProTurf(R)  
BRAND

30-3-10  
Fertilizer Plus  
Weedgrass Control

10/12

---

**Directions for use**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on bentgrass or Poa annua where it is the desired species or on dichondra.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bermudagrass (non-winter overseeded), St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass Precautions."

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

Grassy Weeds

crabgrass  
foxtail  
goosegrass\*  
Poa annua  
barnyardgrass  
fall panicum

Dicot Weeds

chickweed  
oxalis  
prostrate spurge\*

To prevent crabgrass, foxtail, barnyardgrass, fall panicum, chickweed, oxalis and spurge, apply product to dry foliage at the RECOMMENDED RATE in the spring before they sprout.

\*To prevent goosegrass and heavy infestations of spurge, repeat application at the RECOMMENDED RATE after 6-8 weeks. Apply to established, dense turf.

To prevent Poa annua and fall germinating chickweed, apply product to dry foliage in late summer-early fall (before germination) at the RECOMMENDED RATE. Repeat application yearly for continued control.

For best results, water after all applications (1/4-1/2 inch) to remove the granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying after application.

11/12

Precautions

Do not use on bentgrass and Poa annua where it is the desired species or on dichondra.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature - below 40°F, and excessive soil moisture).

Bermudagrass Precautions

Do not apply to dormant bermudagrass (turf that is not actively growing or has experienced frost). However, an application can be made if the bermudagrass is maintained at a high cut (1/2 inch or more) on medium-fine textured soils such as clays and clay loams.

Where turf thinning or damage has occurred due to winter injury of bermudagrass, wait until greenup, fill-in and good rooting before making an application.

Application to overseeded bermudagrass may injure and thin the cool season variety.

Other Precautions

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly planted areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Applications should not exceed twice yearly.

Avoid applications to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water.

STORAGE AND DISPOSAL

**STORAGE:** Store in a clean, dry place. Reseal opened bag by folding top down and securing.

**PESTICIDE DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Suggested Spreader Settings

to provide proper distribution calibrate spreader before application

66 lbs (29.94 kg) treats 11,000 ft<sup>2</sup> (1/4 acre/1022 M<sup>2</sup>) at recommended rate.

<u>SPREADER</u>	<u>GROUND OR PTO SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>RECOMMENDED RATE</u>	<u>HALF RATE with 1/2 SWATH WIDTH**</u>
-----------------	--------------------------------	------------------------------	-----------------------------	---

Scott Drop Type

Gandy (30")

Scott Rotaries\*

R-7, R-7X

R-8 (cone X)

Lely Models

\*Set pattern slide to 1/4 closed position.

\*\*At the HALF RATE setting, apply at a swath width of XXXXX ft. for the Scott Rotary and X ft. for the Lely to achieve the overall RECOMMENDED RATE.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.