

PM 25

5/8/91

538-237

LOW END OF RANGE

97 124

Private Label Fertilizer and Crabgrass Preventer XXXX

04/12/91 VS/cr

FRONT

XXXX

NEW EASY CARRY HANDLE

SCOTTS(R)



Private Label Fertilizer and Crabgrass Preventer

TREATS 5,000 - 6,000 SQUARE FEET

SCOTTS(R)


(RIBBON)
THE NAME TO
GROW WITH!

(Logo)

Premium

Private Label Fertilizer and Crabgrass Preventer

Not for use on dichondra or bentgrass lawns



26-3-3

ACTIVE INGREDIENT:

Pendimethalin, N-(1-Ethylpropyl)-3,4-dimethyl-2,6-dinitrobenzenamine* 0.50%

INERT INGREDIENTS: 99.50%

TOTAL 100.00%

* Licensed under one or more of the following U.S. patents: 3,920,742; 4,066,441.

EPA Reg. No. 538-237

EPA Est. XXX-XX-X

CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Net Weight 34 5/8 LBS.

SCOTTS(R)

Private Label Fertilizer and Crabgrass Preventer

XXXX

TREATS 5,000 - 6,000 SQ. FT.

XXXX

INSTRUCTIONS

SCOTT'S Private Label Fertilizer and Crabgrass Preventer prevents germination of crabgrass and certain other annual weeds. In addition, a XX-X-X fertilizer provides a supply of plant food needed for growth.

1. Apply in early spring before the 3rd or 4th mowing or before temperatures are regularly in the 80's. Rain, snow or freezing weather will not affect the performance of this product.
2. Rake, then mow lawn to normal height 1 to 2 days before application. Important: Do not rake or aerify for 4 months after application to avoid disturbing the crabgrass barrier on the soil surface.
3. Apply when lawn is dry. It is not necessary to water it in after application.
4. Apply to prevent germination of crabgrass, foxtail, barnyardgrass, fall panicum, spurge* and oxalis (common yellow woodsorrel):
 - . in early spring (before dandelions reach the puffball stage)
 - . before March in southern California
 - . before mid-March in northern California

* For heavy infestations, a repeat treatment is recommended after 6-8 weeks.
5. Apply in August or September to prevent germination of Poa annua and chickweed.

SPREADER SETTINGS

Always apply fertilizer products with a spreader. Avoid applying this product on flower, vegetable and shrub plantings.

Scotts SpeedyGreen Rotary X	Scotts EasyGreen Rotary X
Scotts Drop X	Sears Drop (#19810) X
Greenview Drop (Spred II). X	Ortho Drop (#5000) X
Central Drop (#205). X	Precision Drop (Red Devil). X

Above spreader settings are approximate. You may need to adjust for proper coverage.

WEEDS PREVENTED

Crabgrass, foxtail, oxalis (common yellow wood sorrel), spurge*, Poa annua, barnyardgrass, chickweed, fall panicum.

*For heavy infestations, a repeat treatment is recommended after 6 to 8 weeks.

STORAGE

Store in clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL

Securely wrap original bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL

Do not reuse bag. Discard bag in trash.

<p style="text-align: center;">UNCONDITIONALLY GUARANTEED</p> <p>If you are not satisfied with the results from using this product, we will refund your purchase price.</p> <p>Send evidence of your purchase to: THE O.M. SCOTTS & SONS COMPANY MARYSVILLE, OHIO 43041</p>

<p style="text-align: center;">Proof of Purchase UPC</p>

(Logo) Distributed By:
THE O.M. SCOTTS & SONS COMPANY
MARYSVILLE, OHIO 43041

TREATS 5,000 - 6,000 SQ. FT.

(Logo)

SCOTTS(R)

Private Label
Fertilizer and Crabgrass Preventer

Gussets

XXXX

(Logo)

SCOTTS(R)

Private Label
Fertilizer and Crabgrass Preventer

(Logo) SCOTTS(R)

Premium

Private Label
Fertilizer and Crabgrass Preventer

26-3-3

GUARANTEED ANALYSIS

Total Nitrogen (N)	26%
0.7% ammoniacal nitrogen	
24.0% urea, methylene ureas nitrogen**	
1.3% water insoluble nitrogen	
Available phosphoric acid(P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	3%

Derived from: monoammonium phosphate, urea, methylene ureas, muriate of potash.

Chlorine, not more than 2.38%

**Contains 8.0% slowly available methylenediurea and dimethylenetriurea nitrogen.

Product of USA

**KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS
Hazards to Humans and Domestic Animals**

CAUTION, causes eye irritation. Do not get in eyes. If in eyes, flush with plenty of water and get medical attention if irritation persists. Wash after handling. May be harmful if swallowed. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply to water or wetlands. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. In case of spills, either collect for use or dispose of properly.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE PRECAUTIONS

Do not use this product on dichondra or bentgrass lawns. Do not seed or sod for 4 months after using. The crabgrass barrier prevents grass seed from sprouting. Do not apply to newly seeded or sodded areas until after the fourth mowing. On newly sprigged areas, wait 5 months before application. Don't apply this product more often than twice annually. Except for heavy infestations of spurge, allow at least 4 months between applications. Do not apply material to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse off with water. This product does not control actively growing crabgrass and other existing weeds.