


FRONT:

STOCK #8549  
03/22/89 SEY 7/17/89 SC  
11/07/91 VS/cr

06/29/93 VS/cr  
01/28/94 VS/cr  
05/17/94 VS/cr

(Scotts(R))  
ProTurf(R)  
BRAND



Fungicide IX

o provides protection from and control of gray snow mold, Pythium blight, dollar spot, brown patch, copper spot and pink snow mold

Net Weight 42.24 lbs (19.15 kg)

KEEP OUT OF REACH OF CHILDREN  
CAUTION  
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling.

**Statement of Practical Treatment**

If on skin: Wash with plenty of soap and water. Get medical attention.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

Environmental Hazards: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

**BEST AVAILABLE COPY**

ACTIVE INGREDIENTS:

Chloroneb, (1,4-Dichloro-2,5-dimethoxybenzene . . . . .	3.26%
Thiophanate-methyl, Dimethyl 4,4'(-o-phenylenebis) (3-thioallophanate) . . . . .	1.63%

INERT INGREDIENTS: . . . . .	95.11%
Total	100.00%

U.S. Patent No. 4,028,464

EPA Reg No 538-242

EPA Est 538-OH-1 (S)

EPA Est 9198-OH-1 (M)

Subscript used in first letter of  
run code on end of container.

Product of U.S.A.

Distributed by:  
ProTurf Division  
The O M Scott & Sons Company  
14111 Scottslawn Road  
Marysville, Ohio 43041

(c) 1994. The O M Scott & Sons Company

All rights reserved

---

Recommended For Use By Professional Turfgrass Managers

---

8549

Fungicide IX

**BEST AVAILABLE COPY**

RIGHT GUSSET

8549

(Scotts(R))  
ProTurf(R)

Fungicide IX

LEFT GUSSET

(Scotts(R))  
ProTurf(R)

Fungicide IX

8549 (Packaging Calendar)

**BEST AVAILABLE COPY**

Fungicide IX

(Scotts (P))  
ProTurf (R)

Fungicide IX

Directions for use

---

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Adapted for use in all areas where listed turf diseases attack turfgrasses.

TO PREVENT GRAY AND PINK SNOW MOLD, apply to dry foliage at the NORMAL or MEDIUM RATE prior to the first snow. Repeat in mid-Winter when turf is free of snow and again immediately after the final snow melts in the Spring. In areas where snow cover is present the entire Winter, apply at the HIGH RATE prior to the first snow and follow with a NORMAL RATE when the snow melts in the Spring, if needed.

TO PREVENT PYTHIUM BLIGHT (when weather conditions are favorable: high temperature and humidity, day and night), apply to moist foliage at the NORMAL or MEDIUM RATE and repeat at 5-7 day intervals while weather conditions persist.

TO CONTROL ACTIVE PYTHIUM BLIGHT, apply to moist foliage at the MEDIUM RATE as soon as symptoms are observed and repeat as needed if weather conditions remain favorable. For severe attack, apply a HIGH RATE application to moist foliage and repeat at NORMAL RATE as needed. If disease symptoms do not subside, use a second HIGH RATE application in 5-7 days.

TO PREVENT DOLLAR SPOT\*, COPPER SPOT AND BROWN PATCH, apply to dry or moist foliage at the NORMAL or MEDIUM RATE when disease activity is noted or suspected, repeat at 2-3 week intervals. During periods of high day and night temperatures and humidity, treatment intervals may need to be shortened for best brown patch control.

TO CURE DOLLAR SPOT, COPPER SPOT AND BROWN PATCH, when symptoms indicate advanced stages, apply at the MEDIUM or HIGH RATE and follow with the NORMAL RATE preventive program.

\*If dollar spot control is not effective after using this product as recommended, a thiophanate-methyl or benomyl-tolerant strain of fungi may be present. Consult your ProTurf Tech Rep for recommendations of suitable ProTurf Fungicides to use promptly.

ON NEW SEEDINGS, apply at MEDIUM RATE just after sowing seed. After germination or at first sign of disease, apply to moist foliage and repeat at 7-10 day intervals if evidence of disease persists.

DRY FOLIAGE APPLICATIONS should be made to prevent snow mold. For prevention or control of Pythium blight, prolonged contact with the foliage enhances activity.

ON GREENS AND FAIRWAYS, it is suggested to apply the product to moist foliage in the morning and evening, followed by a light syringing before mowing or allowing heavy play.

Do not apply in a brisk wind.

#### PRECAUTIONS

Allow one week between the application of this and other control products. (This guideline can be relaxed where severe insect or disease attack requires immediate treatment).

When treating Pythium and brown patch for control on established turf, do not fertilize or water heavily during periods when disease is active.

#### Storage and disposal

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Distributed by:  
ProTurf Division  
The O M Scott & Sons Company  
14111 Scottlawn Road  
Marysville, Ohio 43041

### Suggested Spreader Settings

To provide proper distribution calibrate spreader before application.

42.24 lbs. treats 22,000 sq ft (1/2 acre) at the NORMAL RATE  
 42.24 lbs. treats 11,000 sq ft (1/4 acre) at the MEDIUM RATE  
 42.24 lbs. treats 7,333 sq ft (1/6 acre) at the HIGH RATE

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS		
			NORMAL RATE	MEDIUM RATE	HIGH RATE
Scotts Drop Type	3 mph	overlap wheels	5 3/4	7 1/2	9
Gandy (30")	3 mph	overlap wheels	22	28	33
Scotts Rotaries R-7, R-7X*, PT-10, PT-11	3 mph	8 ft.	E 1/2	I	L
R-8A (cone 8)	3 mph	9 ft.	K	N 1/2	Q 1/2
Lely Models (see below)	4.5 mph	16 ft.	4 II	5 1/4 II	6 II
Vicon (Model 02)	4.5 mph	18 ft.	20	30	37
Vicon (Model 03)	4.5 mph	10 ft.	16	22	27

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.

\* 1/4 slide

PRODUCT RATE: NORMAL = 1.92 lbs. product/1,000 sq ft (1.0 + 0.5 oz ai/M)  
 MEDIUM = 3.84 lbs. product/1,000 sq ft (2.0 + 1.0 oz ai/A)  
 HIGH = 5.76 lbs. product/1,000 sq ft (3.0 + 1.5 oz ai/A)

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY

This product has been researched to provide necessary data to support its uses listed on the label. The user should always follow the label directions and exercise his judgement and caution when using the product under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS, OR IMPLIED, CONCERNING THE RESULTS TO BE OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. The exclusive remedy of the user or Buyer, and the limit of liability of The O M Scott & Sons Company or any other Seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The O M SCott & Sons Company.

Distributed by:  
ProTurf Division  
The O M SCott & Sons Company  
14111 Scottslawn Road  
Marysville, Ohio 43041