


FRONT LABEL

Liq Plant Disease Control
#XXXX
7/26/94 ER

Ready to use...
Simply attach to a garden hose
and spray!



Apple Scab
Illus.

Black Spot
Illus.

Powdery
Mildew
Illus.

Plant Disease Control

The Scotts Difference™

- Controls apple scab, black spot, leaf spot, powdery mildew, and other plant diseases listed on back
- Also controls summer patch, dollar spot, stripe smut, and other lawn diseases listed on back
- Immediate contact action PLUS long-lasting systemic control
- No messy mixing or measuring required
- Scotts No-Quibble Guarantee

ACTIVE INGREDIENT:	
Thiophanate-methyl (dimethyl(1,2-phenylene)-bis (iminocarbonothioyl))bis(carbamate).....	3.85%
INERT INGREDIENTS:	96.11%
TOTAL	100.00%

EPA Reg. No. 4581-352-XXX
EPA Est. No. XXXXX-XX-XX

CAUTION: KEEP OUT OF REACH OF CHILDREN
See back label for additional precautionary statements and statement of practical treatment.
Treats 5 to 20 trees (10-ft. tall) or 15 to 30 shrubs (4-ft. tall).

Net Contents: 1 qt. (32 fl.oz.)(946 ml)

BEST AVAILABLE COPY

274

OUTSIDE BACK LABEL

Liq Plant Disease Control
#XXXX
7/26/94 ER

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals:

Harmful if swallowed, absorbed through skin, or inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Statement of Practical Treatment:

IF SWALLOWED: Induce vomiting immediately by giving two glasses of water and sticking finger down throat. Call a physician. Never give anything by mouth to an unconscious person.
IF IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention if irritation persists.
IF ON SKIN: Immediately flush skin with plenty of water.
IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration. Get medical attention.

Environmental Hazards:

This pesticide is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not contaminate water by disposal of equipment washwaters.

Storage and disposal

Do not contaminate water, food or feed by storage or disposal.
STORAGE: Store in the original container in a dry area. Do not store in a manner where cross-contamination with other pesticides, fertilizers, food or feed could occur. If spilled during storage or handling, sweep up spillage and dispose of according to the Pesticide Disposal Instructions listed below.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: After bottle has been emptied, dispose of empty bottle in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Product of USA
Distributed by
The O M. Scott & Sons Company
14111 Scottslawn Rd.
Marysville, OH 43041

Complete directions for use inside.

Questions?

Call Scotts'

toll-free

Consumer


Hotline:

1-800-543-TURF

P
U
L
L
H
E
R
E
T
O
R
E
S
E
A
L
O
P
E
N

Scotts No-Quibble Guarantee
If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

...the lawn and garden people (SM)
Marysville, Ohio 43041




0 32247 0XXXX X
Plant Disease Control XXXXS
PROOF OF PURCHASE

1st INSIDE BACK LABEL

Liq Plant Disease Control

#XXXX

7/26/94 ER



Directions for use

Product Control Button

Turn red knob clockwise until slot in knob aligns with white tab. Push in to ALLOW product flow. When finished, push red button in from opposite side to STOP product flow. Turn red knob counterclockwise to prevent accidental discharge of product.

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

SHAKE WELL BEFORE USE

Easy to use

1. Make sure the ON/OFF Water Control Knob is in the OFF position.
2. Shake well before applying.
3. Connect spray nozzle to hose with convenient swivel end.
4. Turn the red knob on the Product Control Button to align with the white tab, then push the button in all the way. Avoid squeezing bottle after this, as contents may spray through nozzle opening.
5. Turn water on at faucet.
6. When ready to spray, point nozzle toward lawn, then turn Water Control Knob to ON position.
7. When finished, push red Product Control Button to OFF position to stop product flow. Turn Water Control Knob to OFF and turn off faucet. Turn Water Control Knob to ON to release water pressure, then turn OFF again.
8. To prevent accidental discharge when not in use, turn red knob so it is not aligned with white tab and cannot be pushed in. Store out of reach of children.

How to spray

Shake bottle often while spraying to keep contents from settling.

Apply until leaves glisten and runoff begins to occur. This bottle will treat 2,450 to 4,900 sq. ft. of area depending on the tree or shrub size and the density of the leaf canopy.

BEST AVAILABLE COPY

2nd INSIDE BACK LABEL

Liq Plant Disease Control
#XXXX
7/26/94 ER

REVISED

DISEASE CONTROLLED	TYPE OF PLANT THE DISEASE ATTACKS			WHEN TO APPLY
	Woody *	Herbaceous **	Specific Plants	
Black Spot			Roses	Begin application when disease first appears and repeat at 10- to 14-day intervals during the growing season. Shorten the intervals during humid, rainy weather.
Ascochyta Blight		X		
Botrytis (gray mold)	X	X		
Diplodia Tip Blight (diplodia die)			Shade & ornamental trees	
Leaf Spot, caracopora	X	X		
Leaf Spot, corynespora			Ligustrum	
Leaf Spot, didymella			Iris	
Leaf Spot, entomosporium	X	X		
Leaf Spot, phomaeae	X	X		
Leaf Spot, ramularia		X		
Leaf Spot, seporia	X	X		
Powdery Mildew	X	X	Ornamental nut & fruit trees	
Anthracoese	X		Shade trees	
Scale			Pyracantha, Flowering crab, ornamental fruit & nut trees	
Ovulinia			Azalea, madodendron	

* Woody plants have strong stems, such as shrubs or trees.
** Herbaceous plants are mostly foliage (leaves) such as perennials or bedding plants.

For Lawns

Use on bentgrass, bluegrass, bermudagrass, fescues, ryegrass, St. Augustinegrass, or mixtures of these grasses for the prevention and control of the following diseases:

• **FUSARIUM BLIGHT, SUMMER PATCH and NECROTIC RING SPOT**
Apply this bottle to 225 to 450 sq. ft. Apply 1 inch of water immediately after application. Make two applications 10 to 14 days apart, beginning when disease first appears.

• **DOLLAR SPOT and ANTHRACNOSE**
Apply this bottle to 1,250 to 2,500 sq.ft. beginning when disease first appears. Do not water in. Repeat every 10 to 14 days through the season as needed.

• **LARGE BROWN PATCH, COPPER SPOT, RED THREAD and FUSARIUM PATCH:**
Apply this bottle to 1,250 to 2,500 sq.ft. beginning when the disease first appears and continue every 10 to 14 days through the season as needed. Do not water in. When conditions are unusually favorable for development of disease, reduce intervals to 5 to 7 days.

• **STRIFE SMUT**
Apply this bottle to 225 to 450 sq.ft. Apply 1 inch of water immediately after application. Make two applications at 14 to 21 day intervals. Begin applications when the disease first appears, either in the spring or fall.

BEST AVAILABLE COPY