

U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (H7505C) 401 "M" St., S.W. Washington, D.C. 20460

EPA Reg. Number:

Date of Issuance:

538-254

045 JUN 1995

Tei of Issuance: Conditional

NOTICE OF PESTICIDE:

<u>x</u> Registration Reregistration

(under FIFRA, as amended)

Name or Laucide Product:

Fertilizer Plus Diazinon

Name and Address of Registrant (include ZIP Code):

The Scotts Company 14111 Scottslawn Road Marysville, OH 43041

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number

On the basis of information furnished by the registrant, the above named posticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

- 1. Submit and/or cite all data required for registration/ reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
- 2. Make the labeling changes listed below before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 538-254".
 - b. In the Environmental Hazards, add a comma after "Birds, especially waterfowl". Also, in this same section, revise "0.25 inch" to "1/4 inch" for consumer clarity.
- 3. Submit three (3) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

yikusmic	or	Approving	Official

Date:

EPA Form 8570-6

page 2 EPA Reg. No. 538-254

4. You have agreed in writing that The Scotts Company will make any changes at the next printing of the label which the Agency should deem necessary once it has completed review of the data submitted in support of this registration.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosures

MASTER LABEL

FRONT:

PACKAGE COPY FOR FERTILIZER PLUS DIAZINON

03/22/95 VS/kl

Fertilizer Plus Diazinono

(Scotts ®) (Lawn Pro ®)

stepTM3

Best

Annual

Program

3105-3/39065-4 (placed vertically)

(Above copy to be inverted.)

Treats 5,000 sq. ft (Bag)

(Scotts®)
(Lawn Pro®)

"Available only at authorized Scotts® Lawn Pro® dealers"

Scotts® Best Annual Program SCOTTS POLICY No-Quibble

Guarantee

SEE BACK FOR DETAILS

stepTM3

of 4 easy steps to a better lawn

(ILLUS.)

Corrects most lawn insect problems

OPTIONAL FRONT PANEL CLAIMS

---The Scotts Difference®---

- Contains diazinon to control white grubs, sod webworms, fleas, Brown dog ticks, chinch bugs, ants, European crane fly larvae, and other insects listed on back
- Timed-release nitrogen won't burn your lawn or cause surge growth
- Scotts No-Quibble Guarantee
- Toll-free Consumer Hotline: 1-800-543-TURF

For outdoor use on the home lawn only. This product must 197 be used on golf courses and sod farms.

Patented Particle Technology

[Art]

A JUN 1995

Under the Federal lower lide.

Fundide, and Rodenti and amended, for the posterior registered under IPA, hep. No.

EPA Letter Diteil

OPTIONAL FRONT PANEL CLAIMS (continued)

Unlike other insect control products, every particle of Fertilizer Plus Diazinon contains

- Scotts patented timed-release nitrogen
- Phosphorus
- Potassium
- Insect control

Treats and fertilizes 5,000 sq. ft.

Kills chinch bugs

Kills sod webworms

Kills white grubs

NET WEIGHT 12.5 LBS. (5.68 kg)

Made in U.S.A.

32-3-3

CAUTION KEEP OUT OF REACH OF CHILDREN


See BACK for additional precautionary statements.

ACTIVE INGREDIENT:

Diazinon [0,0-Diethyl 0-(2-iso	propyl-6-meth	ıyl-4
pyrimidinyl) phosphorothioate		
NÉRT INGRÉDIENTS		
	Total	100.00%

EPA Reg No 538-ELU

EPA Est 538-OH-l



[RIGHT GUSSET]

3105 3

[UPC] 0 32247 03105 9

(LawnPro®) Treats and fertilizes 5,000 sq. ft. (Scotts®) Fertilizer Plus Diazinon

Best Annual Program

stepTM 3

[LEFT GUSSET]

(Scotts®)

(LawnPro®) Treats and fertilizes 5,000 sq. ft. (Scotts®) Fertilizer Plus Diazinon

[UPC] 0 32247 03105 9

3105

Best Annual Program

[BACK]

PULL TAB and remove for pour spout

Easy Open [ILLUS.] PULL HERE

Treats and fertilizes 5,000 sq. ft.

KEEP OUT OF REACH OF CHILDREN CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION. Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin or clothing. Wash thoroughly after handling and before smoking or eating. Do not store near food or feed. Do not contaminate ornamental fish ponds. Do not allow children or pets onto the treated area until granules have been watered in and the grass or soil has dried.

Statement of Practical Treatment: If swallowed: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing. If inhaled: Remove victim to fresh air and apply artificial respiration if indicated. If on skin: Remove contaminated clothing and wash affected areas with soap and water. If in eyes: Rinse eyes with plenty of water and call a physician immediately.

Note to Physician: This product contains an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Environmental Hazards: This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed unaximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping granules. Where irrigation (watering) is recommended in the directions or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), apply a minimum of 0.25 inch of water immediately after applying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

Fertilizer Plus Diazinon®

643-xxxx

Totalizor Tido Didziliono	0.0	
Guaranteed Analysis	222	
Total nitrogen (N)	32%	
0.4% ammoniacal nitrogen		
31.6% urea, methylene urea, nitrogen* Available phosphate acid (P ₂ O ₅) Soluble potash (K ₂ O)	, , , ,	•
Available phosphate acid (P ₂ O ₅)	2%	
Soluble potash (K ₂ O)	3%	, , , , ,
(2)	• •	, ,
Derived from: monoammonium phosphate, urea, methylene ureas and potassi		• • •
*Contains xx% slowly available methylenediurea and dimethylenetriurea nitro	gen.) 1 P
(Scotts®) Fertilizer Plus Diazinon BACK Cont'd		

STORAGE: Store in original container in a cool, dry, well-ventilated area inaccessible to children, and domestic animals. Do not store near fertilizers, herbicides, seeds, food, or feed. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

(OPTIONAL BACK PANEL CLAIMS)

Now it's easy for you to have a better lawn

Save money

Why even consider a lawn care service? Scotts Annual Program will give you a beautiful lawn for about one-third the cost.

Simple and easy

No more guesswork. You simply apply the right product - one Step at a time.

Start anytime

You can start into the Annual Program anytime of the year. Just check the month and select the appropriate Step listed below.

Guaranteed results

Each Step is designed to the specific season to help give you a greener lawn.

[The 4 Steps listed below will be placed horizontally in one row.]

step™1. Early Spring February-April

Crabgrass Preventer plus Fertilizer - prevents crabgrass, foxtail, spurge, oxalis, chickweed and other annual weeds while providing deep, thick greening for up to 2 months. In the Pacific Northwest, StepTMl is "Moss Control plus Lawn Fertilizer" or "Turf Builder® Plus Moss Control". For St. Augustinegrass StepTMl is "Super Bonus® S."

step[™]2. Late Spring April-June

Weed Control plus Fertilizer - controls dandelion, clover, chickweed and other broadleaf weeds while greening your lawn for up to 2 more months. For St. Augustinegrass StepTM2 is "Super Turf Builder®."

step™3. Summer June-August

Insect Control plus Fertilizer - controls white grubs, chinch bugs, sod webworms, molé crickets, flea beetles, and cutworms, plus keeps your lawn green through the summer.

12 41

step™4. Fall August-November

Lawn Fertilizer - specially formulated for the fall - the best time to develop a stronger root system for a more vigorous, thicker lawn. Helps your lawn green-up even quicker next spring.

ants
armyworms
Bermudagrass mites
billbugs
chiggers
chinch bugs
clover mites
crickets
cutworms

earwigs
European crane fly larvae
(leather jackets)
fleas
flea beetles (in dichondra)
leafhoppers
mole crickets
sod webworms (lawn moth
larvae)

Brown dog ticks
vegetable weevil (in dichondra)
white grubs (Japanese beetle,
European
chafer, Masked chafer,
Phyllophaga May or June beetle)

(OPTIONAL BACK PANEL CLAIMS)

What to expect

Grubs and other insects listed will be controlled and prevented and your lawn will be greener in one week.

Fertilizer Plus Diazinon kills these active insects and provides effective control for approximately 6 weeks:

ants
armyworms
Bermudagrass mites
billbugs
chiggers
chinch bugs
clover mites
crickets
cutworms

earwigs
European crane fly larvae
(leather jackets)
fleas
flea beetles (in dichondra)
leashoppers
mole crickets
sod webworms (lawn moth larvae)
Brown dog ticks

vegetable weevil (in dichondra)
white grubs (Japanese beetle,
European
chafer, Masked chafer,
Phyllophaga May or June beetle)

EACH FERTILIZER PLUS DIAZINON GRANULE CONTAINS THE THREE PRIMARY NUTRIENTS AND INSECT CONTROL. We bond Scotts unique controlled-release nitrogen with other essential nutrients and an effective insect control so your lawn receives a safe, uniform feeding.

FERTILIZER PLUS DIAZINON WON'T BURN YOUR I AWN. The nitrogen is buffered. This means Fertilizer Plus Diazinon greens your lawn without the danger of burning. Part of the hittogen in Fertilizer Plus Diazinon is time-released to provide a supply of this estimated nutrient to help your lawn stay thick and green for up to two months.

How and when to apply

step ™ 3 DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Application

- Water the lawn thoroughly, allow to dry, and then apply product.
- Water lawn after application to wash granules from the lawn foliage. If treating for grubs, or mole crickets, water thoroughly so active ingredients will be carried into the root zone.
- For best results on grub control, apply from mid-July through August.
- Don't use on fruit or vegetable plantings.
- Do not allow traffic on your lawn until it has dried.
- Allow at least 4-6 weeks between program applications.
- Some lawn situations may require a supplemental application of Scott Lawn Insect Control (without fertilizer). You may all Scotts Toll-free to ask about your lawn (see below).

SPECIAL PRECAUTIONS TO REDUCE RISKS TO BIRDS

Birds, especially waterfowl (ducks and geese), feeding or drinking on treated areas may be killed. To reduce risks to birds:

- a) Do not treat areas larger than 15,000 sq. ft.
- b) Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20.
- c) Do not exceed maximum permitted label rates since rates above those recommended significantly increase potential hazards to birds, especially waterfowl.
- d) Avoid overlapping treated areas.
- e) Follow all watering instructions included within the directions for use, however, stop watering before puddling occurs.
- f) Do not apply this product to sites occupied by waterfowl or to sites within 75 feet of any body of water that may attract waterfowl, such as lakes or ponds.
- g) If necessary to repeat application, wait a minimum of 3 weeks before applications. Make no more than 4 applications per year.

Tips on Spreading

Scotts Drop:

- Treat entire lawn.
- First apply 2 header strips.
- Overlap spreader application area to avoid strips of untreated lawn.
- Shut off spreader when you reach the header strip or any time you stop.

EasyGreen® Rotary:

- Slide shut-off lever to the right, application automatically starts and stops when you do.
- Avoid misses by making each pass about 30 inches from the previous one so some overlapping occurs.

Speedy Green® Rotary:

- Shut off spreader anytime you stop.
- Avoid misses by making each pass about 5 feet from the previous one so some overlapping occurs.

Scotts Spreader Settings

Scotts	Scotts	Scotts
Drop	EasyGreen®	SpeedyGreen®
Spreaders	Řotary	Rotary
X	x	x

Additional Spreader Settings (Non-Scotts spreader settings are approximate. You may need to adjust for proper coverage.)

EZ Drop, Republic (#530)x	
EZ Broadcast, Republic (#540)x	
EZ Grow, Republic (#545)x	
Red Devil Drop (#FH-22)x	(To be determined at
Red Devil Rotary (#SB-50)x	time of production)
Quaker Drop (Model #10850)x	•
Quaker Rotary (Model #10301)x	
Ortho Drop (#5000)x	
Ortho Rotaryx	

If you have questions

We designed this Program to make having a better lawn as simple and easy as possible: But you may still have some questions about your lawn during the year. If so, you can contact the Scotts Lawn Pro retailer where you bought this Program. The the ready to answer your questions and provide any help or information that you may need. OR-You can get the answers by talking directly with one of Scotts lawn counselors. Simply dial this toll-free number 1-800-543-TURF.

Step 3 5M 39065-4

Scotts No-Quibble Guarantee

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

[UPC CODE] 0 32247 39065 1 PROOF OF PURCHASE

SCOTTS . . . the lawn people Marysville, Ohio 43041

©1995, The Scotts Company. World rights reserved.

[NOTE: Description below will be inverted.]

39065

Sec Sec.

Lawn Pro® (Scotts®)

Best Annual

stepTM 3

Program

EPANEW DOC/9

(ALTERNATE DIRECTIONS FOR USE)

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO apply when insect activity or damage (chewed or browned leaves) is first observed in the lawn.

NOTE: For Grub Control:

Northern States - The most likely period for grubs is late July through September, but in some cases grubs are active in the spring. An application for spring grub activity requires a repeat summer-early fall application for effective control.

Southern States - Apply during July-August. A second application may be needed six weeks later. Applications made later in the year, when grubs are more mature, are less effective.

For chinch bugs start treatments when daily high temperatures are 80°F or above and repeat at 6 week intervals.

For mole crickets, the best time to apply is in May to July.

DO apply when lawn is dry. For white grubs and mole crickets, water thoroughly, let lawn dry and then apply product.

DO water lawn after application to wash granules from the lawn foliage. If treating for grubs or mole crickets, water thoroughly (so that active ingredient will be carried into the rootzone).

DO make a repeat application of Scotts Lawn Insect Control (with or without fertilizer) if there are signs of renewed insect activity. (Normally, application gives effective control for approximately 6 weeks).

DON'T treat just the browned areas of your lawn. To prevent further damage, treat entire lawn.

DON'T use any other lawn control product for at least one week after application.

SPECIAL PRECAUTIONS TO REDUCE RISKS TO BIRDS

Birds, especially waterfowl (ducks and geese), feeding or drinking on treated areas may be killed. To reduce risks to birds:

a) Do not treat areas larger than 15,000 sq. ft.

- b) Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20.
- c) Do not exceed maximul. permitted label rates since rates above those recommended significantly increase potential hazards to birds, especially waterfowl.

d) Avoid overlapping treated areas.

- e) Follow all watering instructions included within the directions for use, however, stop watering before puddling occurs.
- f) Do not apply this product to sites occupied by waterfowl or to sites within 75 feet of any body of water that may attract waterfowl, such as lakes or ponds.
- g) If necessary to repeat application, wait a minimum of 3 weeks before applications. Make no more than 4 applications per year.

ALTERNATE DIRECTION FOR USE cont'd

STORAGE: Store in original container in a cool, dry, well-ventilated area inaccessible to children, and domestic animals. Do not store near fertilizers, herbicides, seeds, food, or feed. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

Spreading Tips

Scotts Drop: Mow lawn first if grass is tall enough to touch the bottom of the spreader. First apply two header strips. Operate spreader the longest way of the lawn. Overlap wheels to insure uniform feeding. shut off spreader when you have reached the

[Illus.]

header strips or stopped.

Scotts EasyGreen® Rotary: Slide shut-off lever to the right; application automatically starts and stops when you do. For complete coverage make each pass about 30 inches from the previous one so some overlapping occurs.

[]llus.]

Scotts SpeedyGreenTM Rotary: Shut off spreader anytime you stop. I or complete coverage make each pass about five feet from the previous one so some overlapping occurs. Avoid applying this product on flowers, vegetables and shrub plantings.

	Spreader Settings (covers 5,000 sq. ft.)
Scotts Drop Spreaders	X
Scotts EasyGreen®	XX
Rotary' Scotts SpeedyGreen®	Ÿ
Rotary	

Additional Spreader Settings

EZ Drop, Republic (#530) EZ Broadcast, Republic (#540) EZ Grow, Broadcast (#545)	xxx	
Ortho Drop	xxxx	
Ortho Rotary	xxxx	(to be determined)
Red Devil Drop (#FH-22)		
Red Devil Rotary (#SB-50)		
Quaker Drop (Model #10850)		
Quaker Rotary (Model #10301)		

(Non-Scotts spreader settings are approximate. You may need to adjust for proper coverage.)

Questions?

Call Scotts' toll-free Consumer Hotline: 1-800-543-TURF Scotts Policy No-Quibble Guarantee: If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

(Scotts®) . . . the lawn people Marysville, Ohio 43041

FERTILIZER PLUS DIAZINON
[UPC CODE]
0 32247 03105 9
PROOF OF PURCHASE

©1995, The Scotts Company. World rights reserved.

[BOTTOM]

[NOTE: Description below will be inverted.]

3105 Treats and fertilizes 5,000 sq. ft. (Scotts®) Fertilizer Plus Diazinon

