



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Registration Division (7505C)
 401 "M" St., S.W.
 Washington, D.C. 20460

EPA Reg. Number:
 538-255

Date of Issuance:
 OCT 22 1997

NOTICE OF PESTICIDE:
 Registration
 Reregistration

Term of Issuance:
 Conditional

Name of Pesticide Product:
 Fungicide XI

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

The Scotts Company
 14310 Scottslawn Road
 Marysville, OH 43041

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c)(5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following labeling change before you release the product for shipment:
 - a. Add the phrase "EPA Registration No. 538-255".
 - b. Under Precautionary Statements, add "Wear long-sleeved shirt and long pants, shoes and socks, and waterproof gloves (see enclosed review)".
 - c. Correct the typographical error under Recommended Rate in the Spreader Settings ("1,000").
3. Submit one copy of the final printed label for the record.

Signature of Approving Official:

Mary J. Walker

Date:

OCT 22 1997

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Mary L. Waller

Mary L. Waller
Acting Product Manager (21)
Fungicide Branch
Registration Division (7505C)

Enclosures

2/3/97 CW,ME

ACCEPTED
with COMMENTS
In EPA Letter Dated

OCT 22 1997
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
538-255

[FRONT]

(Scotts®)

ProTurf®

BRAND

Fungicide XI

- systemic fungicide that prevents and controls dollar spot (including benzimidazole tolerant strains), brown patch, anthracnose, rust, and other diseases.

Net Weight 00 lbs. (00 kg)

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION. Harmful if swallowed or absorbed through the skin. Causes eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Environmental Hazards: Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 to 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

ACTIVE INGREDIENT:

myclobutanil, a-butyl 2a-(chlorophenyl)-1H-1,2,4, triazol-1-propanenitrile..... 0.3 - 1.0%

INERT INGREDIENTS:..... 99.0-99.7% Total 100.00%

EPA Reg No 538-00 EPA Est 538-OH-1 (S) Product of U.S.A.

Distributed by: Professional Business Group, The Scotts Company 14111 Scottslawn Rd. Marysville, OH 43041 ©1997, The Scotts Company. World rights reserved.

Recommended For Use By Professional Turfgrass Managers

[TOP]

Fungicide XI 0000

[Note: Above copy will be inverted]

[RIGHT GUSSET]

0000 [UPC] (Scotts®) ProTurf® Fungicide XI VII 0 00000 00000 0 BRAND [Pkg. Date]

[LEFT GUSSET]

(Scotts®) ProTurf® Fungicide XI 0000 (Packaging Calendar) BRAND [UPC] 0 00000 00000 0

[BOTTOM]

0000 Fungicide XI

[Note: Above copy will be inverted]

[BACK]

(Scotts®)

ProTurf®

BRAND

Fungicide XI

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

For use on turfgrass areas consisting of Kentucky bluegrass, annual bluegrass, bentgrass, perennial ryegrass, fine fescue, zoysiagrass, tall fescue, bermudagrass, St. Augustinegrass, or mixtures where these grasses predominate.

TO PREVENT DOLLAR SPOT, ANTHRACNOSE, POWDERY MILDEW, AND RUST apply to moist or dry foliage at the RECOMMENDED RATE when environmental conditions favor disease activity. Repeat the application at 2 week intervals throughout the expected disease period.

TO CONTROL ACTIVE DOLLAR SPOT, ANTHRACNOSE, POWDERY MILDEW, AND RUST apply to moist or dry foliage at two week intervals. Use the DOUBLE RATE for light to moderate disease infestations and the HEAVY RATE for severe disease infestations. When the disease activity is under control, apply at the RECOMMENDED RATE at two week intervals.

TO PREVENT BROWN PATCH apply to moist or dry foliage at the DOUBLE RATE when environmental conditions favor disease activity. Repeat the application at 2 week intervals throughout the expected disease period.

TO CONTROL ACTIVE BROWN PATCH apply to moist or dry foliage at the HEAVY RATE every two weeks. When the disease activity is under control, apply at the DOUBLE RATE at two week intervals.

TO PREVENT NECROTIC RING SPOT AND SUMMER PATCH apply to moist or dry foliage at the HEAVY RATE. Begin in the spring when conditions are favorable for disease development. Make 2-4 applications at monthly intervals. For best results water in after application.

Note carefully

To avoid product pick-up, lightly irrigate treated areas soon after application.

On short cut bentgrass (1/2" or less) when temperatures are above 80°F, apply only to dry foliage.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reseal opened bag by folding down top and securing. Protect from water and extreme heat.

PESTICIDE DISPOSAL: Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY

This product has been researched to provide necessary data to support its uses listed on the label. The user should always follow the label directions and exercise his judgement and caution when using the product under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS, OR IMPLIED, CONCERNING THE RESULTS TO BE OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. The exclusive remedy of the user or Buyer, and the limit of liability of The Scotts Company or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The Scotts Company.

Professional Business Group
The Scotts Company
Marysville, OH 43041

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application.

00 lbs. treats 22,000 ft² (1/2 acre) at the RECOMMENDED RATE
(0.125 oz. ai/1,00 sq. ft.)

00 lbs. treats 11,000 ft² (1/4 acre) at the DOUBLE RATE
(0.25 oz. ai/1,000 sq. ft.)

00 lbs. treats 5,500 ft² (1/8 acre) at the HEAVY RATE
(0.5 oz. ai/1,000 sq. ft.)

<u>SPREADER</u>	<u>GROUND SPEED</u>	<u>WIDTH OF COVERAGE</u>	<u>SPREADER SETTINGS</u>		
			<u>RECOMMENDED</u>	<u>DOUBLE</u>	<u>HEAVY</u>
Gandy (30")	3 mph	overlap wheels	0	0	0
Scotts Drop	3 mph	overlap wheels	0	0	0
Scotts Rotaries					
R-7, R-7X*	3 mph	9 ft.	0	0	0
R-8 (cone 9)	3 mph	9 ft.	0	0	0
Lely Models					
(see below)	4.5 mph	18 ft.	0	0	0
Vicon (02)	4.5 mph	18 ft.	0	0	0
Vicon (03)	4.5 mph	10 ft.	0	0	0

Lely Models WTR, WFR, HR and 1250. PTO at 450 rpm.

Metric coverage: 00.00 kg treats 0000 m² at the RECOMMENDED RATE, 0000 m² at the DOUBLE RATE and 000 m² at the HEAVY RATE.

* 1/4 slide