

 <p>U.S. ENVIRONMENTAL PROTECTION AGENCY Office of Pesticide Programs Registration Division (E7505) 401 "M" St., S.W. Washington, D.C. 20460</p> <p>NOTICE OF PESTICIDE: <input checked="" type="checkbox"/> Registration <input type="checkbox"/> Reregistration</p> <p>(under FIFRA, as amended)</p>	<p>EPA Reg. Number: 538-258</p>	<p>Date of Issuance: JAN 13 1998</p>
	<p>Term of Issuance: Conditional</p>	
	<p>Name of Pesticide Product: Fertilizer and Diazinon</p>	
<p>Name and address of Registrant (include ZIP Code):</p> <p>The Scotts Company 14111 Scottslawn Road Marysville, OH 43041</p>		
<p><small>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</small></p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.</p> <p>Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p>		
<p>This product is conditionally registered in accordance with FIFRA sec. 3(c) (7) (A) provided that you:</p> <ol style="list-style-type: none"> 1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c) (5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4. 2. Make the labeling changes listed below before you release the product for shipment: <ol style="list-style-type: none"> a. Add the phrase "EPA Reg. No. 538-258". b. Revise the top of the bag claim "Kills fleas and ticks before they get to your pets" to "Kills fleas and Brown Dog ticks before they get to your pets". 		
<p>Signature of Approving Official:</p>		<p>Date: JAN 13 1998</p>

- c. Please be aware that the limitation "This product must NOT be used on golf courses and sod farms" must appear at the top (i.e., within the top half) of the front panel.
- d. Revise the optional front panel claim "Corrects most lawn insect problems" to "Corrects many lawn insect problems".
- e. On page 2, revise the claim "Kills fleas and ticks before they get on your pets" to "Kills fleas and Brown Dog ticks before they get on your pets".
- f. In the Environmental Hazards text, revise "Where irrigation (watering) is recommended in the directions or if waterfowl, i.e., ducks or geese, can be expected..." to read "Where watering is recommended in the directions or if waterfowl (ducks or geese) can be expected...".
- g. The SPECIAL PRECAUTIONS TO REDUCE RISKS TO BIRDS heading and text should appear within an outline box to make it prominent to the user.
- h. There appears to be discrepancies between the main and alternate directions for use. Specifically, the main directions instruct the user to water the lawn thoroughly before application (presumably for all pests), while the alternate directions instruct the user to thoroughly water before application when treating (only) for the pests white grubs and mole crickets. Also, while the main directions instruct the user to allow at least 4-6 weeks between program applications, no mention of specific time limitations between program applications appears in the alternate directions, which just indicate "Normally, application gives effective control for approximately 6 weeks." and to make repeat applications if there are signs of renewed insect activity (again, no specific minimum time interval is given if retreatment is required), potentially leaving the user to interpret the limitation to not use any other lawn control product for at least one week after application (which does not appear in the main directions) as an indication it could be appropriate to move on to the next program step (or even to reapply this product) after only one week, contradicting the minimum of 3 weeks between applications limitation that appears later in the SPECIAL PRECAUTIONS TO REDUCE RISKS TO BIRDS. The main and alternate directions should be clarified and made consistent to address the above concerns.

page 3 of 3
EPA Reg. No. 538-258

- i. Be sure to meet type size requirements for elements of the label with minimum type size requirements.

- 3. Submit three (3) copies of your final-printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

4/13

- Guaranteed
- Builds thick, green turf from the roots up without burning your lawn
- For best results, apply a Turf Builder[®] product four times a year (see back for details)
- Kills insects that cause your lawn to thin and turn brown
- Kills fleas and ticks before they get on your pets

MASTER LABEL

FRONT:

PACKAGE COPY FOR
FERTILIZER AND DIAZINON

10/10/96 CW/Ad 9/22/97VS
12/3/97 VS/Ad 12/12/97 CW/Ad

gw
12/12/97

Fertilizer and Diazinon

(Scotts[®])
(Lawn Pro[®])

step^{TM3}

Best
Annual
Program

ACCEPTED
with COMMENTS
in EPA Letter Dated

xxxx- (placed vertically) (Above copy to be inverted.)

Treats 5,000 sq. ft (Bag)

JAN 13 1998

"Available only at authorized Scotts[®] Lawn Pro[®] dealers"

Scotts[®] Best
Annual Program

SCOTTS POLICY
No-Quibble
Guarantee
SEE BACK FOR DETAILS

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

538-258

Step^{TM3}
of 4 easy steps to
a better lawn

Summer Lawn Fertilizer and Insect Control

For outdoor use on the home lawn only. This product must NOT be used on golf courses and sod farms.

OPTIONAL FRONT PANEL CLAIMS

—The Scotts Difference[®]—

- Contains diazinon to control white grubs, sod webworms, fleas, Brown dog ticks, chinch bugs, ants, European crane fly larvae, and other insects listed on back
- Timed-release nitrogen won't burn your lawn or cause surges growth
- Scotts No-Quibble Guarantee
- Toll-free Consumer Hotline: 1-800-543-TURF
- Scotts Best Annual Program for a Picture-Perfect lawn
- Apply in Summer: June - August

Corrects most lawn insect problems

Patented Particle Technology

- Thicker, greener turf - satisfaction guaranteed

538-XXX Fert + 3.2% Diazinon.doc

- Guaranteed
- Builds thick, green turf from the roots up without burning your lawn
- For best results, apply a Turf Builder® product four times a year (see back for details)
- Kills insects that cause your lawn to thin and turn brown
- Kills fleas and ticks before they get on your pets

Unlike other insect control products, every particle of Fertilizer and Diazinon contains

- Scotts patented timed-release nitrogen
- Phosphorus
- Potassium
- Insect control

Treats and fertilizes 5,000 sq. ft.

Kills chinch bugs
 Kills sod webworms
 Kills white grubs

NET WEIGHT 14.35 LB (6.50 kg)

Made in U.S.A.

XX-XX-XX

KEEP OUT OF REACH OF CHILDREN
CAUTION

See BACK for additional precautionary statements and directions for use.

ACTIVE INGREDIENT:

Diazinon [†]	3.20%
OTHER INGREDIENTS:	<u>96.80%</u>
Total	100.00%

Note to EPA: bag weights provided for 5000 sq ft bag coverage equals a rate of diazinon of 4.0 lbs per acre AI.

[†] CAS # 333-41-5

EPA Reg No 538-XXX

EPA Est 538-OH-1

[GUSSETS]

[RIGHT GUSSET]

XXXX X	[UPC] 0 32247 03105 9 Step™ 3	(Lawn Pro®) Treats and fertilizes 5,000 sq. ft. (Scotts®) Fertilizer and Diazinon	Best Annual Program
-----------	-------------------------------------	--	---------------------------

[LEFT GUSSET]

(Lawn Pro®) (Scotts®)	Treats and fertilizes 5,000 sq. ft. Fertilizer and Diazinon	[UPC] 0 32247 03105 9	XXXX X	Best Annual Program
--------------------------	--	--------------------------	-----------	---------------------------

[BACK]

Easy Open	PULL TAB and remove for pour spout [ILLUS.] PULL HERE	-OR-	CUT HERE TO OPEN
--------------	---	------	------------------

Treats and fertilizes 5,000 sq. ft.

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: CAUTION. Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust and contact with eyes, skin or clothing. Wash thoroughly after handling and before smoking or eating. Do not store near food or feed. Do not contaminate ornamental fish ponds. Do not allow children or pets onto the treated area until granules have been watered in and the grass or soil has dried.

First Aid: If swallowed: Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing. If inhaled: Remove victim to fresh air and apply artificial respiration if indicated. If on skin: Remove contaminated clothing and wash affected areas with soap and water. If in eyes: Rinse eyes with plenty of water and call a physician immediately.

Note to Physician: This product contains an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

Environmental Hazards: This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping granules. Where irrigation (watering) is recommended in the directions or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after applying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

Fertilizer and Diazinon

643-XXXX

Guaranteed Analysis

Total nitrogen (N).....	XX%
X.X% ammoniacal nitrogen	
XX.X% urea, methylene urea, nitrogen*	
Available phosphate (P ₂ O ₅).....	X%
Soluble potash (K ₂ O).....	X%

Derived from: monoammonium phosphate, urea, methylene ureas and potassium sulfate.

*Contains xx% slowly available methylenediurea and dimethylenetriurea nitrogen.

STORAGE AND DISPOSAL

STORAGE: Store in original container in a cool, dry, well-ventilated area inaccessible to children, and domestic animals. Do not store near fertilizers, herbicides, seeds, food, or feed. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

OPTIONAL BACK PANEL CLAIMS

Now it's easy for you to have a better lawn

Save money

Why even consider a lawn care service? Scotts® Lawn Pro® Annual Program will give you a beautiful lawn for about one-third the cost.

Simple and easy

No more guesswork. You simply apply the right product - one Step at a time.

Start anytime

You can begin the Annual Program anytime of the year. Just check the month and select the appropriate Step listed below. Apply the remaining Steps at two-month intervals.

Guaranteed results

Each Step is designed to the specific season to help give you a greener lawn.

[The 4 Steps listed below will be placed horizontally in one row.]

Step™1.

Early Spring

February-April

Crabgrass Preventer plus Fertilizer - prevents crabgrass, foxtail, spurge, oxalis, chickweed and other annual weeds while providing deep, thick greening for up to 2 months. In the Pacific Northwest, Step™1 is "Moss Control plus Lawn Fertilizer" or "Turf Builder® Plus Moss Control". For St. Augustinegrass Step™ 1 is "Super Bonus S®."

**Step™2.
Late Spring
April-June**

Weed Control plus Fertilizer - controls dandelion, clover, chickweed and other broadleaf weeds while greening your lawn for up to 2 more months. For St. Augustinegrass Step™2 is "Super Turf Builder®."

**Step™3.
Summer
June-August**

Insect Control with Fertilizer - controls white grubs, chinch bugs, sod webworms (lawn moths), mole crickets, flea beetles, cutworms, and other lawn insects, plus keeps your lawn green through the summer. For grub control, apply "Scotts® GrubEx®" anytime from April to mid-August.

**Step™4.
Fall
August-November**

Lawn Fertilizer - specially formulated for the fall - the best time to develop a stronger root system for a more vigorous, thicker lawn. Helps your lawn green-up even quicker next spring.

Turf Builder Annual Programs

- No more guess work . . . apply the right product at the right time for a thick, green, weed-free lawn.
- Start anytime . . . just check the month below and apply products at 2 month intervals.

[The 4 products listed below will be placed horizontally in one row.]

Turf Builder With Halts — Apply in early spring (February-April). Prevents crabgrass and other annual weeds while feeding your lawn. (In the Pacific Northwest, substitute Turf builder Plus Moss Control. For St. Augustinegrass, substitute Bonus S. For dormant bermudagrass substitute Halts.)

Turf Builder With Plus 2 — Apply in late spring (April-June). Kills dandelions and other broadleaf weeds while feeding your lawn. (for St. Augustinegrass, substitute Turf Builder.)

Turf Builder With Insect Control — Apply in summer (June-August). Kills lawn damaging insects, fleas and ticks while feeding your lawn. For grub control, apply Scotts® GrubEx® anytime from April to mid-August.

Turf Builder — Apply in fall (August-November). Important time to feed for a thicker, greener lawn.

What to expect

Grubs and other insects listed will be controlled and prevented and your lawn will be greener in one week.

Fertilizer and Diazinon kills these active insects and provides effective control for approximately 6 weeks:

ants	cutworms	sod webworms (lawn moth larvae)
armyworms	earwigs	Brown dog ticks
Bermudagrass mites	European crane fly larvae	vegetable weevil (in dichondra)
billbugs	(leather jackets)	white grubs (Japanese beetle, European
chiggers	fleas	chafer, Masked chafer Phyllophaga -
chinch bugs	flea beetles (in dichondra)	May or June beetle)
clover mites	leafhoppers	
crickets	mole crickets	

EACH FERTILIZER PLUS DIAZINON GRANULE CONTAINS THE THREE PRIMARY NUTRIENTS AND INSECT CONTROL. We bond Scotts unique controlled-release nitrogen with other essential nutrients and an effective insect control so your lawn receives a safe, uniform feeding.

FERTILIZER AND DIAZINON WON'T BURN YOUR LAWN. The nitrogen is buffered. This means Fertilizer and Diazinon greens your lawn without the danger of burning. Part of the nitrogen in Fertilizer and Diazinon is time-released to provide a supply of this essential nutrient to help your lawn stay thick and green for up to two months.

-OR-

The Scotts Difference®

Unlike ordinary insect control products, every particle of Fertilizer and Diazinon contains: Scotts patented timed-release nitrogen, phosphorus, potassium and insect control.

Scotts Fertilizer and Diazinon

Fertilizer and Diazinon particles deliver the same ratio of nitrogen, phosphorus, potassium and insect control to every square inch of turf for even, consistent feeding plus better insect control.

Ordinary Mixed Insect Controls

Granules contain different nutrients. Nutrient and insect control granules of different sizes and shapes can separate during application resulting in uneven, inconsistent feeding and insect control.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Application

- Water the lawn thoroughly, allow to dry, and then apply product.
- Water lawn after application to wash granules from the lawn foliage. If treating for grubs, or mole crickets, water thoroughly so active ingredients will be carried into the root zone.
- For best results on grub control, apply from mid-July through August.
- Don't use on fruit or vegetable plantings.
- Do not allow traffic on your lawn until it has dried.
- Allow at least 4-6 weeks between program applications.
- Some lawn situations may require a supplemental application of Scotts Lawn Insect Control (without fertilizer), or Scotts Grubex® (without fertilizer). You may call Scotts toll-free to ask about your lawn (see below).

SPECIAL PRECAUTIONS TO REDUCE RISKS TO BIRDS

Birds, especially waterfowl (ducks and geese), feeding or drinking on treated areas may be killed. To reduce risks to birds:

- a) Do not treat areas larger than 15,000 sq. ft.
- b) Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20.
- c) Do not exceed maximum permitted label rates since rates above those recommended significantly increase potential hazards to birds, especially waterfowl.
- d) Avoid overlapping treated areas.
- e) Follow all watering instructions included within the directions for use, however, stop watering before puddling occurs.
- f) Do not apply this product to sites occupied by waterfowl or to sites within 75 feet of any body of water that may attract waterfowl, such as lakes or ponds.
- g) If necessary to repeat application, wait a minimum of 3 weeks before applications. Make no more than 4 applications per year.

Tips on Spreading

Scotts Drop:

- Treat entire lawn.
- First apply 2 header strips.
- Overlap spreader application area to avoid strips of untreated lawn.
- Shut off spreader when you reach the header strip or any time you stop.

EasyGreen® Rotary:

- Slide shut-off lever to the right, application automatically starts and stops when you do.
- Avoid misses by making each pass about 30 inches from the previous one so some overlapping occurs.

Speedy Green® Rotary:

- Shut off spreader anytime you stop.
- Avoid misses by making each pass about 3 feet from the previous one so some overlapping occurs.

Scotts Spreader Settings

Scotts Drop Spreaders	Scotts EasyGreen® Rotary	Scotts SpeedyGreen® Rotary
x	x	x

Additional Spreader Settings (Non-Scotts spreader settings are approximate. You may need to adjust for proper coverage.)

EZ Drop, Republic (#530)	x	
EZ Broadcast, Republic (#340)	x	
EZ Grow, Republic (#545)	x	
Red Devil Drop (#FH-22)	x	
Red Devil Rotary (#SB-50)	x	
Quaker Drop (Model #10850)	x	
Quaker Rotary (Model #10301)	x	
Ortho Drop (#5000)	x	
Ortho Rotary	x	

(To be determined at time of production)

This package should cover 5,000 sq. ft. (for example 50 ft. x 100 ft.). Close spreader when filling, stopping or turning.

11/8/13

If you have questions

We designed this Program to make having a better lawn as simple and easy as possible. But you may still have some questions about your lawn during the year. If so, you can contact the Scotts Lawn Pro retailer where you bought this Program. They are ready to answer your questions and provide any help or information that you may need. **-OR-** You can get the answers by talking directly with one of Scotts' Lawn Counselors. Simply dial this toll-free number 1-800-543-TURF.

(OPTIONAL)

Your Do-It-Yourself Lawn Care Headquarters

Scotts® Lawn Pro® retailers can provide you with the do-it-yourself expertise and products to have a beautiful Scotts lawn simply and easily . . . for less money than hiring a lawn service. That's why we call them Lawn Pros!

Questions?

If you have any questions you can talk directly with a Scotts® Lawn Counselor. Simply call the toll-free number 1-800-543-TURF.

Questions, Comments, Information

For Additional Questions, Comments and Information contact the Scotts® Lawn Pro® retailer where you bought this Program OR
Call Scotts' toll-free Consumer Helpline: 1-800-543-TURF
"Lawn Success Made Easy"
Visit Scotts' website at www.scottslawn.com

Scotts No-Quibble Guarantee™

If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

[UPC CODE]
0 32247 39065 1
PROOF OF PURCHASE

SCOTTS . . . the lawn people
Marysville, Ohio 43041

©1998, The Scotts Company. World rights reserved.

[NOTE: Description below will be inverted.]

~~XXXXX~~
X

Lawn Pro®
(Scotts®)

Best
Annual
Program

Step™ 3

(ALTERNATE DIRECTIONS FOR USE)

How and when to apply

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO apply when insect activity or damage (chewed or browned leaves) is first observed in the lawn.

NOTE: For Grub Control:

Northern States - The most likely period for grubs is late July through September, but in some cases grubs are active in the spring. An application for spring grub activity requires a repeat summer-early fall application for effective control.

Southern States - Apply during July-August. A second application may be needed six weeks later. Applications made later in the year, when grubs are more mature, are less effective.

For chinch bugs start treatments when daily high temperatures are 80°F or above and repeat at 6 week intervals.

For mole crickets, the best time to apply is in May to July.

DO apply when lawn is dry. For white grubs and mole crickets, water thoroughly, let lawn dry and then apply product.

DO water lawn after application to wash granules from the lawn foliage. If treating for grubs or mole crickets, water thoroughly (so that active ingredient will be carried into the rootzone).

DO make a repeat application of Scotts Lawn Insect Control (with or without fertilizer) if there are signs of renewed insect activity. (Normally, application gives effective control for approximately 6 weeks).

DONT treat just the browned areas of your lawn. To prevent further damage, treat entire lawn.

DONT use any other lawn control product for at least one week after application.

SPECIAL PRECAUTIONS TO REDUCE RISKS TO BIRDS

Birds, especially waterfowl (ducks and geese), feeding or drinking on treated areas may be killed. To reduce risks to birds:

- a) Do not treat areas larger than 15,000 sq. ft.
- b) Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20.
- c) Do not exceed maximum permitted label rates since rates above those recommended significantly increase potential hazards to birds, especially waterfowl.
- d) Avoid overlapping treated areas.
- e) Follow all watering instructions included within the directions for use, however, stop watering before puddling occurs.
- f) Do not apply this product to sites occupied by waterfowl or to sites within 75 feet of any body of water that may attract waterfowl, such as lakes or ponds.
- g) If necessary to repeat application, wait a minimum of 3 weeks before applications. Make no more than 4 applications per year.

STORAGE: Store in original container in a cool, dry, well-ventilated area inaccessible to children, and domestic animals. Do not store near fertilizers, herbicides, seeds, food, or feed. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

Spreading Tips

Scotts Drop: Mow lawn first if grass is tall enough to touch the bottom of the spreader. First apply two header strips. Operate spreader the longest way of the lawn. Overlap wheels to insure uniform feeding. Shut off spreader when you have reached the header strips or stopped.

Scotts EasyGreen® Rotary: Slide shut-off lever to the right; application automatically starts and stops when you do. For complete coverage make each pass about 30 inches from the previous one so some overlapping occurs.

Scotts SpeedyGreen™ Rotary: Shut off spreader anytime you stop. For complete coverage make each pass about five feet from the previous one so some overlapping occurs. Avoid applying this product on flowers, vegetables and shrub plantings.

	Spreader Settings (covers 5,000 sq. ft.)
Scotts Drop Spreaders	X
Scotts EasyGreen® Rotary	XX
Scotts SpeedyGreen® Rotary	X

Additional Spreader Settings

EZ Drop, Republic (#530)	XXXX	
EZ Broadcast, Republic (#540).....	XXX	
EZ Grow, Broadcast (#545).....	X	
Ortho Drop.....	XXXX	
Ortho Rotary	XXXXX	(to be determined)
Red Devil Drop (#FH-22).....	XX	
Red Devil Rotary (#SB-50).....	XXXXX	
Quaker Drop (Model #10850).....	X	
Quaker Rotary (Model #10301).....	X	

(Non-Scotts spreader settings are approximate. You may need to adjust for proper coverage.)

Scotts Policy No-Quibble Guarantee™: If for any reason you are not satisfied with results after using this product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

(Scotts®) . . . the lawn people
Marysville, Ohio 43041

FERTILIZER AND DIAZINON
[UPC CODE]
0 32247 03105 9
PROOF OF PURCHASE

©1998, The Scotts Company. World rights reserved.

[BOTTOM] [NOTE: Description below will be inverted.]

XXXX Treats and fertilizes 5,000 sq. ft.
x (Scotts®) Fertilizer and Diazinon