


Anderson's ProTurf Fertilizer Plus Weedgrass Control
538-XXX

Anderson's ProTurf Fertilizer Plus Turf
Weedgrass Control



XX-X-X
Anderson's
Fertilizer Plus
Turf Weedgrass
Control

- preemergent control of crabgrass, goosegrass, foxtail, chickweed, oxalis, spurge, and *Poa annua*
- provides controlled-release nitrogen feeding

Net Weight 66.0 lbs. (XX.XX kg)

KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS

Hazards to Human and Domestic Animals: CAUTION, causes eye irritation. Avoid contact with eyes, skin, or clothing. Wash after handling. May be harmful if swallowed. Avoid inhalation. Do not allow people or pets to enter the treated area until dusts have settled. Do not contaminate feed or foodstuffs. Do not graze treated areas. Do not feed clippings to livestock.

FIRST AID

If in eyes: Call A poison control center immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-t-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling poison control center or doctor, or going for treatment.

Environmental Hazards: This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment washwaters. In case of spills, either collect for use or dispose of properly.

ACTIVE INGREDIENT:

Pendimethalin† 1.15%

INERT INGREDIENTS: 98.85%

Total 100.00%

EPA Reg. No. 538-XXX

EPA Est 538-OH-1

Fertilizer Plus Weedgrass Control XX-X-X

XXXX

Guaranteed Analysis

Total nitrogen (N)	XX%
X.X% ammoniacal nitrogen	
X.X% urea, methylene ureas nitrogen	
X.X% water insoluble nitrogen	
Available phosphate (P ₂ O ₅)	X%
Soluble potash (K ₂ O)	X%

Derived from: urea, methylene ureas, monoammonium phosphate and potassium sulfate

Product of USA
U.S. Patent No's 5,219,465; 5,300,135

The Scotts Company
14111 Scottslawn Road
Marysville, Ohio 43041
© 2000. The Scotts Company. World rights reserved.

Recommended for Use by Professional Turfgrass Managers

XXXX	XX-X-X
	Anderson's ProTurf®
	Fertilizer Plus
	Turf Weedgrass
	Control

GUSSETS:

XXXX	Scotts®	XX-X-X
	ProTurf®	Anderson's
		Fertilizer Plus
		Turf Weedgrass
		Control

BACK:

Scotts®	XX-X-X
ProTurf®	Anderson's
BRAND	Fertilizer Plus
	Turf Weedgrass
	Control

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Do not apply this product in a way that will contact workers or other persons or pets, either directly or through drift. Only protected handlers may be in areas during application. Keep people and pets out of the area during application.

For use on established turfgrasses including Kentucky bluegrass, perennial ryegrass, fine fescue, tall fescue, bentgrass and *Poa annua* (except putting greens and tees), bermudagrass, St. Augustinegrass, bahiagrass, zoysiagrass, centipedegrass and blends of these grasses. See "Bermudagrass, Bentgrass and *Poa annua* Precautions."

Not for use on bentgrass and *Poa annua* putting greens and tees, or on dichondra.

Prevents sprouting of the following grassy and dicot weeds and provides a controlled-release nitrogen feeding:

<u>Grassy Weeds</u>	<u>Dicot Weeds</u>	
barnyardgrass	chickweed	knotweed
crabgrass	cudweed	oxalis
fall panicum	eveningprimrose	prostrate spurge*
foxtail	henbit	purslane
goosegrass*	hop clover	
<i>Poa annua</i>		

To prevent crabgrass, foxtail, barnyardgrass, fall panicum, *Poa annua*, cudweed, eveningprimrose, chickweed, oxalis, spurge, henbit and knotweed, apply product to dry foliage at the recommended rate in the spring or fall before they sprout.

* To prevent goosegrass and heavy infestations of spurge, repeat application at the suggested rate after 6-8 weeks.

To prevent *Poa annua*, henbit and fall germinating chickweed, apply product to dry foliage at the suggested rate in late summer-early fall before germination. A repeat application after *Poa annua* treatments should not be made until the following spring.

Apply product uniformly to dry foliage at the suggested rate with a properly calibrated spreader. For best result, water after application (1/4 to 1/2 inch) to remove granules from the foliage and to move the product to the soil surface for an effective chemical barrier. Do not destroy the chemical barrier created at the soil surface by raking or aerifying subsequent to application. Allow at least a four month span between any two application except when repeat treatments are necessary after 6-8 weeks.

Do not mow treated area until product has been watered in and turf foliage has dried.

Precautions

Do not use on bentgrass and *Poa annua* putting greens and tees where it is the desired species, or on dichondra.

Do not apply this product to cool season turfgrasses when temperatures are regularly in the 80's.

This product may cause discoloration and thinning where overapplications have been made if the turfgrass is under severe stress conditions (disease, drought, low temperature – below 40°F, scalping, and excessive soil moisture).

Bermudagrass, ½ Inch Height or Taller Bentgrass and *Poa annua* Precautions

Do not apply to dormant bentgrass and *Poa annua* (turf that is not actively growing or has experienced frost).

When applied to dormant bermudagrass, green-up may be delayed to 2 to 3 weeks in the spring. The delay in green-up may be more extensive on low-cut bermudagrass and on sandy soils.

Where turf thinning or damage has occurred due to winter injury, wait until greenup, fill-in and good rooting before making an application.

Do not apply to bentgrass or *Poa annua* after late summer.

Application of this product to bermudagrass overseeded with cool season turfgrass species may injure and thin the overseeding.

Other Precautions

Delay seeding by 4 months and sprigging by 5 months after application. Prior to planting, for best results, the soil surface should be disturbed or worked to dilute any remaining chemical residue.

On newly seeded areas, wait until the grass has been mowed at least 4 times and has filled in before making an application.

Do not apply more than a cumulative total of 261 lbs. of product/acre (3.0 lbs. active ingredient/acre) per year on bentgrass and *Poa annua*; 522 lbs. of product/acre (6.0 lbs. active ingredient/acre) per year on cool season grasses; and 783 lbs. of product/acre (9.0 lbs. active ingredient/acre) per year on warm season grasses.

Avoid application to cement, stone, clothing or shoes. If contact occurs, brush off particles to prevent staining. If necessary, rinse with water. If metallic surfaces (golf course equipment, automobiles, etc.) are

present at the application site, do not use this product under windy conditions (5-10 MPH) due to its staining potential.

Where turf has been severely thinned due to winter damage, insects, disease, scalping, etc., an application should be withheld until filling-in and rooting of stolons/rhizomes is complete and desired turf density is achieved.

Turfgrass Species	Weeds	Rates of Anderson's Fertilizer Plus Weedgrass Control	Comments
Cool Season Grasses Kentucky bluegrass Perennial ryegrass Fine fescue Tall fescue	barnyardgrass chickweed crabgrass fall panicum foxtail knotweed oxalis <i>Poa annua</i> prostrate spurge purslane	132-176 lbs./acre as the initial application prior to weed germination	Apply a repeat application of 110-152 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected
	goosegrass	132-176 lbs./acre in the spring prior to weed germination. Apply a repeat application after 6-8 weeks at 132-154 lbs./acre	
Bentgrass and established <i>Poa annua</i> (1/2 inch height or taller. Not for use on greens and tees)	barnyardgrass crabgrass fall panicum foxtail knotweed oxalis <i>Poa annua</i> prostrate spurge purslane	132 lbs./acre in the spring prior to weed germination	Apply a repeat application of 110-132 lbs./acre after 6-8 weeks where heavy infestations are expected
	goosegrass	132 lbs./acre in the spring prior to weed germination with a repeat application of 132 lbs./acre after 6-8 weeks	
Warm Season Grasses Bermudagrass St. Augustinegrass Bahiagrass Zoysiagrass Centipedegrass Tall fescue	barnyardgrass chickweed crabgrass fall panicum foxtail knotweed oxalis <i>Poa annua</i> prostrate spurge purslane	132-176 lbs./acre prior to weed germination	Apply a repeat application of 110-132 lbs./acre after 6-8 weeks for extended control or where heavy weed infestations are expected
	cudweed eveningprimrose goosegrass henbit hop clover	132-176 lbs./acre prior to weed germination	Apply a repeat application of 132-154 lbs./acre after 6-8 weeks

Bermudagrass Greens	barnyardgrass chickweed crabgrass fall panicum foxtail knotweed oxalis Poa annua prostrate spurge purslane	132 lbs./acre prior to weed germination	Apply a repeat application of 132 lbs./acre after 6-8 weeks if necessary
	goosegrass cudweed hop clover eveningprimrose henbit	132 lbs./acre prior to weed germination with a repeat application of 132 lbs./acre after 6-8 weeks	

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in a clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Suggested Spreader Settings

To provide proper distribution calibrate spreader before application.
Do not adjust rates in order to meet fertilizer requirements of your soil.

66.0 lbs. treats 22,000 ft² (1/4 acre) at the 132 lbs. product/acre rate
(equivalent to 1.5 lb. active ingredient/acre and 2.0 lb. N/1,000 sq. ft.)

Spreader	Ground or PTO Speed	Width of Coverage	Recommended Rate Range Lbs. product/acre			
			110	132	154	176
Scotts Drop	X mph	overlap wheels	X	X	X	X
Gandy 30"	X mph	overlap wheels	X	X	X	X
Scotts Rotary* R-7, R-7X	X mph	X feet	X	X	X	X
Scotts Rotary* R-8, (cone 8)	X mph	X feet	X	X	X	X
Lely Models	X mph	X feet	X	X	X	X

- Use discharge hole cover #13030 (already attached to the PT-10 and PT-11 models). For the R-7X model use with adjustable pattern slide ¼ closed.

Lely models WTR, WFR, HR and 1250. PTO at 450 rpm.

Metric Coverage: XX.XX kg treats 1022 m² at the 132 lbs. product/acre rate.

LIMITATION OF LIABILITY

It must be understood that this product has not been evaluated on all bentgrass varieties and *Poa annua* under all possible growing conditions. The user should exercise judgement and caution when using this product on a given variety until familiar with the performance under his growing conditions. NO WARRANTY OR REPRESENTATION IS MADE, EXPRESS OR IMPLIED, CONCERNING THE RESULTS OBTAINED FROM THE USE OF THIS PRODUCT ON BENTGRASS AND *POA ANNUA* IF NOT USED IN ACCORDANCE WITH DIRECTIONS AND ESTABLISHED SAFE PRACTICES. The exclusive remedy of the user or Buyer, and the limit of liability of The Scotts Company or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or Buyer for the quantity of this product involved. The Buyer and all users are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing signed by a duly authorized representative of The Scotts Company.

The Scotts Company
14111 Scottslawn Road
Marysville, Ohio 43041