

538-286

04/16/2001

1/8



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (7505C)
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg.
Number:

538-286

Date of Issuance:

APR 16 2001

NOTICE OF PESTICIDE:
 X Registration
 Reregistration

Term of Issuance:

Conditional

Name of Pesticide Product:

Proturf Anderson's
Fungicide X

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

The Scotts Company
14111 Scottslawn Road
Marysville, Ohio 43041

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA sec. 3(c) (7) (A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product under FIFRA sec. 3(c) (5) when the Agency requires all registrants of similar products to submit such data; and submit acceptable responses required for reregistration of your product under FIFRA section 4.
2. Make the following labeling change before you release the product for shipment: add the phrase "EPA Registration No. 538-286".
3. Submit one copy of the final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Enclosure

Signature of Approving Official:

Mary L. Waller
Mary L. Waller, PM (21)

Date:

APR 16 2001

MASTER LABEL

FRONT

**ANDERSONS' FUNGICIDE X
EPA REG. NO 538-XXX**

(Scotts®)
ProTurf®
BRAND

(Scotts®)
ProTurf®
BRAND

(Scotts®)
ProTurf®
BRAND

Andersons' Fungicide X

Optional Front Panel Claims

- Prevents and controls brown patch, leaf spot, red leaf spot, pink snow mold and gray snow mold.

NET WEIGHT XX.XX lbs (XX.XX kg)

**KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS**

Hazards to Humans and Domestic Animals: CAUTION, Harmful if absorbed through skin or inhaled. Avoid contact with eye, skin or clothing. Avoid breathing dust. Wear long-sleeved shirt and long pants, socks and shoes and chemical-resistant gloves (such as Barrier Laminate, Butyl rubber, Nitrile rubber and Viton)

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call A poison control center immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling poison control center or doctor, or going for treatment.

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**

APR 16 2001

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.**

538-286

User Safety Recommendations

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide get inside. Then wash thoroughly and put on clean clothing.

Users should remove personal protective equipment immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothes.

Environmental Hazards:

This chemical can contaminate surface water through aerial and ground applications. Under some conditions, it may also have a high potential for runoff into surface water application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water areas not separated from adjacent surface waters with vegetated filter strips, and areas over-laying tile drainage systems that drain surface water. This pesticide is toxic to invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Cover, incorporate, or clean up spills. Do not contaminate water when disposing of equipment washwater to rinsate.

ACTIVE INGREDIENT:

Iprodione * 1.30%

INERT INGREDIENTS: 98.70%

Total 100.00%

*CAS #36734-19-7

EPA Reg No 538-XXX

EPA Est 538-OH-1

Product USA

Distributed By:

Professional Business Group

The Scotts Company

14111 Scottslawn Road

Marysville, Ohio 43041

(c) 2001. The Scotts Company, Marysville, Ohio 43041 World rights reserved

Recommended for Use by Professional Turfgrass Managers

GUSSETS:




RIGHT GUSSET:

538-XXX

(Scotts®)

Andersons' Fungicide X
(ProTurf®)
BRAND



LEFT GUSSET:

(Scotts®)
(ProTurf®)
BRAND

Andersons' Fungicide X

538-XXX

BACK:

538-XXX



Andersons' Fungicide X

(Scotts®)
(ProTurf®)
BRAND

Andersons' Fungicide X

Directions for Use

It is a violation of federal law to use this product in manner inconsistent with its labeling.

Use of this product at residential sites is prohibited.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

Entry Restrictions

Do not enter or allow others to enter the treated area until dusts have settled.

This fungicide product prevents and controls the following turfgrass diseases.

- brown patch
- leaf spot (Helminthosporium spp.)
- red leaf spot
- pink snow mold (Fusarium nivale)
- gray snow mold (Typhula spp.)

For use on fairways, putting greens and other turf areas consisting of Kentucky bluegrass, perennial ryegrass, fine fescue and bermudagrass, or where these mixtures predominate. Product may also be applied to St. Augustinegrass turf areas.

WHEN TO USE

Brown Path and Red Leaf Spot

To prevent, apply at the NORMAL or HEAVY RATE (1.0 to 1.5 oz ai/1000 sq ft) to moist or dry foliage when disease activity is anticipated and repeat at 2-week intervals. If heavy disease pressures are encountered, apply at the HEAVY RATE to moist or dry foliage. When diseases are brought under control, revert back to bi-weekly NORMAL RATE application intervals.

To control, apply to moist or dry foliage at the HEAVY RATE (1.5 oz ai/1000 sq ft) as soon as disease activity is noticed. Repeat at bi-weekly until disease is under control, then revert back to NORMAL or HEAVY RATE applications at bi-weekly.

Pink and Gray Snow Mold

To prevent. apply at the LIGHT or NORMAL RATE (0.5 to 1.0 oz ai/1000 sq ft) to moist or dry foliage when disease activity is anticipated. Repeat at 2 to 3 week intervals.

To control. apply to dry foliage at the NORMAL or HEAVY RATE (1.0 to 1.5 oz ai/1000 sq ft) when disease activity is first noticed. Repeat as needed at this rate at 2 to 3 week intervals.

When disease activity is under control, revert to the LIGHT or NORMAL RATE (0.5 to 1.0 oz ai/1000 sq ft) at 2 to 3 week intervals.

For suppression. apply to dry foliage at the NORMAL or HEAVY RATE (1.0 to 1.5 oz ai/1000 sq ft) prior to the first snow. Repeat in mid-winter when turf is free of snow and again immediately after the final snow melts in the spring.

Not a season long (all winter) control for snow molds unless applied on a 2 to 3 week application program.

Leaf Spot

To prevent. apply to moist or dry foliage at the LIGHT RATE (0.5 oz ai/1000 sq ft) when disease activity is anticipated. Repeat if needed after 3 to 4 weeks.

To control. apply to moist foliage at the NORMAL to HEAVY RATE (1.0 to 1.5 oz ai/1000 sq ft) when disease activity is first noticed. Repeat as needed at intervals not closer than 2 weeks.

HOW TO USE

For LIGHT, NORMAL or HEAVY RATE applications (0.5, 1.0 or 1.5 oz ai/1000 sq ft), apply to moist or dry foliage for summer diseases. Applications need not be watered in. Apply to dry foliage for snow mold prevention/control.

Delay mowing for 24 hours after application to prevent particle pickup by mowers.

Expected time disease activity:

Spring – leaf spot, pink snow mold (Fusarium patch)
gray snow mold

Summer – leaf spot and brown patch

Fall – leaf spot and pink snow mold (Fusarium patch)

Winter – pink snow mold (Fusarium patch),
and gray snow mold

Allow one week between the application of this and other control products. This guideline may be relaxed where a severe disease attack requires immediate treatment.

NOTE TO USER

Do not apply this product in a way that will contact workers or other persons, either directly or indirectly or through drift. Only protected handlers may be in the area during application.

Except for use on golf courses, if applying this product adjacent to a water body such as a lake, estuary, reservoir, river, permanent stream, marsh or natural pond, or commercial fish pond, there must be at least a 25-foot vegetative buffer strip between the water body and the point of application.

For golf courses only, do not apply to turf cut higher than 1" on golf holes where water bodies are present.

Do not apply this product when the wind direction is toward aquatic areas.

May apply up to but no more than 24 pounds active ingredient (ai.) annually. Do not apply more than 6 times per year.

Do not apply this product using "belly grinder" or other handheld application equipment.

Storage and disposal

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in clean, dry place. Reseal opened bag by folding top down and securing.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment, Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out smoke.

IMPORTANT NOTICE: DISCLAIMER AND LIMITATION OF LIABILITY

This product has been researched to provide the necessary data to support its uses listed on the label. The user should always follow the label directions and exercise his judgment and caution when using the product under his growing conditions. NO WARRANTY OR REPRESENTATION IF MADE, EXPRESS, OR IMPLIED, CONCERNING THE RESULTS TO BE OBTAINED FROM THE USE OF THIS PRODUCT IF NOT USED IN ACCORDANCE WITH DIRECTIONS OR ESTABLISHED SAFE PRACTICE. The exclusive remedy of the user or buyer, and the limit of liability of The Scotts Company or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user are deemed to have accepted the terms of this Notice, which may be varied only by agreement in writing, signed by a duly authorized representative of The Scotts Company.

Professional Business Group
The Scotts Company
Marysville, Ohio 43041

BACK CONTINUED:

Suggested spreader Settings

To provide proper distribution calibrate spreader before application

- 52.86 lbs (23.97 kg) treats 22,000 ft² at the LIGHT RATE
- 52.86 lbs (23.97 kg) treats 11,000 ft² at the NORMAL RATE
- 52.86 lbs (23.97 kg) treats 7,330 ft² at the HEAVY RATE

SPREADER	GROUND OR PTO SPEED	WIDTH OF COVERAGE	SPREADER SETTINGS		
			LIGHT RATE	NORMAL RATE	HEAVY RATE
Scotts Drop Type	X	X	X	X	X
Scotts Rotaries	X	X	X	X	X
Lely Models (see below)	X	X	X	X	X

Lely Models WTR, WFR, HR and 1250. PTO at 450 RPM