



U.S. ENVIRONMENTAL PROTECTION AGENCY
Office of Pesticide Programs
Registration Division (H7505C)
401 "M" St., S.W.
Washington, D.C. 20460

EPA Reg. Number:
538-300

Date of Issuance:
JUN - 7 2005

NOTICE OF PESTICIDE:
 Registration
 Reregistration

Term of Issuance:
Conditional

Name of Pesticide Product:
Scotts Granular MSM

(under FIFRA, as amended)

Name and Address of Registrant (include ZIP Code):

The Scotts Company
14111 Scottslawn Rd.
Marysville, OH 43041

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act.

Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit and/or cite all data required for registration/reregistration of your product when the Agency requires all registrants of similar products to submit such data.
2. Add the phrase "EPA Registration No. 538-300" to your label before you release the product for shipment.
3. Submit two (2) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of this product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Signature of Approving Official:

Date:

6-7-05



EPA #538-GNN



2/12

FRONT

SCOTTS®
Granular MSM

NOTE TO PM: OPTIONAL FRONT PANEL CLAIMS

[Claims regarding "Problem Control"]


- Kills/controls/eliminates/eradicates/treats weeds from your whole lawn, unlike spot treatment sprays.
- Kills/controls/eliminates/eradicates/treats weeds spot treatment sprays miss.
- Stop weeds in their tracks.
- Make weeds/dollarweed disappear/history.
- Say goodbye to weeds/dollarweed
- Get the upper hand on tough [summer] weeds/dollarweed/dandelions/clover.
- [Summer] weed control for broadleaf weeds.
- Kills/controls/eliminates/eradicates/treats dollarweed [and] [over 50] other [common] [lawn] weeds [as listed].
- Kills/Controls/Eliminates/Eradicates/Treats the most [majority of] common lawn weeds (as listed)
- Kills/controls/eliminates/eradicates/treats weeds/dollarweed/dandelions/clover [in lawns]
- Kills/controls/eliminates/eradicates/treats weeds/dollarweed/dandelions/clover and [many/other] [common] lawn weeds.(as listed)
- Eliminates/Eradicates/Treats/Cures your lawn of tough [summer] weeds.
- Kills/Controls/Eliminates/Eradicates/Treats/Clears out [tough] [existing] weeds.
- Kills/Controls/Eliminates/Eradicates/Treats/Clears out embarrassing lawn weeds.
- Corrects/Controls/Eliminates/Eradicates/Treats most/many lawn weed problems

[Claims regarding "How it works"]

- Scotts weed killer [killing technology].
- With [to be named], Scotts weed killing technology.
- Contains a granular/proven formula that attacks weeds through the leaves killing them while leaving your grass [St. Augustine/bermudagrass/zoysia Lawn) completely unharmed – guaranteed.
- Contains [a proven] formula that targets/kills/eliminates/controls/destroys weeds.
- Works through the leaves and roots of the weeds.
- Small, lightweight granules provide complete uniform coverage and are quickly absorbed to help ensure complete control of weeds.
- [Contains a formula that] works through the leaves and roots of the weeds.

[Claims regarding "Where to use/Grass types"]

- For use on [all] lawns [and] /grass types.
- Won't harm your lawn.
- For use on St. Augustinegrass (including Floratam) bermudagrass, zoysia (Meyers & Emerald) and buffalograss lawns.
- Made (formulated) (specially formulated) for [use on] St. Augustine, bermudagrass, zoysia and buffalograss lawns.
- For/Made for/Specially formulated for/Formulated for [use on] St. Augustinegrass [(including Floratam)], bermudagrass, zoysia and buffalograss lawns.



- Treats [and feeds] bermudagrass [too]!
- Now a [St. Augustinegrass/bermudagrass/zoysia] weed control [formula].
- Specially formulated for buffalograss
- Now for use on Bermuda and buffalograss.
- Kills weeds, not St. Augustine/bermudagrass lawns.
- Won't harm St. Augustine/Southern/bermudagrass lawns/grass)!
- Specially formulated to kill weeds, won't harm lawns [grass].
- Targets/Kills/seek out and destroys weeds, while leaving the [your] grass [lawn/St. Augustine/bermudagrass/zoysia lawn] unharmed – guaranteed.
- Targets/Kills/seek out and destroys weeds, leaves grass (lawn) unharmed.
- Apply to wet grass.

[Claims regarding "Positioning"]

- Scotts best.
- Our/Scotts best weed killing formula.

[Claims regarding "Product Form"]

- Easy to spread [granules].
- Ready to Use [granules].

[Claims regarding "Coverage"]

- Treats/Covers [up to] 5,000 sq. ft.

[Claims regarding "Application Timing"]

- Apply/Use anytime weeds are [actively] growing/active/present/visible.
- Apply/Use in the spring, summer or fall or anytime weeds are growing/active/present/visible [actively growing].

[Claims regarding "Guarantee"]

- Guaranteed [performance/results].
- Scotts® No-Quibble Guarantee®

[Claims regarding "Cross-Sell"]

- For a beautiful lawn, apply [a] Scotts fertilizer [product] four times a year [every six to eight weeks].

NOTE TO PM: END OF OPTIONAL FRONT PANEL CLAIMS

NET WEIGHT 11.50 LBS

XX-X-X

KEEP OUT OF REACH OF CHILDREN
CAUTION
See back for PRECAUTIONARY STATEMENTS

ACTIVE INGREDIENT:

Metsulfuron-Methyl†	0.03%
OTHER INGREDIENTS:	99.97%
Total	100.00%

†CAS #74223-64-6

ACCEPTED

JUN - 7 2005

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended for the pesticide registered under EPA Reg. No. 538-300

BACK

PULL TAB
and remove for
pour spout
(ILLUS.)

XXXXX
XM


TREATS XX,XXX SQ FT

PROOF OF
PURCHASE

Easy
Open PULL HERE -

Cut Here To Open
(Scissors Illust)

NOTE TO PM: OPTIONAL BACK PANEL MARKETING CLAIMS

Product Facts	
Treats up to XXXX sq ft	
KILLS WEEDS	Dollarweed and over 50 other [common] lawn weeds
WHERE TO USE	For use on St. Augustinegrass (including Floratam), bermudagrass, zoysia (Meyers & Emerald) and buffalograss lawns [only] - won't harm lawn grasses
AMOUNT TO USE	Sprinkle product lightly over weeds. 5.43 grams (X tbsp) covers 4 sq ft. Spreader instructions
 Questions, Comments or Medical Information call 1-800-225-2883 (1-800-543-TURF) www.ortho.com (www.scotts.com)	

WEEDS CONTROLLED/CONTROLS THESE COMMON LAWN WEEDS:

Grasses: bahiagrass, ryegrass

Broadleaves: Annual blueeyedgrass, annual sowthistle, aster, bittercress, blue mustard, Brazil & Florida pusley, bur buttercup, Carolina geranium, carpetweed, chamberbitter, chickweed, clover (white & sweet), common groundsel, common mullein, common purslane, common yarrow, creeping beggarweed, creeping indigo, crown vetch, cudweed, curly dock, cutleaf eveningprimrose, dandelion, dollarweed, dogfennel, field pennycress, Florida betony, ground ivy, henbit, lambsquarters, lawn burweed, lespedeza, matchweed, miners lettuce, old world diamond-flower, parsley-piert, Pennsylvania smartweed, pigweed (redroot & smooth), plantains, prickly lettuce, prostrate knotweed, ragweed, redstem (filaree), shepherdspurse, smallseed falseflax, spurge, spurweed, tansy mustard, treacle mustard, tumble mustard, wild carrot, wild celery, wild garlic, wild lettuce, wild mustard, wild onion, and woodsorrel (oxalis).

Scotts Tips

The best weed prevention begins with a complete lawn care program. Tips for a weed-free lawn:

1. Feed your lawn regularly using Scotts Annual Lawn Care Program. It provides the correct balance of nutrients to help the lawn crowd out weeds, making it difficult for them to establish.
2. Mow your lawn at the recommended height. Mowing too short weakens the lawn and allows sun to reach weed seeds.
3. Make sure the lawn gets 1-inch per week of water. Weeds thrive in drought conditions, and steal water and nutrients from the grass.

Controls/Kills**Grasses:**

Bahigrass, ryegrass

Broadleaves

annual blue-eyed
grass
annual sowthistle
aster
bittercress
black medic
blue mustard
Brazil & Florida
Pusley
broadleaf plantain
buckhorn plantain
bushy buttonweed
bur buttercup
Canada thistle
Carolina mullein
catsear, common
chickweed, common
creeping
beggarweed
creeping indigo
crown vetch
cudweed
curly dock
cutleaf evening
primrose
dandelion, common
dollarweed
dogfennel
eclipta

English daisy
Evening primrose
field pennycress
field birdweed
field violet
Florida betony
Florida pusley
ground ivy
groundsel, common
henbit
hop clover
Jersey cudweed
lambquarters,
common
lawn burweed
lespedeza
matchweed
miners lettuce
mouse-ear
chickweed
nimblewill
old world diamond-
flower
parsley-piert
Pennsylvania
smartweed
pigweed (redroot &
smooth)
plantains

puncturevine
prickly lettuce
prostrate knotweed
purslane, common
ragweed
redroot pigweed
redstem (filaree)
shepherdspurse
smallseed falseflax
spotted spurge
spurge
spurweed
tansy mustard
treacle mustard
tumble mustard
Virginia
buttonweed
white clover
wild carrot
wild celery
wild garlic
wild lettuce
wild mustard
wild onion
yarrow, common
yellow woodsorrel
(oxalis)

What to expect

- Weeds in treated areas will stop growing 2 – 3 days after application.
- Affected weeds will gradually die and disappear with 3 – 4 weeks after application.

Where to use

- For use on St. Augustinegrass (including Floratam), bermudagrass, zoysia (Meyers & Emerald), buffalograss, and centipedegrass lawns only.
- Do not apply to lawns or mixed lawns of bahigrass, bluegrass, fine fescue, bentgrass, ryegrass, tall fescue or dichondra because injury may occur.

Coverage

- Covers X,XXX sq. ft.

NOTE TO PM: END OF OPTIONAL BACK PANEL MARKETING CLAIMS

Note to PM: Direction for Use Option 1

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling

When to use/apply

- Apply anytime weeds are actively growing in spring, summer, or fall.
- Apply in the fall to control winter weeds such as henbit, chickweed, cudweed and evening primrose.
- This product applies 0.03 lbs active ingredient per acre.

Reapplication

- For severe weed problems, a second treatment may be required.
- Allow at least two-months after the first application.

[Optional] With other products

- Does not interfere with the performance of other lawn products such as fertilizers, weed controls and insect controls.
- Allow at least one-week between product applications.

With new lawns or seeding/When seeding, sprigging, or sodding

- Can be applied after sprigging/sodding a new St. Augustine, bermuda, or zoysia lawn to prevent competition from newly emerging weeds. Water the new sprigs/sod and then make the application.
- Don't plant sprigs for one month, or sow new grass seed for two months following application.

[Optional] Reentry

For best results, avoid disturbing particles by keeping children and pets off of treated area until product has been watered in per instructions and foliage is dry.

How to apply

- Apply to wet grass blades. Do not apply if rain is expected within 24-hours
 - Apply with a drop or rotary spreader. Do not apply by hand or hand-held spreader.
- [Optional] For best results, use a Scotts Spreader.

[Optional]	[Optional]
Step	Action

- | | |
|---|--|
| 1 | Mow Lawn <ul style="list-style-type: none"> • If grass is tall enough to touch bottom of spreader, mow before application. • Mow tall grass a day or two before application |
| 2 | Water Lawn <ul style="list-style-type: none"> • Grass blades must be wet for product to work properly. • Apply when the grass is wet from dew or rainfall. Water lawn lightly if grass blades are dry. |
| 3 | Set Spreader |
| 4 | Fill spreader <ul style="list-style-type: none"> • Fill spreader over bare ground or hard surface. • Sweep or brush any particles off of hard surfaces and apply on lawn or dispose of properly. |
| 5 | Apply Product <ul style="list-style-type: none"> • Apply one or two header strips, then apply product in longest direction of lawn. • Shut off spreader each time you reach the header strip, or stop. |
| 6 | DO NOT Water <ul style="list-style-type: none"> • Do not water lawn for 24-hours following application. |

Note to PM: End of Option 1

Note to PM: Directions for Use Option 2

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling

When to use/apply

- Apply anytime weeds are actively growing in spring, summer, or fall.
- Apply in the fall to control winter weeds such as henbit, chickweed and evening primrose.
- This product applies 0.03 lbs active ingredient per acre.

Reapplication

- For severe weed problems, a second treatment may be required.
- Allow at least two-months after the first application.

[Optional] With other products

- Does not interfere with the performance of other lawn products such as fertilizers, weed controls and insect controls.
- Allow at least one-week between product applications.

With new lawns or seeding/When seeding, sprigging, or sodding

- Can be applied after sprigging/sodding a new St. Augustine, bermuda, or zoysia lawn to prevent competition from newly emerging weeds. Water the new sprigs/sod and then make the application.
- Don't plant sprigs for one month, or sow new grass seed for two months following application.

[Optional] Reentry

For best results, avoid disturbing particles by keeping children and pets off of treated area until product has been watered in per instructions and foliage is dry.

How to apply

- Apply to wet grass blades. Do not apply if rain is expected within 24-hours
 - Apply with a drop or rotary spreader. Do not apply by hand or hand-held spreader.
[Optional] For best results, use a Scotts Spreader.
1. If grass is tall enough to touch bottom of spreader, mow before application. Mow tall grass a day or two before application.
 2. Grass blades must be wet for product to work properly. Apply when the grass is wet from dew or rainfall. Water lawn lightly if grass blades are dry.
 3. Set Spreader
 4. Fill spreader over bare ground or hard surface. Sweep or brush any particles off of hard surfaces and apply on lawn or dispose of properly.
 5. Apply one or two header strips, then apply product in longest direction of lawn. Shut off spreader each time you reach the header strip, or stop.
 6. **DO NOT** water lawn for 24-hours following application.

Note to PM: End of Option 2

Note to PM: Directions for Use Option 3

Directions for use

It is a violation of federal law to use this product in a manner inconsistent with its labeling

- **Do not apply to lawns or mixed lawns of bahiagrass, centipedegrass, bluegrass, fine fescue, bentgrass, ryegrass, tall fescue or dichondra because injury may occur.**
- For best results, apply to wet grass.
- Do not apply on or under the branch spread (rootzone) area of trees, shrubs, bedding plants, flowers or garden plants, particularly in compact, slowly draining soils.
- This product applies 0.03 lbs active ingredient per acre.

Control

- Apply anytime weeds are actively growing. For severe weed problems, a second treatment may be required 2 months after the first.
- Apply in the fall to control winter weeds such as henbit, chickweed and eveningprimrose.

Seeding/sodding/sprigging

- Apply after sprigging/sodding a new St. Augustine, Bermuda, or zoysia lawn to prevent competition from germinating newly emerging weeds. Water the new sprigs/sod and then make the application of Bonus M.
- Do not plant sprigs/sod for one month or sow new grass seed for 2 months following application.

Note to PM: End of Option 3

Note to PM: Directions for Use Option 4

- For best results, apply to wet grass.
- Don't apply to lawns or mixed lawns of bahiagrass, centipedegrass, bluegrass, fine fescue, bentgrass, ryegrass, tall fescue or dichondra because injury may occur.
- Don't plant sprigs for one month, or sow new grass seed for two months following application.
- **DESIRABLE PLANTS: DON'T APPLY ON OR UNDER THE BRANCH SPREAD (ROOTZONE) AREA OF TREES, SHRUBS, BEDDING PLANTS, FLOWERS OR GARDEN PLANTS, PARTICULARLY IN COMPACT, SLOWLY DRAINING SOILS.**
- An application of Scotts Lawn Supplement may be used anytime during the year if extra greening is desired.
- This product applies 0.03 lbs active ingredient per acre.

Note to PM: End of Option 4

OPTIONAL DIRECTIONS FOR USE Option 5 – For Shaker/Canister/ Bottle Applicator

IMPORTANT

Do not apply to lawns or mixed lawns of bahiagrass, centipedegrass, bluegrass, fine fescue, bentgrass, ryegrass, tall fescue or dichondra because injury may occur.

Do not plant sprigs for one month, or sow new grass seed for two months in treated area following application.

Desirable plants: Do not apply on or under the branch spread (rootzone) area of trees, shrubs, bedding plants, flowers or garden plants, particularly in compact, slowly draining soils.

FOR BEST RESULTS

AMOUNT TO USE *[Image/illustration of amount to use]*

Using a quarter as a size reference, this illustration shows the amount of *(Name to be determined)* with grams (5.43 tbs) sprinkled evenly over 4 sq ft (square feet). This product applies 0.03 lbs active ingredient per acre.

WHERE TO USE

Use to control weeds on St. Augustine (including Floratam), bermudagrass, zoysia (Meyers & Emerald), buffalograss, and carpetgrass lawns only.

WHEN TO APPLY

Apply anytime weeds are actively growing.

Optional - Ideal for spot treating lawn weeds in between fertilizer [and granular weed & feed] (Scotts Bonus M®) applications.

For best results, apply to wet grass.

HOW TO APPLY Spreader instructions

——SHAKER BAG, CANISTER, SHAKER BOTTLE——

Open container and shake as needed to lightly sprinkle product over weeds. Cover as indicated in the picture above (below). 5.43 grams (X tbs) treats 4 square feet (sq ft).

[OR]

——GRASSHOPPER APPLICATOR ——

Easy to Use Applicator

1. Remove top overcap. Unscrew nozzle and remove safety seal from jug. [illustration #1]
2. Extend nozzle applicator and rotate as shown [illustration #2]
3. Sprinkle product over weeds in lawn. Cover as indicated in above instructions. 5.43 grams (X tbs) treats 4 square feet (sq ft) [illustration #3]

END OF OPTIONAL DIRECTIONS FOR USE Option 5– For Shaker/Canister/ Bottle Applicator

**KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS**

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye irritation. Avoid contact with eyes and clothing.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye.

Call poison control center or doctor for further treatment advice. Have the product container or label with you when calling poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

STORAGE & DISPOSAL

Pesticide Storage: Store in a clean, dry place. Reseal opened bag [by folding top down and securing].

Pesticide disposal: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions.

Never place unused product down any indoor or outdoor drain.

Container disposal: Do not reuse bag. Discard bag in trash or offer for recycling if available.

Spreader Setting

For best results, use a Scotts spreader.

Apply header strips. Then apply product in longest directions of lawn.

[ART]

		Directions
Scotts AccuGreen® Drop Spreaders	X	Mow lawn; apply 2 header strips; apply longest direction of lawn, overlapping wheel marks.
Scotts PrecisionGreen® Drop Spreader	X	Mow lawn; apply 2 header strips; apply longest direction of lawn, overlapping wheel marks.
Scotts SpeedyGreen & EdgeGuard™ Rotary Spreader	X	Make each pass about 5 feet from previous pass to overlap spread width.
Scotts EasyGreen® Rotary Spreader	X	Make each pass about 5 feet from previous pass to overlap spread width.

Additional Spreader Settings:

EZ Drop, Republic (#530)	X
EZ Broadcast, Republic (#540)	X
Scotts Broadcast Tractor Spreader	X

(Non-Scotts spreader settings are approximate. You may need to adjust for proper coverage.)

[OR]

Spreader	Setting Covers x,xxx sq. ft.	Coverage Tip
Rotary Spreaders		

Pending EPA Approval

11/12

Scotts SpeedyGreen® (1000, 2000, 3000 series) & EdgeGuard™	X	Make each pass about 5 feet from the previous pass.
Scotts EasyGreen®	X	Make each pass about 30 inches from the previous pass.
Scotts Tractor Spreader	X	
Drop Spreaders		
Scotts Accugreen® (1000, 2000, 3000 series)	X	Overlap wheel marks to ensure uniform coverage.
Scotts PrecisionGreen®	X	

Product of USA

U.S. Patent No. XXXXXXXXXXXXXXX

EPA Reg. No. 538-xxx

EPA Est. Nos. 538-OH-1(S)

Superscript used is first letter of run code on end of container.

If you have questions

We designed this Program to make having a better lawn as simple and easy as possible. But you may still have some questions about your lawn during the year. If so, you can contact the Scotts LawnPro retailer where you bought this Program. They are ready to answer your questions and provide any help or information that you may need. -OR- You may get the answers by talking with one of Scotts lawn counselors. Simply dial this toll-free number 1-800-543-TURF (Monday through Friday 9:00 AM to 5:00 PM Eastern time (in Ohio 1-800-762-4010.)

Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied with results after using the product, you are entitled to get your money back. Simply send us evidence of purchase and we will mail you a refund check promptly.

NOTICE: Buyer assumes all risks of use, storage or handling of this product not in accordance with directions.

The ORTHO Guarantee

If for any reason you are not satisfied with this product, mail us proof of purchase to obtain a full refund of your purchase price.

Dedicated To A Beautiful World™

Trusted by generations of homeowners, Scotts is dedicated to helping you enhance and protect the natural beauty of your home environment
Scotts®

Questions / Information

Consumer Helpline
1-800-543-TURF(8873)

Visit our website
www.scotts.com

[Illus]SMG listed NYSE

The Scotts Company
14111 Scottslawn Road
Marysville, Ohio 43041

The Ortho Group
P.O. Box 1749
Columbus, OH 43216

12/12

Pending EPA Approval

SCOTTS.....the lawn people
MARYSVILLE, OHIO 40341

©2004, The Scotts Company. World rights reserved.