

538-316

02/12/2010

1/15



U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs
Registration Division (7505P)
Ariel Rios Building
1200 Pennsylvania Ave., NW
Washington, D.C. 20460

EPA Reg. Number:

538-316

Date of Issuance:

FEB 12 2010

NOTICE OF PESTICIDE:

- Registration
 - Reregistration
- (under FIFRA, as amended)

Term of Issuance: unconditional

Name of Pesticide Product:

Scotts Lawnservice Turf Fertilizer
Plus 0.28% Halts Pro

Name and Address of Registrant (include ZIP Code):

The Scotts Company
14111 Scottslawn Rd.
Marysville, OH 43041

~~Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.~~

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

The basic formulation dated 8/18/09 and alternate formulation 'A' dated 1/7/10 are acceptable.

This product is registered in accordance with FIFRA provided that you:

1. Submit and/or cite all data required for registration review/reregistration of your product when the Agency requires all registrants of similar products to submit data.
 - a. One year storage stability (830.6317) and Corrosion Characteristics (830.6320) studies must be generated and submitted to the Agency. It is recommended that the observations must be made at 0, 3, 6, 9, & 12 month intervals. Please submit the results electronically if possible along with the hard copy.
2. Make the following label revisions:
 - a. Revise "EPA REG. NO. 538-NEW" to "EPA REG. NO. 538-316"
 - b. Add a descriptor below the product name indicating that "Pro" refers to the active ingredient, prodiamine, and not professional grade.

Continued on Page 2

Signature of Approving Official:

Kathryn V. Montague
Project Manager 23
Herbicide Branch
Registration Division (7505P)

Date:

FEB 12 2010

c. Revise the Environmental Hazards statement to read as follows:

“This pesticide may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters or rinsate.”

d. Replace the word, “recommended” with “listed” in the sentence, “Sequential application may be made as long as the total amount of product applied does not exceed the maximum annual application rates *recommended* for each turf species.”

3. Submit one (1) copy of the revised final printed label for the record.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

NOTE:

Bold, italicized text is information for the reader and is not part of the label.
[Bracketed information is optional text]. Text separated by / denotes and/or options.
Underline text is new. ~~Strike-through text is being deleted.~~

**ACCEPTED
with COMMENTS
In EPA Letter Dated:**


FEB 12 2010
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

538-316

[For preemergence control of grass and broadleaf weeds in established turf grasses (excluding golf course putting greens) and lawns.]

[Landscape ornamentals, established perennials and wildflower plantings]

SCOTTS LAWN SERVICE TURF FERTILIZER PLUS .28% HALTS PRO



ACTIVE INGREDIENT

Prodiamine: [N³,N³-Di-n-propyl-2, 4-dinitro-6-(trifluoromethyl)-m-phenylenediamine] 0.28%

OTHER INGREDIENTS.....99.72%

TOTAL 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See back / side / top / bottom [panel / label] for additional precautionary statements.

NET WEIGHT: 50 POUNDS (22.7 kg.)

4/15

NOTE:

EPA Reg. No. 538-NEW

Bold, italicized text is information for the reader and is not part of the label.
[Bracketed information is optional text]. Text separated by / denotes and/or options.
Underline text is new. ~~Strike through text is being deleted.~~

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals.

FIRST AID

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for further treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for further treatment advice.

IF SWALLOWED: Call a poison control center or physician immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or physician. Do not give anything by mouth to an unconscious person.

IF INHALED: Move person to fresh air. If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container with you when calling a poison control center or doctor, or going for treatment. Questions? Call 1-877-561-6584.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Application Precautions

Not for use on turf or plants being grown for sale or other commercial use, for commercial seed production, or for research purposes. For use on turf or plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses or lawns and grounds.

NOTE: Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

NOTE:

Bold, italicized text is information for the reader and is not part of the label.
[Bracketed information is optional text]. Text separated by / denotes and/or options.
Underline text is new. ~~Strike through text is being deleted.~~

How it Works

This product is a turf fertilizer and a selective pre-emergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses (excluding golf course putting greens), and lawns, landscape ornamentals and established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seeds, germination and root development.

ESTABLISHED TURF

Application Directions: Apply this product uniformly, using a suitable spreader that has been properly calibrated. Most effective weed control will be obtained when this product is activated by at least 1/2 inch of rainfall or irrigation prior to weed seed germination and within 14 days of application.

Use Precautions: The following precautions apply to the use of this product in turf grasses and lawns.

1. Application of this product may thin emerged annual bluegrass and newly overseeded grasses.
2. Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil (not thatch plus soil).
3. Do not cut (harvest) treated sod before 120 days after application. Do not apply newly set sod until the following year.
4. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury.
5. Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
6. Do not apply to putting greens or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

Rates of Application: This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential application may be made as long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. All applications must be made prior to germination of the target weeds. This product will not control established weeds. Maximum use rate selection should be based on turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

Maximum Annual Rates: This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not apply more than 535 lbs. of this product per calendar year. Use higher rates of product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum yearly rate listed.

NOTE:

Bold, italicized text is information for the reader and is not part of the label.
 [Bracketed information is optional text]. Text separated by / denotes and/or options.
Underline text is new. ~~Strike-through text is being deleted.~~

MAXIMUM APPLICATION RATE/CALENDAR YEAR (by Turf Grass Species)		
Turf Species	Lbs. Product/Acre	Lbs. a.i./Acre
Creeping Bentgrass	233 lbs.	0.65 lb.
Creeping Red Fescue	267 lbs.	0.75 lb.
Buffalograss Kentucky Bluegrass Perennial Ryegrass	357 lbs.	1.00 lb.
Bahiagrass ¹ Bermudagrass ² Centipedegrass ¹ Seashore Paspalum ¹ St. Augustinegrass Tall Fescue (including turf-Type) Zoysiagrass	535 lbs.	1.50 lb.

¹These are the maximum rates per calendar year.

²May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 180 pounds/acre. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

When To Apply After Overseeding Turf: Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

When To Overseed After Application: This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/reseeding results.

Product Rate Lbs./Acre		Months Before Overwintering		
Lbs. Product/Acre	Lbs. a.i./Acre	North	Transition	South
180	0.50	4	4	4
233	0.65	5	4	4
267	0.75	6	5	5
285	0.80	-	6	6
357	1.00	-	7	7
407	1.14	-	-	9
465	1.30	-	-	10
535	1.50	-	-	12

Application Directions: Apply uniformly with suitable, calibrated application equipment. Calibrate your spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended application rate under actual application conditions. Apply this product uniformly over the area to be treated. Avoid streaking, skips and overlaps during application. Calibration of spreader may be needed if conditions of area to be treated changes and rate or speed of application are changed.

NOTE:

Bold, italicized text is information for the reader and is not part of the label.
[Bracketed information is optional text]. Text separated by / denotes and/or options.
Underline text is new. ~~Strike through text is being deleted.~~

SPREADER RECOMENDATIONS				
Spreader Settings for 1,000 sq. ft.	180 lbs. product/A	267 lbs. product/A	357 lbs. product/A	535 lbs. product/A
Spreader A	<i>Spreader settings available at printing</i>			
Spreader B				
Spreader C				
<i>May contain additional spreader brands/models, spreader settings and spreader use instructions as appropriate for the product</i>				

LANDSCAPE ORNAMENTALS
(including established perennials and wildflower plantings)

This product may be applied for residual pre-emergence weed control in ornamentals. Apply at the rate of 5.35 to 12.28 pounds/1,000 square feet (233 to 535 pounds/acre) in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 12.28 pounds/1,000 square feet (535 pounds/acre) per year.

Application Timing And Information: This product may be applied to newly transplanted and established ornamentals as a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grafts have taken to avoid any inhibition of the tissue union. This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 1/2 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when applied to soil free of clods, weeds and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate post-emergence herbicide.

Tolerant Ornamental Species: This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

SCIENTIFIC NAME

- Abies spp.
- Abelia grandiflora
- Acer palmatum
- A. Platanoides
- Achillea spp.
- Actindia chinensis
- Agapanthus africanus
- Agapanthus orientalis
- Akebiaq quintata
- Allium cemuum

COMMON NAME

- Fir species** (Balsam, Frasier, Noble, etc.)
- Abelia: Sherwood
- Japanese Maple
- Norway Maple
- Yarrow: King Edward
- Kiwi*
- Lily-of-the-Nile (African Lily)

- Five-leaf or Chocolate Vine
- Lady's Leek, Nodding Onion

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by / denotes and/or options.

Underline text is new. ~~Strike through text is being deleted.~~

Anemone hybrida	Japanese Anemone
Aquilegia spp.	Aquilegia: Red and Gold
Arctostaphylos densiflora	Vine Hill Manzanita
Arctotheca calendula	Cape Weed
Artemesia spp.	Wormwood; Silver Mound, Castle
Aster spp.	Aster: Bonnie Blue, Purple Dome
Aster X Frikartii	
Athrium Filiz-femina	Lady Fern; Fern Lady
Aucuba japonica	Japanese Aucuba
Begonia spp.	Fibrous Begonia: Hardy Grandis
Berberis gladwynensis	Barberry
B. Julianae	Wintergreen Barberry
B. mentorensis	Mentor Barberry
B. Thunbergil	Japanese Barberry
B. Verruculosa	Warty Barberry
Bergenia cordifolia	
Boltonia asteroides	Snowbank
Buddleia davidii	Butterfly-Bush (Dwarf Blue); Royal Red
Buxus microphylla	Japanese Boxwood
Callistemon viminalis	Weeping Bottlebrush
Calluna vulgaris	Scotch Heather
Campanula carpatica	Tussock Bellflower, (White Clips)
Campis X Tagliabuana	Trumpet Flower, Madame Galen
Carpobrotus edulis	Hottentot Fig (Ice Plant)
Cassia artemisoides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Ceratostigma plumbagonoides	
Chamaecyparis pisifera	False Cypress
Cleyera japonica	Cleyera
Citrus spp.	Citrus species*
Coreopsis spp.	Coreopsis (Calliopsis) Early Sunrise, Moonbeam
Cornus florida	Flowering Dogwood
C. Stolonifera	American Dogwood
Cortaderia seloana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. dammeri	Bearberry Cotoneaster
C. Microphyllus	Rockspray Cotoneaster
Cretaegus spp.	Hawthorne
Crocosmia spp.	Lucifer
Cupressus sempervirens	Italian Cypress
Delosperma alba	White Trailing Ice Plant
Delosperma spp.	Cooperi Pink
Delphinium spp.	Larkspur: Blue Elf
Dianthus deltoides	Dianthus: Maiden Pinks "Zing"
D. gratianopolitanus	Cheddar Pink
Dodonea viscosa	Hop Bush
Echinacea purpurea	Coneflower, Purple; Magnus

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by / denotes and/or options.

Underline text is new. ~~Strike through text is being deleted.~~

- Elaeagnus pungens
- Euonymus fortunei
- E. japonica
- E. kiautschovica
- Fatsia japonica
- Forsythia intermedia
- Forsythia suspensa
- F. viridissima
- Gaillardia spp.
- Gardenia jasminoides
- Gaura spp.
- Gentiana dahurica
- Geranium cinereum
- Gladiolus spp.
- Gypsophilia repens
- Hedera helix
- Hellanthemum spp.
- Hemerocallis spp.
- Hibiscus
- Hibiscus spp.
- Hibiscus rosa-sinensis
- Hosta plantaginea
- H. Sieboldiana
- Houttuynia cordata var. variegata
- Hydrangea macrophylla
- Ilex Bernyi
- Ilex cornuta
- I. crenata
- I. opaca
- I. vomitoria
- Inula ensifolia
- Iris ensata
- Iris siberica
- Iris spp.
- Jasminium nudiflorum
- Juniperus chinensis
- J. conferta
- J. davurica
- J. horizontalis
- Juglans spp.
- Justicia brandegeana
- Lagerstromia indica
- Lagerstromia indica X Fauriei
- Lantana montevidensis
- Lavender spp.
- Leonopodium alpinum
- Ligustrum amurense
- Ligustrum sinense
- Silverberry
- Wintercreeper
- Evergreen Euonymus
- Spreading Euonymus
- Japanese Aralia
- Border Forsythia
- Weeping Forsythia
- Greenstem Forsythia
- Gaillardia, Blanket Flower: "Goblin"
- Gardenia, Cape-Jasmine
- Gentian
- Cranesbill
- Gladiolus species**
- Baby's Breath
- English Ivy
- Sunrose
- Daylily: Aztec Gold, Stella de Oro, Tender Love
- Rose of Sharon**
- Mallow: Disco Belle White
- Chinese Hibiscus
- Hosta, Plantain Lily (Fragrant)
- Hosta, "Seersucker"
- Bigleaf Hydrangea
- Holly
- Chinese Holly
- Japanese Holly
- American Holly
- Yaupon Holly, Schillings
- Sword-Leaved Iris; Jodlesong
- Siberian Iris; Cabernet
- Iris species**
- Winter Jasmine
- Chinese Juniper; Nick's Compact, Parsonnii
- Shore Juniper; Blue Pacific
- Parsoni
- Creeping Juniper
- Walnut*
- Shrimp Plant
- Crape Myrtle
- Crape Myrtle; Tuscarora
- Weeping Lantana
- Lavender; Munstead
- Edelweiss
- Amur. Privet
- Chinese Privet; Variegata

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by / denotes and/or options.

Underline text is new. ~~Strike through text is being deleted.~~

- L. japonicum
- L. lucidum
- Lilium spp.
- Liriope muscari
- Liriope muscari var. variegata
- Liriope spicata
- Lobelia Cardinalis
- Lonicera japonica
- L. tatarica
- Lorapetalum chinense
- Lythrum spp.
- Magnolia spp.
- Maleophora luteola
- Malus spp.
- Miscanthus sinensis
- Nandina domestica
- Narcissus spp.
- Nerium oleander
- Olea europaea
- Ophiopogon japonicus
- Osmanthus heterophyllus
- Osteospermum fruticosum
- Oxydendrum arboreum
- Paeonia suffruticosa
- Pennisetum setaceum
- Perovskia atriplicifolia
- Persea americana
- Photinia Fraseri
- Physotegia Virginiana
- Picea spp.
- Pieris japonica
- Pinus brutia
- P. canariensis
- P. Elliottii
- P. halepensis
- P. nigra
- P. palustris
- P. radiata
- P. strobus
- P. sylvestris
- P. taeda
- P. Thunbergiana
- P. Virginiana
- Pistachio spp.
- Pittosporum rhombifolium
- P. tobira
- Podocarpus macrophyllus
- Prunus laurocerasus

- Japanese Privet
- Glossy Privet (Waxleaf)
- Lily; Jazz
- Big Blue Lily, Silver Mound, Evergreen Giant
- Liriope, Variegated
- Liriope, Creeping
- Cardinal Flower, Indian Pink
- Japanese Honeysuckle
- Tatarian Honeysuckle
- Burgundy
- Loosestrife; Modern Pink
- Magnolia species**
- Ice Plant
- Crabapple*
- Yaku Jima, Silberfeder
- Heavenly Bamboo
- Narcissus species**
- Oleander
- Olive*
- Mondo Grass
- Osmanthus (False Holly); Gulf Tide
- Trailing African Daisy
- Sourwood
- Tree Peony
- Fountain Grass (Dwarf)

- Avocado*
- Fraser's Photinia (Redtip)
- Dragonhead, False; Vivid
- Spruce species** (Colorado Blue, Norway, etc.)
- Lily-of-the-Valley Bush
- Calabrian Pine
- Canary Island Pine
- Slash Pine
- Aleppo Pine
- Austrian Pine
- Longleaf Pine
- Monterey Pine
- Eastern White Pine
- Scotch Pine
- Loblolly Pine
- Japanese Black Pine
- Virginia Pine
- Pistachio*
- Queensland Pittosporum
- Japanese Pittosporum
- Japanese Yew
- English Laurel

NOTE:

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by / denotes and/or options.

Underline text is new. ~~Strike-through text is being deleted.~~

Prunus spp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
Pseudotsuga menziesii	Douglas Fir**
Pyracantha coccinea	Firethorn (Scarlet)
P. fortuneana	Firethorn
P. koidzumii	Firethorn
Pyrus spp.	Bradford Pear spp.
Quercus Shumardii	Oak, Shumard's Red
Quercus spp.	Oak species
Raphiolepis indica	Indian Hawthorne
Raphiolepis umbellata	Yedda Hawthorne
Rhododendron	'Coral Bells,' Formosa,' Hino-crimson," PJM'
(including Azalea)	'Delaware Valley White', ' Roseum Elegans,'
	'Flame Creeper', Girard Crimson', 'Wakeilebisu
	'White Gumbo', 'George L. Tabor'
Rosa banksiae	Lady Bank's Rose
Rosmarinus officinalis	Rosemary*
Rudbeckia spp.	Black-Eyed Susan; Goldsturm
Santolina virens	Santolina
Saxifraga spp.	Saxifrage; Purple Dome
Scabiosa spp.	Pincushion Flower
Sedum album	Stonecrop
Sedum caudicicola	Stonecrop; Lidakense
Sedum dasyphyllum	Stonecrop
Sedum spurium	Stonecrop; Dragon's Blood
Spiraea bumalda	Spirea; Anthony Waterer
Syzygium paniculatum	Australian Brushcherry
Taxus cuspidata	Japanese Yew
T. media	Yew
Teucrium spp.	Germander
Thalictrum dipterocarpum	Meadow Rue
Thuja occidentalis	American Arborvitae
Trachelospermum asiaticum	Star Jasmine
Tsuga canadensis	Canada Hemlock
Tulipa spp.	Tulip species
Veronica spp.	Veronica, Speedwell; Sunny Border
Viburnum japonicum	Japanese Viburnum
V. odoratissimum	Sweet Viburnum
V. plicatum	Japanese Snowball
V. rigidum	Canary Island Viburnum
V. suspensum	Arrowood Viburnum
V. tinus	Laurustinus
V. trilobium	Cranberry Bush
V. wrightii	Leatherleaf Viburnum
Vinca major	Greater Periwinkle, Vinca
Vinca Minor	Common Periwinkle, Vinca
Vitis spp.	Grape*
Weigela florida	Old Fashioned Weigela

12/15

NOTE:

EPA Reg. No. 538-NEW

Bold, italicized text is information for the reader and is not part of the label.
[Bracketed information is optional text]. Text separated by / denotes and/or options.
Underline text is new. ~~Strike through text is being deleted.~~

Yucca aloifolia
Y. filametosa
Yucca

Spanish Bayonet, Yucca
Adam's Needle,

*Do not use on food producing trees, vines or plants.
**Not for use on container grown plants.

WEEDS CONTROLLED

When used in accordance with label directions, this product will provide control of the following weeds:


Barnyardgrass	Henbit	Purslane, Common
Bluegrass, Annual (<i>Poa annua</i>)	Itchgrass	Pusley, Florida
Carpetweed	Johnsongrass (from seed)	Rescuegrass ³
Chickweed, Common	Junglerice	Shepherd's Purse ²
Chickweed, Mouseear (from seed)	Knotweed	Signalgrass, Broadleaf
Crabgrass (large, smooth)	Kochia	Speedwell, Persian
Crowfootgrass	Lambsquarter, Common	Sprangletop
Cupgrass, Woolly	Lovegrass	Spurge, Prostrate
Foxtails, Annual	Panicum (Texas, fall, browntop)	Witchgrass
Goosegrass ¹	Pigweed	Woodsorrel, Yellow (from seed)

¹In many areas, a single application of 233 to 535 pounds/acre will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, most effective weed control may be obtained with an initial application of 233 to 357 pounds/acre, followed after 60 to 90 days by a second application at doses that would not exceed the maximum annual rate (see "Maximum Application Rate by Turfgrass Species" table in the Rates of Application section of this label).

²For this weed, apply in late summer, fall or winter prior to germination.

³Suppression only.

GERMINATION DATES



13/15

NOTE:

EPA Reg. No. 538-NEW

Bold, italicized text is information for the reader and is not part of the label.
[Bracketed information is optional text]. Text separated by / denotes and/or options.
Underline text is new. ~~Strike through text is being deleted.~~

FERTILIZER GUARANTEE INFORMATION

The following table may be used in whole or in part. Fertilizer information denoted by "X" will be added to FPL. NPK will not exceed the endpoints tested for acute toxicology studies.

GUARANTEED ANALYSIS

[Product Name] [Scotts LawnService Turf Fertilizer Plus 0.28% Halts Pro]

XX-XX-XX *insert N-P-K ratio*

FXXX *Insert FL Fertilizer License No.*

Total Nitrogen	XX%
Soluble Potash	XX%
Sulfur	XX%
XXXXXX.....	XX%

Derived from: ***insert ingredients as required to comply with State fertilizer regulations***

*Contains X.X% slowly available ***insert source of slowly available nitrogen consistent with State fertilizer regulations***

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.regulatory-info-sc.com>

Florida Applications: X annual feedings with the exception of bahiagrass and centipedegrass (X feedings in the north and X feedings in the central and south)] ***Feedings information denoted by "X" will be added to FPL. Feedings will not exceed agronomic recommendations nor RED application requirements.***

[Following are optional referral statements for Florida BMP information sources]
[In the state of Florida, it is recommended that the user of this product follow the Green Industries and Golf Course Best Management Practices Found at:
http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/BMP_Book_final.pdf
<http://www.dep.state.fl.us/water/nonpoint/docs/nonpoint/glfbmp07.pdf>]

14/15

NOTE:

EPA Reg. No. 538-NEW

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by / denotes and/or options.

Underline text is new. ~~Strike through text is being deleted.~~

STORAGE AND DISPOSAL [icon]

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE AND CONTAINER DISPOSAL Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain. **If spilled:** Sweep up and use as per label instructions or dispose of as indicated above.

OPTIONAL CLAIMS *Claims may appear on any or all label panel(s)*

- Stops many ugly weed grasses before they grow.
- Prevents crabgrass, annual bluegrass, foxtail & more.

Scotts No-Quibble Guarantee®

If for any reason you, the consumer, are not satisfied after using this product, you are entitled to get your money back. Simply send us proof of purchase and we will mail a refund check to you promptly.

Questions / Information

Consumer Helpline
1-800-543-TURF (8873)

NOTICE TO BUYER – Purchase of this material does not confer any rights under patents of countries outside of the United States.

LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as eater conditions, soil factors, off target movement, unconventional technique, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of The Scotts Company or Seller. These risks can cause: ineffectiveness of the product or other unintended consequences. **WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.**

The Scotts Company warrants that this product conforms to the chemical description on the label

15/15

NOTE:

EPA Reg. No. 538-NEW

Bold, italicized text is information for the reader and is not part of the label.

[Bracketed information is optional text]. Text separated by / denotes and/or options.

Underline text is new. ~~Strike through text is being deleted.~~

thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to inherent risks described above, when used in accordance with the Directions for Use under normal conditions.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, The SCOTTS COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL THE SCOTTS COMPANY OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF THE SCOTTS COMPANY OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHER LEGAL THEORY) RESULTING FROM THE STORAGE, USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF THE SCOTTS COMPANY OR SELLER, THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applicable law that allows such requirement, The Scotts Company or its distributor must have prompt notice of any claim so that an immediate investigation of buyer's or user's claim can be made. Buyer and all users shall promptly notify The Scotts Company or a Scott's Company distributor or any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

EPA Reg. No. 538-NEW

EPA Est. No.

[Superscript used is first letter of run code on end of container]

[Manufactured by] [Sold by] [Sold and Guaranteed by]

The Scotts Company
14111 Scottslawn Rd.
Marysville, OH 43041

© 2009. Licensed by The Scotts Company LLC.

World rights reserved

[Made in USA] [Product of USA]

[Product of Country Name] [Made in Country Name]