

DIRECTIONS FOR USE

When powdery mildew, rust, or red spider first appear and attack the plants, start dusting plants thoroughly or spray plants thoroughly with 2 table-spoonfuls in one gallon of water, or 2 lbs. of Sulphur in 32 gallons of water. Repeat application at 7 day intervals as necessary.

FOR POWDERY MILDEW ON: Roses, chrysanthemums, zinnias, crape myrtle, euonymus, hollies, sweet peas, beans, and peas.



FOR RUST ON: Roses, beans, carnations, chrysanthemums, hollyhock, snapdragons, and peas.

FOR RED SPIDER ON: Marigolds, Petunias, Roses, Verbena, Zinnias, Arbor Vitae, Boxwood, Juniper, Beans, Potatoes, Peas, Tomatoes.

DOSE RATE: Dust 40 pounds per acre; Spray 100 to 150 gallons per acre.

CAUTION: Do not use on cucurbits. Do not use if temperature is 95° F. or above.

Conditions of use are beyond seller's control. Seller's liability is limited to product's replacement or purchase price refund. Seller makes no warranty, express or implied, concerning product's use beyond claims on label.



WETTABLE DUSTING SULPHUR

CONTROLS POWDERY MILDEW, RUST, RED SPIDER ON CERTAIN FLOWERS, SHRUBS AND VEGETABLES.

ACTIVE INGREDIENTS	
Sulphur (fineness 93% thru 325 mesh)	95%
INERT INGREDIENTS	5%
Total	100%

CAUTION: KEEP OUT OF REACH OF CHILDREN.
SEE BACK FOR ADDITIONAL CAUTIONS.