

71 11516

FRUIT SPRAY

KILLS MANY INSECTS
and CONTROLS DISEASES
COMMON TO FRUIT

ACTIVE INGREDIENTS:

Captan* 16.0%
Malathion** 8.0%
Methoxychlor, Technical 16.0%

INERT INGREDIENTS:

50.0%
TOTAL 100.0%

*N-trifluoromethylthio-4-
chlorophenyl-2,2-dimethyl-1,3-dioxane-5-carboximide
**O,O-dimethyl dithiophosphate of
diethylmercaptosuccinate
***Equivalent to 11.28% of 1,2-bis-
[p-methoxyphenyl]-1,1,1-trichloro-
ethane and 1.92% other isomers
and related compounds

CAUTION

KEEP OUT OF REACH OF CHILDREN

(See side panel for additional cautions)

NET WEIGHT 12 OUNCES

E.P.A. Reg. No. 539-76



Conditions of use are beyond
seller's control. Seller's liability
is limited to product's replace-
ment or purchase price refund.
Seller makes no warranty, ex-
press or implied, concerning
product's use beyond claims on
label.

DISTRIBUTED BY SEARS, ROEBUCK AND CO.
CHICAGO, ILLINOIS 60607

HOW TO USE WITH SEARS SPRAYER:

1. Fill wettable sprays with spray powder. 2. Fill can of water.
3. Make a paste of the correct amount of powder and a small portion
of water. 4. Mix thoroughly with balance of water.
5. For best results use a Sears Sprayer, agitator sprayer which allows
to keep mixture in suspension.

Not recommended for hose end sprayers.

Spray all sides of leaves and fruit thoroughly.
Do not use sulfur with captan in the petal fall and flower sprays on
Red Delicious or other sulfur sensitive varieties. Do not use on
large pears. Some russeting may occur on Bosc pears.

CAN BE USED ON:

Fruit	Timing	Insects or Diseases
Apple and Pear	Make first application when leaves begin to show and at 7 day intervals until bloom. Additional sprays should be made at petal fall and 7 days later. Repeat at two week intervals. Do not treat within 14 days of harvest.	Aphids, Mites, Codling moth, Leafhoppers, Leafrollers, Scale crawlers, Pear slug, Scab, Black rot, Brown spot, Sooty blotch, Bitter rot.
Peach, Apricot, Cherry, Plum	Apply first spray when blossoms show pink. Spray again when petals fall. Repeat every 7 to 10 days. Do not treat within 21 days before harvest. If a later spray is needed for fruit rot, use Captan 50% wettable powder.	Cherry fruit fly, Plum curculionid, Mites, Brown rot, Leaf spot and Fruit rot.
Strawberry, Raspberry	Spray plants at 7 to 10 day intervals beginning when plants start to grow, until bloom. During bloom use only Captan 50% wettable powder. After bloom spray at 7 to 10 day intervals. Do not treat within 14 days before harvest. If a later spray is needed for fruit rot, use Captan 50% wettable powder.	Strawberry weevil, Mites, Leaf miner, Leaf spot, Fruit rot, Gray mold and Anthracnose.
Grape	Spray when new shoots are 7 to 12 inches long, pre-bloom, post-bloom. Repeat every 10 days. Do not treat within 14 days before harvest.	Grape berry moth, Japanese beetle, Leaf hopper, Leaf miner, Leaf chafer, Flea beetle and Downy mildew.
Blueberry	Spray when leaf buds are now green and beginning to open. Again when buds are about 1/2 inch in diameter. Repeat every 7 to 10 days. Do not treat within 14 days before harvest.	Fruitworm, Curculionid, Leafhopper.

CAUTION

Do not use with Captan in the petal fall and flower sprays on Red Delicious or other sulfur sensitive varieties. Do not use on large pears. Some russeting may occur on Bosc pears.

Do not reuse empty container. Destroy when empty.

NOTE: For more information on
this product, contact your
local distributor.