ACCEPTED

FRONT LABEL

NOV 4 1968

SEARS(logo) SEVIN *50W INSECTICIDE SPRAYABLE POWDER

- CONTROLS MANY INSECTS ON VEGETABLES, FRUITS, FLOWERS, SHRUBS, TREES, LAWNS
- . KILLS RESISTANT DOG TICKS, FLEAS

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-Methylcarbamate)

50.0%

INERT INGREDIENT:

50.0%

TOTAL 100.0%

*Registered Trade Mark of Union Carbide Corporation

CAUTION: Keep Out of Reach of Children. See additional cautions on back.

71 11575

Net Contents 8 ounces.

USDA Reg.No. 539-169

NOTE: This material settles after filling.

DISTRIBUTED BY SEARS, ROEBUCK AND CO., CHICAGO, ILLINOIS 60607

BACK LABEL

DIRECTIONS FOR USE

LAWN PESTS: For control of dog ticks, fleas, ants, chinch bugs, earwigs, fall armyworms, leafhoppers, sod webworms, millipedes, mosquitoes, use one level cupful in 15 to 20 gals. of water for each 500 sq. ft. of established lawn area. For best results mow lawn and apply immediately after rain or watering. For dog ticks or fleas, spray inside and outside of dog house & other areas where dogs sleep. Put fresh bedding in dog house after spraying. Repeat in 2 or 3 weeks if necessary. Spray dog at same time with 4 tbspns. per 1 gal. water.

FLOWERS, SHRUBS, EVERGREENS, SHADE TREES: To control bagworm, Birch leaf miner, box elder bug, leaf miner, blister beetle, elm leaf beetle, flea beetle, gypsy moth, inchworm(canksrworm), Japanese beetle, June beetle, lace bug, leafhopper, leaf roller, mealy bug, mimosa webworm, plant bug, psyllid, cicada, puss caterpillar, rose slug or aphid, scale insects, tent caterpillar, willow leaf beetle, thrips use 2 Tospns. per gal. water or 2 lb. in 25 gals. water, & spray plants thoroughly. Do not use on Boston ivy or Virginia creeper. Repeat weekly or as needed.

VEGETABLE Insects: Beans for bean beetles, stink bugs; cucumber & squash for melonworms, cucumber beetles, squash bugs; tomato, potato, eggplant & pepper for potato beetles, flea beetles, leafhoppers, stink bugs & grass-hoppers; cabbage & broccoli for flea beetles, harlequin bug & cabbage caterpillars, use 2 Tospns. per 1 gal. water, or 8 oz. in 25 gals. water. Some leaf injury may occur on cucurbits if treatments are made when tender foliage is wet or in presence of prolonged high humidity. Repeat at 5 to 7 day intervals as needed. Do not apply spray to vegetables within 3 days of harvest.

FRUIT Insects: Peaches, apricots, plums for coddling moth, oriental fruit moth, peach twig borer, plus curculio, San Jose scale use 2 Tbspns. in 1 gal. water, or 8 oz. in 25 gals. Make first spray when one half petals have fallen. Repeat spray 3 or 4 times at 7 to 10 day intervals. Do not apply within 3 days of harvest.

CITRUS Insects: To control red scale, black scale, brown soft scale, orange tortrix worms use same dosage recommended for peaches. Apply for full coverage in normal spray schedules. Consult local agricultural authorities for proper timing of sprays. Do not apply within 5 days of harvest.

GRAPES: For leaf folder, leafhopper, leaf skeletonizer, grape berry moth use 4 Tospns. per 1 gal. water. Make no application within 1 day of harvest. Consult local agricultural authorities for timing of sprays for grapes.

CAUTION: Keep out of reach of children. Harmful if inhaled or swallowed. Do not breathe dust. Do not take internally. Avoid prolonged or repeated breathing of spray. Skin contact may be harmful. Avoid skin contact. Wash thoroughly after using. Wash hands before eating. Avoid contamination of food. Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warm beekeepers well in advance to locate hives at safe distance till I week after application.

Conditions of use are beyond seller's control. Seller's liability is limited to product replacement or purchase price refund. Seller makes no warranty, express or implied, concerning product use beyond claims on label.