

WARNING:

CHLORDANE IS TOXIC. KEEP AWAY FROM CHILDREN. HARMFUL IF SWALLOWED. AVOID INHALING SPRAY OR MIST. MAY BE ABSORBED THROUGH THE SKIN. CONTACT WITH SKIN CAUSES TOXIC SYMPTOMS. WASH THOROUGHLY AFTER USE. DO NOT USE ON HUMANS. AVOID CONTAMINATION OF FEED OR FOODSTUFFS. DO NOT USE ON HOUSEHOLD PETS OR ANIMALS. DO NOT USE ON DAIRY ANIMALS OR THOSE BEING FINISHED FOR SLAUGHTER. THIS PRODUCT WILL KILL FISH AND WILDLIFE.

DO NOT RE-USE EMPTY CONTAINER. CONTAINERS SHOULD BE DESTROYED BY CRUSHING AND BURYING IN A SAFE PLACE.

ANTIDOTE

External — In case of spillage on skin, wash promptly with soap and water.

Internal — Emetic of 1 tablespoonful mustard in tumbler of water. Call a physician.

Fire hazard. Do not spray into open fire or open flame.
Do not smoke while spraying.

U.S.D.A. Reg. No. 539-191

Sold by SEARS, ROEBUCK AND CO.
CHICAGO, ILLINOIS, and SIMPSONS-SEARS
LIMITED, Toronto, Canada — Made in U.S.A.

GENERAL DIRECTIONS TREATMENT FOR TERMITES

STEP ONE: PREPARE THE BUILDING

Repair existing damage. Eliminate wood-to-ground contacts. Remove wood debris, trash and rubbish from around and beneath the building. Scrape away termite shelter tubes from foundation walls and piers. Tunnel into dirt-filled areas beneath porches and terraces. Drill access holes through mortar joints into voids in masonry foundations, fireplaces and flues (8 inches apart, 6 inches above grade level).

STEP TWO: DIG THE TRENCH

Dig a narrow trench along the outside and inside of foundation walls to a depth of about 15-18 inches, but no deeper than the top of the footings. Trench around other danger areas such as piers, conduits and all pipes entering house from ground. If the building has deep foundations, trench to a depth of 30 inches.

STEP THREE: PREPARE THE MIXTURE

Dilute at the rate of 1 quart SEARS TERMITE KILLER AND CONTROL with 25 gallons of water or 1 gallon with 99 gallons of water. Keep the mixture stirred up. If allowed to stand, restir before using.

STEP FOUR: APPLY THE TREATMENT

Inside foundation walls: Apply 4 gallons to 10 lineal feet of trench along foundation walls and around piers and utility entrances. If trench is deeper than 15-18 inches or 30" deep for buildings with deep foundations, apply 8 gallons per 10 lineal feet. Replace soil and apply another 4 gallons per 10 feet. Apply 1 gallon to each 10 square feet of soil surface in crawl space.

Outside foundation walls: Apply 4 gallons per 10 lineal feet in a trench along the outside of foundation walls, including entrance platforms and porches. If trench is deeper than 15-18 inches but not over 30 inches deep, apply 8 gallons per 10 lineal feet. Replace soil and apply another 4 gallons per 10 feet.

Porches and terraces: Drench soil under porches and terraces at the rate of 1 gallon per 10 square feet.

Voids: Flood voids in concrete block, brick or tile foundations, fireplaces and flues with 4 gallons each 10 feet.

Net Contents Makes 100 Gallons

SEARS TERMITE KILLER AND CONTROL

ACCEPTED WITH COMMENTS

2-47051

R
L

structions

EN

NEL

72.80%

21.10%

6.10%

100.00%

9.12% re-
bs F).

Net Weight 30 Pounds

AGSCO DB-GREEN-A

Seed Disinfectant and Insecticide For Drill or Planter Box Application

WHEAT - BARLEY - OATS - FLAX - RYE
CORN - SOYBEANS

INGREDIENTS

ACTIVE INGREDIENTS:

Hexachlorobenzene	10.00%
Maneb (Manganese ethylene bisdithiocarbamate)	40.00%
Lindane (Gamma Isomer of Benzene Hexachloride)	18.75%
INERT INGREDIENTS	31.25%

100.00%

A READY TO USE, DUAL PURPOSE FORMULATION, FOR THE CONTROL OF SEED-BORNE AND SOIL-BORNE DISEASES AND FOR TEMPORARY PROTECTION OF SEEDS & SEEDLINGS AGAINST WIREWORMS.

WARNING

KEEP OUT OF REACH OF CHILDREN

(For Additional Precautionary Statements See Back Panel)

Manufactured by

AGSCO CHEMICALS, INC.

Grand Forks, North Dakota

Jan 3, 1972

554-127

Product Information

AGSCO DB-GREEN-A is a special seed treatment formulation of Lindane, Maneb and Hexachlorobenzene which, when used in accordance with Directions For Use, protects crops against seed-borne diseases such as stinking smut of wheat, oat smut, covered and false loose smut of barley, and bunt of rye. It also gives temporary protection to cereal crops, corn and soybeans against certain soil-borne diseases such as root rot, seedling blight and for temporary protection of seeds and seedlings against wireworms.

AGSCO DB-GREEN-A is especially designed for on-the-farm application as a Drill Box Treatment.

Directions For Use

WHEAT, BARLEY, OATS, RYE—Use rate of 2 oz. per bushel.

FLAX—Use rate of 4 oz. per bushel.

CORN AND SOYBEANS—Use rate of 2½ oz. per bushel.

MIX AGSCO DB-GREEN-A THOROUGHLY into the seed until uniformly applied by:

- (1) Use an AGSCO Automatic Seed Treater that meters in the proper amount of AGSCO DB-GREEN-A and uniformly mixes it as clean untreated seed is augered from the grain box into the planter or drill box as uniformly treated seed.
- (2) Placing and leveling one-half of the seed in the drill or planter box and sprinkle one-half of the total DB-GREEN-A required (for the full planter box) uniformly on or under the surface of the seed. Mix thoroughly with a stick or paddle, then fill the box with seed and sprinkle the remaining one-half of the powder on or under the surface and mix again until all streaking has disappeared and coloring of seed appears uniform. **DO NOT MIX WITH HANDS.**
- (3) Dribble the required amount of DB-GREEN-A into each bushel of seed as it is poured into the drill box and mix with a paddle until uniform.

Avoid very deep seeding and exceptionally early sowing under poor conditions for maximum benefits. Use only recommended rates. Seed should be well cured, dry and clean before treatment.

WARNING

Avoid contact with AGSCO DB-GREEN-A. It may be absorbed through the skin. Avoid contact with skin. Do not breathe dust. It may be fatal if inhaled or swallowed. Wear mask over mouth and nose while handling. Wash thoroughly with soap and water after handling and before eating or smoking. In case of contact with eyes, flush with plenty of water and get medical attention.

Do not treat seed indoors unless there is thorough ventilation. Store treated seed in a ventilated place away from foodstuffs. Keep container tightly closed when not in use. Avoid any contamination of feed and foodstuffs. Do not use any treated seed for feed.

If Maneb becomes wet in storage, its fungicidal effectiveness may be reduced and it may also produce flammable vapors. Keep away from flames or sparks. Store in a cool, dry place.

ENVIRONMENTAL AND WILDLIFE WARNING: Toxic to fish and wildlife. Do not contaminate any body of water by cleaning of equipment and disposal of wastes and containers. Treated seed exposed on soil surface will be hazardous to birds and other wildlife. Dispose of excess treated seeds by burial. Destroy empty containers by burying.

Treated seed should be labeled as follows:

**DISINFECTED SEED
POISONOUS TO ANIMAL
AND MAN
DO NOT USE FOR FOOD,
FEED OR OIL**

Warranty and Limitation of Damages

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the Directions For Use. AGSCO CHEMICALS, INC. makes no other express or implied warranty, including any other express or implied warranty of fitness or of Merchantability, and no agent of AGSCO CHEMICALS, INC. is authorized to do so except in writing with specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or value, etc.