




ACCEPTED
 10/11/71
 U.S. DEPARTMENT OF AGRICULTURE
 FOR THE NATIONAL PESTICIDE CONTROL BOARD
 THE UNITED STATES OF AMERICA

WEED KILLER

**WITH 2, 4-D*
 10-6-4**

- COVERS 5000 SQUARE FEET FOR A WEED FREE LAWN**
 kills broadleaf weeds such as Dandelion, Plantain, Trifolium, Wild Morning Glory, Marlow, and Mustard.
- STIMULATES GRASS GROWTH**
 Stimulates grass growth to cover spots where weeds have died.
- SAFE FOR ESTABLISHED LAWNS**



**CONTENTS:
 20 POUNDS NET
 Z180214**

CAUTION

KEEP OUT OF REACH OF CHILDREN!
 SEE BACK PANEL FOR ADDITIONAL CAUTIONS

ACTIVE INGREDIENTS:		
2,4-Dichlorophenoxyacetic Acid	10.0%	
INERT INGREDIENTS:	99.0%	
(Relative to herbicidal contents)		
	TOTAL	100.0%
*2,4-Dichlorophenoxyacetic Acid equivalent		
MINIMUM GUARANTEED FERTILIZER ANALYSIS		
Total Nitrogen (N)	10.0%	
Available Phosphorus (As P ₂ O ₅)	6.0%	
Soluble Phosphorus	4.0%	
Potential Acidity 650 lbs. of Calcium Carbonate per ton		

U.S.D.A. REG. NO. 579200

Sold by SEARS, ROEBUCK AND CO., CHICAGO, ILL. 60607

BEST DOCUMENT AVAILABLE

SEARS

WITH
70


COVERS 5000 SQUARE FEET
FOR A WEED FREE LAWN

Kills broadleaf weeds such as Dandelion,
Morning Glory, Mallow, Wild Mustard

STIMULATES GRASS GROWTH

Stimulates grass growth to cover bare spots

SAFE FOR ESTABLISHED LAWNS



CONTENTS:

20 POUNDS NET
7180214

C

KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR PRECAUTIONS

ACTIVE INGREDIENT:

acetone, 2, 4-D

INERT INGREDIENTS:

(Relative to net weight)

*2, 4-D-monoisobutoxyacetate

MINIMUM GUARANTEED

Total Nitrogen (N)

Available Phosphoric Acid (P₂O₅)

Soluble Potash

Potential Acidity 650 lbs. of CaCO₃

Sold by SEARS, ROEBUCK & CO.

FEED AND WEED WITH 2, 4-D* 10-6-4

SPREADER SETTINGS

Sears Calibrated	6½	Jackson	6½
Lawn Beauty	5	Central	6½
Scott	5		

Note: Except for Sears Calibrated models, above spreader settings are approximate. Check your spreader on 100 sq. ft. and adjust to appl. only 6 oz. per 100 sq. ft.

DIRECTIONS

Apply when weeds are in active growth. Better results are obtained if applied when ground moisture is adequate. During dry weather water lawn thoroughly 24 hours prior to application. For persistent weeds or if a heavy rain within 2 days after application, a second application may be necessary.

FOR ESTABLISHED LAWNS

With Sears Calibrated spreader, apply evenly at rate of 4 lbs. per 1000 sq. ft. Do not apply to creeping grass such as Bentgrass or to Dichondra. Do not water, mow, or make heavy use of lawn for at least 24 hours after application.

FOR NEW LAWNS

Do not apply to newly seeded lawn until after the second or third mowing.

CAUTION

Avoid inhaling dust. Avoid contact with skin, eyes, or clothing. Do not apply directly to children, pets, or plants. Do not apply to flowers, fruit trees, ornamentals, or other desirable plants. Do not store near fertilizer, insecticides, or herbicides.

DO NOT REUSE CONTAINER. DESTROY BY BURYING IN A SAFE PLACE. DO NOT BURN.

Sold by SEARS, ROEBUCK AND CO., CHICAGO, ILL.

FRONT LABEL

SEARS (logo) SPOT WEED KILLER

- KILLS KNOTWEED, CHICKWEED, CLOVER, POISON IVY, WOODY PLANTS, MANY OTHER LAWN WEEDS.
- CONTAINS 2,4-D & BANVEL[®]D

CAUTION

539-212

KEEP OUT OF REACH OF CHILDREN. CONTENTS UNDER PRESSURE. (See back panel for additional cautions)

71 11505

Net Weight 15 oz.av.

USDA Reg.No. 539-212

SIDE LABEL

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid*	1.27%
Dimethylamine salt of dicamba (3,6-dichloro-o-anisic acid**)	.64%
Dimethylamine salt of related acid	.10%
INERT INGREDIENTS:	<u>97.99%</u>
TOTAL	100.00%

* Equivalent to 1.05% of 2,4-Dichlorophenoxyacetic acid

**Equivalent to .53% of 3,6-Dichloro-o-anisic acid

PACK LABEL

Destroys many common weeds without injuring most permanent grasses. Ideal for spot treating lawns. Single application to vigorously growing weeds will improve appearance of lawn. Apply in spring, summer and fall to control broadleaf weeds in lawn. Apply when there is no wind.

Do not use on Bent grass, Carpetgrass, Centipede, Dichondra, Lippia, St. Augustine, or any young grasses in seedling stage. Clover can be severely damaged or killed. Wait 3 weeks to reseed after spraying. Treat new lawns after second mowing.

LAWN WEEDS: Shake can well and hold 15 to 18 inches from foliage, push button down using short bursts. Wet all plants thoroughly using care to avoid spray mist drift to desirable plants.	Knottweed, Sheep Sorrel, Dandelion, Plantain, Buckhorn, Ground Ivy, Thistles, Yellow Sorrel, Henbit, Pennywort, Wild Onion, Chickweed, and Clovers, Oxalis, Spotted Spurge, Heal-all, Purslane, Yarrow, Veronica, Wild Garlic.
NOXIOUS WEEDS; WOODY PLANTS: Repeat application 7 to 10 days if necessary.	Thistle, Ragweed, Bindweed, and Dock, Poison Ivy, Poison Oak, ...