


SEARS (Logo)

FRONT PANEL

LAWN ~~FEED~~-RENOVATOR  
AND  
GRASS & WEED KILLER

TREATS \* SQUARE FEET OF LAWN



ACTIVE INGREDIENTS

Sodium Cacodylate .....	1.98%
Dimethylarsinic Acid (Cacodylic Acid) .....	0.34%

<u>INERT INGREDIENTS</u> .....	97.68%
--------------------------------	--------

Total arsenic (as elemental) all in water soluble form .....	1.11%
---	-------

Top kills Bermudagrass, Johnsongrass, Quackgrass and other deep rooted perennials.  
Kills annuals such as: Brome, Foxtail, Wild Oats, Burr Clover, Mallow and Morning  
Glory.

CAUTION: KEEP OUT OF REACH OF CHILDREN. SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

71 11017 Contents 1 Quart

USDA Reg. No. 539-R0

71 11018 Contents 1 Gallon

DISTRIBUTED BY SEARS, ROEBUCK AND CO., CHICAGO, ILLINOIS

\* For 71 11017 Quart: 250

For 71 11018 Gallon 1,000



<p>FOR NOXIOUS WEEDS ON VACANT LOTS: Mix 2 tablespoonsful in 1 gal. water or <math>\frac{1}{2}</math> pt. in 6 gal. water.</p>	<p><del>WHISTLES, BACHMID, BLYNDRED, DOCK, MORNING GLORY</del> Wet all plants thoroughly. Repeat application as necessary.</p>
<p>FOR WOODY PLANTS ALONG DITCHES, ROADWAYS, OR SIMILAR NON-CROP LANDS: Mix <math>\frac{1}{4}</math> tablespoonsful in 1 gal. water or <math>\frac{1}{2}</math> pt. in 3 gal. water.</p>	<p><del>POISON IVY, SUMAC, HONEY SUCKLE, POISON OAK AND BRAMBLES</del> Apply when the new leaves are well developed but before the leaves become fully grown. Wet leaves thoroughly. Repeat application as necessary.</p>

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact, flush with plenty of water; for eyes, get medical attention. Avoid spray drift to susceptible plants such as: garden vegetables, flower beds, shrubbery and other ornamentals. Coarse sprays are less likely to drift. Do not apply on or near desirable trees or in locations where chemical may be washed or moved into contact with their roots.

Do not use same sprayer for other purposes unless you thoroughly clean spray equipment with suitable chemical cleaner. Do not store near fertilizer, seeds, insecticides, fungicides.

WARRANTY. Conditions of use are beyond seller's control. Seller's liability is limited to product's replacement or purchase price refund. Seller makes no warranty, express or implied, concerning product's use beyond claims on label.

BEST DOCUMENT AVAILABLE

CAUTION:  
HAZARDOUS IF SWALLOWED

Avoid contact with skin, eyes or clothing and inhaling spray. In case of contact, flush with plenty of water; for eyes, get medical attention.

Do not store near fertilizers, seeds, insecticides or fungicides.

Avoid contamination of irrigation ditches and water used for domestic purposes.

Avoid spray drift to susceptible plants such as: garden vegetables, flower beds, shrubbery and other ornamentals. Minute quantities can cause severe damage. Vapors from this product may cause injury to susceptible plants growing nearby.

Contents under pressure. Do not puncture, incinerate or store above 120°F.

Conditions of use are beyond seller's control. Seller's liability is limited to product replacement or purchase price refund. Seller makes no warranty, express or implied, concerning product use beyond claims on label.

DISTRIBUTED BY SEARS, ROEBUCK AND CO., CHICAGO, ILLINOIS 60607

BEST DOCUMENT AVAILABLE

539-226

XX

7

**CAUTION:**  
**HARMFUL IF SWALLOWED**

Avoid contact with skin, eyes or clothing. In case of contact, flush with plenty of water.

Do not store near fertilizers, seeds,

or in contamination of irrigation water for agricultural purposes.

Avoid spray drift to susceptible plants, shrubs, and other ornamentals. Vapors from this product may be irritating to growing nearby.

Contents under pressure. Do not puncture.

Warranties of use are beyond seller's control. No product replacement or purchase price refund, express or implied, concerning product.

DISTRICTS BY BRAND, ROBBUCK



(2)

FOR NOXIOUS WEEDS ON VACANT LOTS: Mix 2 tablespoonsful in 1 gal. water or $\frac{1}{2}$ pt. in 6 gal. water.	THISTLES, RAGWEED, BINDWEED, DOCK, MORNING GLORY Wet all plants thoroughly. Repeat application as necessary.
FOR WOODY PLANTS ALONG FENCEDROWS, ROADWAYS, OR SIMILAR NON-CROP LANDS: Mix $\frac{1}{4}$ tablespoonsful in 1 gal. water or $\frac{1}{2}$ pt. in 3 gal. water.	POISON IVY, SUMAC, HONEY SUCKLE, POISON OAK AND BRAMBLES Apply when the new leaves are well developed but before the leaves become fully grown. Wet leaves thoroughly. Repeat application as necessary.

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact, flush with plenty of water; for eyes, get medical attention. Avoid spray drift to susceptible plants such as: garden vegetables, flower beds, shrubbery and other ornamentals. Coarse sprays are less likely to drift. Do not apply on or near desirable trees or in locations where chemical may be washed or moved into contact with their roots.

Do not use same sprayer for other purposes unless you thoroughly clean spray equipment with suitable chemical cleaner. Do not store near fertilizer, seeds, insecticides, fungicides.

WARRANTY. Conditions of use are beyond seller's control. Seller's liability is limited to product's replacement or purchase price refund. Seller makes no warranty, express or implied, concerning product's use beyond claims on label.

Labels (10-6) (10-6) (10-6) (10-6) (10-6)

(Contains 100% D and 1,4-D)

KILLS KNOX, INC., CHICAGO, ILLINOIS  
KILLS MANY OTHER LAMN WEEDS  
KILLS POISON IVY, WOODS PLANTS

ACTIVE INGREDIENTS:

Dimethylamine salt of 2,4-dichlorophenoxyacetic acid*	12.00%
Dimethylamine salt of dicamba (3,5-dichloro- <u>o</u> - anisic acid)**	2.10%
Dimethylamine salt of related acid	.30%

INERT INGREDIENTS:

65.60%

TOTAL 100.00%

\* Equivalent to 1.00% of 2,4-dichlorophenoxyacetic acid  
\*\* Equivalent to 0.90% of 3,5-dichloro-o-anisic acid

CAUTION: Keep Out of Reach of Children  
(See back panel for additional cautions)

- (1) 11601 (Net Contents 8 ounces) (Covers up to 2,400 sq. ft.)
- (2) 11602 ( " 1 pint ) ( " " 1,200 " )
- (3) 11603 ( " 1 quart ) ( " " 9,600 " )

**ACCEPTED**

**SEP 28 1967**

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO. 537-22 (SUBJECT  
TO ATTACHED COMMENTS.)

100% D is a registered trademark of  
Melsco Chemical Corporation.

USDA Reg. No. 539-X

DISTRIBUTED BY SENECA, INC., CHICAGO, ILLINOIS

USE

Directions

Use only on lawns which are not being mowed. Apply  
uniformly when weeds are in early stages of growth.  
Apply in spring, summer and fall. Do not use on  
Broomrape, Bluegrass, St. Augustine, or other  
perennial grasses. Do not use on  
perennial grasses after seeding.

<p>1. Mix 1/2 cup of product with 1 gal. of water.</p> <p>2. Spray on 2,400 sq. ft. of lawn.</p> <p>3. Repeat every 2-3 weeks.</p>	<p>1. Mix 1/4 cup of product with 1 gal. of water.</p> <p>2. Spray on 1,200 sq. ft. of lawn.</p> <p>3. Repeat every 2-3 weeks.</p>
--	--



<p>FOR NOXIOUS WEEDS ON VACANT LOTS:</p> <p>Mix 2 tablespoonsful in 1 gal. water or <math>\frac{1}{2}</math> pt. in 6 gal. water.</p>	<p>WHISTLES, RAGWEED, BINDWEED, DOCK, MORNING GLORY</p> <p>Wet all plants thoroughly. Repeat application as necessary.</p>
<p>FOR WOODY PLANTS ALONG FENCEROWS, ROADWAYS, OR SIMILAR NON-CROP LANDS:</p> <p>Mix 4 tablespoonsful in 1 gal. water or <math>\frac{1}{2}</math> pt. in 3 gal. water.</p>	<p>POISON IVY, SUMAC, HONEY SUCKLE, POISON OAK AND BRAMBLES</p> <p>Apply when the new leaves are well developed but before the leaves become fully grown. Wet leaves thoroughly. Repeat application as necessary.</p>

**CUTION:** Avoid contact with skin, eyes and clothing. In case of contact, flush with plenty of water; for eyes, get medical attention. Avoid spray drift to susceptible plants such as: garden vegetables, flower beds, shrubbery and other ornamentals. Coarse sprays are less likely to drift. Do not apply on or near desirable trees or in locations where chemical may be washed or moved into contact with their roots.

Do not use same sprayer for other purposes unless you thoroughly clean spray equipment with suitable chemical cleaner. Do not store near fertilizer, seeds, insecticides, fungicides.

**WARRANTY.** Conditions of use are beyond seller's control. Seller's liability is limited to product's replacement or purchase price refund. Seller makes no warranty, express or implied, concerning product's use beyond claims on label.

<p>FOR NOXIOUS WEEDS ON VACANT LOTS: Mix 2 tablespoonsful in 1 gal. water or ½ pt. in 6 gal. water.</p>	<p><del>THISSEN, RAGWEED, BINDWEED, DOCK, MORNING GLORY</del> Wet all plants thoroughly. Repeat application as necessary.</p>
<p>FOR WOODY PLANTS ALONG FENCES, ROADWAYS, OR SIMILAR NON-CROP LANDS: Mix 4 tablespoonsful in 1 gal. water or ½ pt. in 3 gal. water.</p>	<p><del>POISON IVY, SIDA, HONEY SUCKLE, POISON OAK AND BRAMBLES</del> Apply when the new leaves are well developed but before the leaves become fully grown. Wet leaves thoroughly. Repeat application as necessary.</p>

CAUTION: Avoid contact with skin, eyes and clothing. In case of contact, flush with plenty of water; for eyes, get medical attention. Avoid spray drift to susceptible plants such as: garden vegetables, flower beds, shrubbery and other ornamentals. Coarse sprays are less likely to drift. Do not apply on or near desirable trees or in locations where chemical may be washed or moved into contact with their roots.

Do not use same sprayer for other purposes unless you thoroughly clean spray equipment with suitable chemical cleaner. Do not store near fertilizer, seeds, insecticides, fungicides.

WARRANTY. Conditions of use are beyond seller's control. Seller's liability is limited to product's replacement or purchase price refund. Seller makes no warranty, express or implied, concerning product's use beyond claims on label.