

ACCEPTED

Beans (Logo) Indiana State University

533 750

For use on Soybeans, Peas, Lentils, Dry Beans, Flowers, Grasses, Fruit and Nut Trees to control nematodes.

May be used on certain lawn grasses to suppress nematode damage.

2-D-thiophene-5-oximidazole	5.0%
Other inactive ingredients	95.0%
Total	100.0%

CAUTION: Keep out of reach of children. See additional cautions on label.

Net weight - 1 lb. USDA Reg. No. 539-X

Manufactured by Deery, Hockaday and Co., Chicago, Ill. 60607

General Directions for Use

The use of this material to control nematodes is approximately 2 ounces per 100 square feet of area. Apply to the soil surface and work into the soil to a depth of 6 to 8 inches as soon as possible. Soil should be in good condition and nematode population should be high. Application can be made before, during, or after planting, depending on the crop. Use only on crops listed in the label. Do not use on alfalfa, clover, sugar beets, and tobacco and do not use on soil surfaces. Do not use on soil surfaces, food or feedstuffs. Do not use on soil surfaces which will be subjected to poultry. Apply in accordance with directions for use on label. Consult State and local authorities for use on other soils.

Directions for Use: For most nematodes, apply 2 ounces per 100 square feet of area. For nematodes which are not listed on the label, apply 2 ounces per 100 square feet of area. For nematodes which are listed on the label, apply 2 ounces per 100 square feet of area. For nematodes which are listed on the label, apply 2 ounces per 100 square feet of area. For nematodes which are listed on the label, apply 2 ounces per 100 square feet of area.

Precautions: Do not use on alfalfa, clover, sugar beets, and tobacco and do not use on soil surfaces. Do not use on soil surfaces, food or feedstuffs. Do not use on soil surfaces which will be subjected to poultry. Apply in accordance with directions for use on label. Consult State and local authorities for use on other soils.

Application of this material to control nematodes is approximately 2 ounces per 100 square feet of area. Apply to the soil surface and work into the soil to a depth of 6 to 8 inches as soon as possible. Soil should be in good condition and nematode population should be high. Application can be made before, during, or after planting, depending on the crop. Use only on crops listed in the label. Do not use on alfalfa, clover, sugar beets, and tobacco and do not use on soil surfaces. Do not use on soil surfaces, food or feedstuffs. Do not use on soil surfaces which will be subjected to poultry. Apply in accordance with directions for use on label. Consult State and local authorities for use on other soils.

Net weight - 1 lb.

For use on Soybeans, Peas, Lentils, Dry Beans, Flowers, Grasses, Fruit and Nut Trees to control nematodes. May be used on certain lawn grasses to suppress nematode damage.

CAUTION: Keep out of reach of children. See additional cautions on label. Net weight - 1 lb. USDA Reg. No. 539-X

Manufactured by Deery, Hockaday and Co., Chicago, Ill. 60607

General Directions for Use

The use of this material to control nematodes is approximately 2 ounces per 100 square feet of area. Apply to the soil surface and work into the soil to a depth of 6 to 8 inches as soon as possible. Soil should be in good condition and nematode population should be high. Application can be made before, during, or after planting, depending on the crop. Use only on crops listed in the label. Do not use on alfalfa, clover, sugar beets, and tobacco and do not use on soil surfaces. Do not use on soil surfaces, food or feedstuffs. Do not use on soil surfaces which will be subjected to poultry. Apply in accordance with directions for use on label. Consult State and local authorities for use on other soils.

Directions for Use: For most nematodes, apply 2 ounces per 100 square feet of area. For nematodes which are not listed on the label, apply 2 ounces per 100 square feet of area. For nematodes which are listed on the label, apply 2 ounces per 100 square feet of area. For nematodes which are listed on the label, apply 2 ounces per 100 square feet of area.

Precautions: Do not use on alfalfa, clover, sugar beets, and tobacco and do not use on soil surfaces. Do not use on soil surfaces, food or feedstuffs. Do not use on soil surfaces which will be subjected to poultry. Apply in accordance with directions for use on label. Consult State and local authorities for use on other soils.