


7180227

NET WT. 20 LBS.
FEEDS 5000 Sq. Ft.



Superfine Weed and Feed

25-3-3 FOR SOUTHERN LAWNS

CAUTION
KEEP OUT OF REACH OF CHILDREN!
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS



USE ON St. Augustine, Bahia, Fescue, Rye,
Bermuda, Centipede and Zoysia Lawns
FOR CONTROL OF ... growing broadleaf weeds
listed on back of bag

ACTIVE INGREDIENTS:

* Dimethylamine Salt of 2, 4-dichlorophenoxyacetic acid	0.16%
** Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	0.52%
*** Dimethylamine Salt of Dicamba (3, 6-dichloro-o-anisic acid)	0.06%
INERT INGREDIENTS:	99.26%
TOTAL	100.00%

Used at 20 lbs. on 5,000 sq. ft. equivalent to:

* 2, 4 D Acid	0.124% (0.23 lbs. acre or 2.63 lbs. per ton)
** MCPA Acid	0.436% (0.86 lbs. acre or 8.72 lbs. per ton)
*** Dicamba Acid	0.054% (0.11 lbs. acre or 1.07 lbs. per ton)

CONTAINS TRIVECT HERBICIDE

* Reg. T. M. IPI Garden Corporation - U.S. Patent No. 3,284,186

EPA Reg. No. 589-072

EPA Est. No. 9404 FL-1

Sold by SEARS, ROEBUCK AND CO. • CHICAGO, ILL. 60684

529-272

BEST DOCUMENT AVAILABLE

NET WT. 20 LBS.
FEEDS 5000 SQ. FT.

SUPERWEED WEED & FEED

25-3-3 FOR SOUTHERN LAWNS

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	25.00%
4.00 Ammoniacal Nitrogen	
21.00 Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	3.00%
Soluble Potash (K ₂ O)	3.00%
Chlorine not more than	3.00%
Derived from Urea, Ammoniated Superphosphate, Sulphate of Ammonia, Muriate of Potash	
Potential Acidity equivalent to 800 pounds of Calcium Carbonate per Ton.	

DIRECTIONS FOR USE

CONTROLS SUCH WEEDS AS: Florida Pusley, Black Medic, Chickweed, Clover, Cockspur, Dandelion, Creeping Charlie, Dock, Henbit, Knotweed, Lambsquarter, Mustard, Oxalis, Peppergrass, Pigweed, Plantains, Poison Ivy, Parslane, Ragweed, Sida, Spurge, Wild Onion, Wild Lettuce, and most other broadleaf weeds.

TIME TO APPLY: Apply any time broadleaf weeds are present, except to avoid treating St. Augustine Grass in summer months (June through September) when it is growing rapidly, because temporary foliage discoloration may occur. Best results are obtained when weeds are young and growing vigorously. If weeds are growing slowly, control will be slow and more than one application may be needed. Disturbance from dew or watering on weed foliage at time of application aids control. Do not water for 24 to 48 hours after application.

HOW MUCH TO APPLY: Use 20 lbs. on 5,000 sq. ft. for Bahia, St. Augustine Grass and most other grasses. Over-application may cause temporary discoloration.

SUGGESTED SPREADER SETTINGS

May vary with age, model and condition of spreader and walking speed. Check your spreader by trial on a known area.

Make of Spreader	Spreader Settings
Sears Broadcast (5 ft. swath)	6½
Sears Calibrated (20")	8
Control (20")	8
Cyclone (5 ft. swath)	6½
Scott (21")	7

CAUTION

Keep out of reach of children. Keep away from feed and food products. Do not take internally. Avoid inhalation or contact with eyes, skin or clothing. Wash after use. Do not use on driveways or light ground cover or similar ground covers. Do not use on lawns. Do not use on roofs. Do not use on areas enclosed by trees and shrubs. Avoid contact with trees and shrubs.


Apply to lawn, not to garden plants, flowers and shrubs. Do not use on newly seeded lawns. Do not use on lawns that have been recently treated with other fertilizers. Keep away from plants, seeds, fertilizers, insecticides and fungicides.


WARRANTY

Guaranteed to control weeds and feed lawns. If it does not, your money will be refunded. No other conditions apply.

Sold by SEARS, ROEBUCK AND CO. • CHICAGO, ILL. 60684

539-272





2130227

NET WT. 20 LBS.

FEEDS 5000 Sq. Ft.

Superfine Weed and Feed

25-3-3 FOR SOUTHERN LAWN

CAUTION
KEEP OUT OF REACH OF CHILDREN!
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS



USE ON St. Augustine, Bahia, Fescue, Rye,
Bermuda, Centipede and Zoysia Lawns

FOR CONTROL OF . . . growing broadleaf weeds
listed on back of bag

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2, 4-dichlorophenoxyacetic acid	0.16%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy)propionic acid	0.52%
***Dimethylamine Salt of Dicamba (3, 6-dichloro o-anisic acid)	0.05%
INERT INGREDIENTS:	99.26%

TOTAL

100.00%

Used at 20 lbs. on 5,000 sq. ft. equivalent to:

*2, 4-D Acid	-0.134%-0.23 lbs. acre or 2.68 lbs. per ton
**MCPP Acid	-0.436%-0.74 lbs. acre or 8.72 lbs. per ton
***Dicamba Acid	-0.054%-0.08 lbs. acre or 1.07 lbs. per ton

CONTAINS TRIMEC® HERBICIDE+

©Reg. T. M. PBI-Gordon Corporation + U.S. Patent No. 3,284,186

EPA Reg. No. 539-272

EPA Est. No. 9404-FL-1

Sold by SEARS, ROEBUCK AND CO. • CHICAGO, ILL. 60684


RECORDED
539-272

BEST DOCUMENT AVAILABLE
412

NET WT. 20 LBS.
FEEDS 5000 SQ. FT.

SUPERFINE WEED & FEED

25-3-3 FOR SOUTHERN LAWNS



GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) 25.00%
4.00 Ammoniacal Nitrogen
21.00 Urea Nitrogen
Available Phosphoric Acid (P₂O₅) 3.00%
Soluble Potash (K₂O) 3.00%
Chlorine not more than 3.00%
Derived from Urea, Ammoniated Superphosphate, Sulphate of Ammonia, Muriate of Potash
Potential Acidity equivalent to 800 pounds of Calcium Carbonate per Ton.

DIRECTIONS FOR USE

CONTROLS SUCH WEEDS AS: Florida Pusley, Black Medic, Chickweed, Clover, Cocklebur, Dandelion, Creeping Charlie, Dock, Henbit, Knotweed, Lambsquarter, Mustard, Oxalis, Peppergrass, Pigweed, Plantains, Poison Ivy, Purslane, Ragweed, Sida, Spurge, Wild Onion, Wild Lettuce, and most other broadleaf weeds.

TIME TO APPLY: Apply any time broadleaf weeds are present, except to avoid treating St. Augustine Grass in summer months (June through September) when it is growing rapidly, because temporary foliage discoloration may occur. Best results are obtained when weeds are young and growing vigorously. If weeds are growing slowly, control will be slow and more than one application may be needed. Moisture from dew or watering on weed foliage at time of application aids control. Do not water for 24 to 48 hours after application.

HOW MUCH TO APPLY: Use 20 lbs. on 5,000 sq. ft. for Bahia, St. Augustine Grass and most other grasses. Over-application may cause temporary discoloration.

SUGGESTED SPREADER SETTINGS

May vary with age, model and condition of spreader and walking speed. Check your spreader by trial on a known area.

Make of Spreader	Spreader Settings
Sears Broadcast (5 ft. swath)	6½
Sears Calibrated (20")	8
Central (20")	8
Cyclone (5 ft. swath)	6½
Scott (21")	7

CAUTION

Keep out of reach of children. Keep away from feed and food products. Do not take internally. Avoid inhalation or contact with eyes, skin or clothing. Wash after use. Do not use on dichondra or lippia ground cover or similar ground covers. Do not re-use empty bag. Destroy when empty. Do not apply in areas underlaid by roots of desirable trees and shrubs. Avoid drift of dust to desirable plants.

Avoid application on garden plants, flowers and shrubs. Do not use on newly seeded lawns. Do not contaminate irrigation ditches or water used for irrigation purposes. Store in a dry place, away from plants, seeds, fertilizers, insecticides or fungicides.

WARRANTY

Since conditions of use are beyond seller's control, Sears and the manufacturer's liability under this warranty is limited to replacement or refund. If this product is unsatisfactory, Sears or the manufacturer will provide a refund or replacement.

Sold by SEARS, ROEBUCK AND CO. CHICAGO, ILL. 60684

ST DOCUMENT AVAILABLE.

49