

**SEARS PRE-EMERGE CRABGRASS CONTROL
and
FERTILIZER**

10-0-4

DIRECTIONS FOR USE

Early spring application of Sears Pre-Emerge Crabgrass Control and Fertilizer, when germinate, provides full season control of crabgrass and adds essential to good grass growth.

Control: Crabgrass, Foxtail, Johnson Grass (from seed), and Younggrass.

When spreader is in good operating condition, apply this at a rate of 8 pounds per 1000 sq. ft.

SUGGESTED SPREADER SETTINGS

Calibrated 7	Central 7 1/2
Sears Broadcast 5 1/2	Jackson 7
Cyclone 5 1/2	Scott (Model 35) 5 1/2

This product is also ideal for controlling above listed weeds around shrubbery and ornamentals.

Treated areas should be watered lightly if no rain falls within one week of application.

Reseeding should be delayed until 60 days after application.

NOTE FOR NEW LAWNS- Do Not Apply to Newly Seeded Lawn Until After Lawn Is Completely Green. Crabgrass Already Growing Will Not Be Controlled.

CAUTION

Avoid contact with eyes. Wash after handling and using. Do not use on Dichondra lawns, Bugle Weed, Button Bush, Carnation, Geum, Germander, Mesembryanthemum, Pansy, Phlox, Sweet William and Telandthera. Do not contaminate feed or foodstuff or store near feed, fertilizer or other pesticides. Do not reuse container. Do not apply to lawns, roads, highways, ditches. Do not graze treated areas or feed plant parts. Destroy when empty.

Non-Warranty: Conditions of sale are payment and seller's control. Seller's liability is limited to the product's replacement or purchase price refund. Seller makes no warranty, express or implied, concerning product's use beyond claims on label.

Sold by SEARS, ROEBUCK AND CO. CHICAGO, ILL. 60684

SEARS PRE-EMERGE CRABGRASS CONTROL & FERTILIZER

ACCEPTED
EPA Reg. No. 539286

SEARS, ROEBUCK AND CO.
CHICAGO, ILL. 60684

DIRECTIONS


Puritan's CURTAINS is highly soluble in water and adaptable to preparation sprays. Always use sufficient water to insure distribution of CURTAINS over area being treated, i.e., about 125 gallons per acre. Spray while weeds are growing actively and after leaves are fully developed, but prior to bud development. Since the response of weeds in different areas to 2,4-D Compounds is so greatly influenced by environmental conditions, users should consult experiment station or extension service weed specialists.

On pastures, lawns, golf courses, parks, playgrounds and other turf areas, for control of broadleaf weeds such as plantain, dandelion and chickweed, and shallow-rooted weeds such as yellow mustard and ragweed. Use 8 to 16 pints per acre. This is equivalent to 3-6 ounces of CURTAINS in 3 gallons of water, for every 1000 sq. ft.

For weeds which are difficult to control such as yarrow, goldenrod and wild onions. Use 8 to 24 pints per acre. This is equivalent to 3-9 ounces of CURTAINS in 3 gallons of water, for every 1000 sq. ft.

In vacant lots and airfields, on roadsides, along fence rows and railroad rights-of-way, around farm buildings and other waste areas, for control of deep-rooted perennial weeds such as Canada thistle, bindweed, poison ivy and sow thistle. Use 16 to 32 pints per acre. This is equivalent to 6-12 ounces of CURTAINS in 3 gallons of water, for every 1000 sq. ft.

8-4-72
541-31



CURTAINS

(Formerly Machete)

Kills Dandelion, Goldenrod, Ragweed, Broad and Narrow Leaf Plantain, Thistle, Dock, Wild Morning-Glory, Shepherd's-Purse, Wild Carrot, Mallow, and many other broad-leaved weeds without injury to most grasses.

ACTIVE INGREDIENTS

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid*

7.5%

INERT INGREDIENTS

*Equivalent to 6.2% (0.534 lbs. per Gallon) of 2,4-Dichlorophenoxyacetic acid

92.5%

CAUTION:

KEEP OUT OF REACH OF CHILDREN.

See right panel for additional caution statement.

E.P.A. Reg. No. 541-31

IMPORTANT

Do not apply CURTAINS directly to, or otherwise permit it to contact, vegetables, flowers, fruit trees, ornamentals, cotton, dichondra or other desirable plants which are sensitive to 2,4-D and similar materials. Do not permit spray mist containing it to drift onto them, since even minute quantities of the spray might cause severe injury. Coarse sprays are less likely to drift.

Do not use on lawns of creeping grasses, except for spot spraying, nor on seeded lawns until the grass has become well established. Most legumes are usually damaged or killed. Excessive amounts of 2,4-D in the soil may temporarily inhibit seed germination or plant growth. Do not use weed control equipment for applying insecticides or fungicides or for other purposes, unless cleaned thoroughly with suitable chemical cleaner. Do not store near fertilizers, seeds, insecticides or fungicides. Do not graze treated areas by dairy animals within 7 days after treatment. Do not contaminate water used for any domestic purposes.

CAUTION

May cause irritation. Avoid contact with eyes, skin or clothing. In case of contact, wash off promptly with plenty of water.

Do not re-use empty container. Wash thoroughly with water and detergent. Discard in a safe place.

QUALITY CONTROL NUMBER

NET CONTENTS