

A.A.I.

Animal Area Insecticide

Indoor Space and Surface Spray

Contains both DDVP and Synergized Pyrethrins

ACCEPTED

FEB 19 1968

Knocks down and kills certain flying and crawling insects including both resistant and non-resistant varieties.

Active Ingredients: 100%

Petroleum Distillate

2,2-dichlorovinyl dimethyl phosphate and related compounds (DDVP)

N-octyl bicycloheptane dicarboximide

Technical piperonyl butoxide - Consists of (butylcarbityl) (6-propyl-piperonyl) ether and related compounds.

Pyrethrins

I. Animal buildings (Horse Barns, Dairy Barns, Shelter Sheds), Milk Sheds and Outdoor Areas.

USE AS A SPACE SPRAY, FOG, OR MIST

TO CONTROL FLIES, GNATS, MOSQUITOES:

Use 1 quart to treat 8,000 cu. ft. Before applying reduce air movement as much as possible by closing doors, windows, and other openings. Do not contaminate water, feed or foodstuffs, milk or milking utensils. Do not use this solution for fogging purposes in areas where animals have received direct application within 8 hours. Keep area closed for 15 - 20 minutes after which air out.

TO CONTROL HOUSEFLIES, HORN FLIES, STABLE FLIES, FACE FLIES AND MOSQUITOES ON BEEF AND DAIRY CATTLE: Apply 1 to 2 ounces per animal as a mist spray daily with hand or automatic sprayer. Spray to thoroughly cover all parts of the animal, including the legs, but do not wet the skin. Do not apply in excess of 2 fluid ounces per animal per day.

USE AS A SURFACE SPRAY

TO CONTROL FLIES, GNATS, MOSQUITOES, ROACHES, FLICKS, ANTS, SOMEBUGS, BROWN DOG TICKS:

Apply as a coarse wet spray at the rate of 1 quart per 1,000 sq. ft. For milk sheds, stables, livestock barns, loafing sheds, pig pens and outdoor areas treat around doorways, feed storage rooms, alleyways and window sills. Do not contaminate milk or milking utensils. For roaches treat cracks and crevices.

In poultry houses, apply to manure, window sills, outside of panned enclosures and on floor of feed storage rooms.

Treat outdoor living areas, picnic grounds, backyard areas, patios, loading docks, outdoor latrines, parking areas, refuse areas around service stations, open-air drive-ins, ice cream stands, garbage disposal and collection areas.

In treating dog kennels, apply to outside dog runways, window sills and ledges. For control of fleas and brown dog ticks increase dosage to 2 quarts per 1,000 sq. ft.

For control of fly maggots in maggot breeding areas such as garbage dumps and manure piles, apply at the rate of 1 to 2 quarts per 100 sq. ft.

Before applying in enclosed areas, reduce air movement as much as possible by closing doors, windows, and other openings. Do not treat animals or humans. Remove fish, birds, and pets in areas to be treated. Do not contaminate water, feed or foodstuffs, milk or milking utensils. Repeat as necessary.

CAUTION: (18 pt.) Keep out of reach of children (12 pt.)

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid breathing spray mist. Wash thoroughly with soap and water after handling and before eating or smoking. Do not contaminate food, water or foodstuffs. If swallowed, induce vomiting immediately and get medical attention. Atropine is antidotal.

U.S.D.A. Registration No. 541-174

Net Vol. Gals.

Manufactured by FURITAN CHEMICAL COMPANY, Atlanta, Georgia
Branches: St. Louis - Houston - Detroit - Newark

18

(All label on one panel)

