


PURITAN #3940
CHEMICAL LAWN TRIMEER



Controls weeds, grasses and other forms of undesirable vegetation without damage to valuable trees, shrubs, or neighboring plants.

Use along fences, sidewalks, driveways, buildings and shrubbery borders; around the base of trees, fire plugs, guy wires and utility poles; and in drainage ditches

Non-flammable contact spray. For general post-emergent weed control. Non-selective in action. Phytotoxic properties are quickly inactivated on contact with soil. Controls plant growth only where applied; does not "spread" or "run" to other plant life. Not a soil sterilant. Non-volatile. Contains wetting agent.

DIRECTIONS:

Apply any time when grass is growing actively. Use pressure sprayer - not a sprinkler can. Wet plants thoroughly but avoid overuse. Excessive application may cause temporary non-ductivity of soil, but this condition usually disappears in the "over-winter" period. Do not apply to foliage of valuable plants. Spray on a warm, sunny day. Do not spray during a rain or when rain is threatening, or when weather is cool. PURITAN #3940 must be thoroughly dissolved. Fill the spray equipment reservoir about half full with water and add the required amount of herbicide with agitation. Finish filling the reservoir with water and apply. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period.

FOR TRIMEER EDGE EDGES: Mix about 1/2 gallon with 4 or 5 gallons of water in sprayer and treat up to 100 square feet.

FOR TEMPORARY WEED CONTROL CLOSE TO DESIRABLE ORNAMENTALS IN NON-CROP AREAS: Mix 1 to 2 gallons in 20 gallons of water. Spray unwanted vegetation thoroughly to just short of run-off. Any spray equipment that gives good coverage may be used. If regrowth occurs reapply as required.

FOR LAWN RENOVATION: Mow lawn to be renovated about 1 inch high before treatment. Mix about 1/2 gallon in 4 gallons of water and apply to 1,000 square feet. Cover foliage thoroughly, preferably on a warm sunny day. Foliage will usually turn brown in about five days. Bermuda-grass and other deep rooted perennials, or uncovered foliage, may require a second treatment. Reapply five days later if green areas remain. Rake matted areas when brown and seed 5 days after last application.

Degree of control and duration of effect will vary with weed species, rainfall, temperature and other conditions.

KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL AFTER FIRST RAIN OR SPRINKLING FOLLOWING TREATMENT. DO NOT TRACK FROM TREATED TO UNTREATED AREAS.

CAUTION (18 pt.). Keep out of reach of children (12 pt.)

CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after using. Avoid storage near feed or food products. Keep children and domestic animals off treated areas until this material has been washed into the soil.

ANTIDOTE: If taken internally, induce vomiting and call a physician at once.

Do not allow solution or spray mist to contact desirable vegetation as product will kill plants with which it comes in contact. Wash equipment thoroughly after use. Frequency of retreatment depends on soil fertility and moisture.

Do not store with foodstuffs. Do not contaminate waters used for domestic consumption, or by animals, wildlife and aquatic life, or for irrigation purposes. Do not graze treated areas to livestock.

Do not reuse empty container. Destroy it by perforating or crushing. Bury or discard in a safe place.

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Active Ingredients:

Sodium Cacodylate	4.5 %
Dimethylarsinic acid (Cacodylic acid)	0.78%

Inert Ingredients: 94.727

Total Arsenic (as elemental) all in water soluble form 2.53%

Product contains 0.494 lbs. cacodylic acid equivalent per gallon

U.S.D.A. Registration No. 541-

Net Vol. Gals.

Manufactured by PURITAN CHEMICAL COMPANY, Atlanta, Georgia
Branches: Detroit -- Houston -- Newark -- St. Louis

65