CROAKS

Residual Insecticide Concentrate

Water Emulsifiable

Contains Baygon, Patented Insecticide

For fast knockdown, quick kill and long re idual kill of resistant insects. Quickly knocks down and kills crawling insects such as cockroaches, waterbugs, silver fish, ants, spiders, crickets, brown dog ticks, centipedes, scorpions, and exposed stages of flour beetles, flour moths, weevils, cadelles.

Active Ingredients:

Aromatic petroleum distillate 88.0% *o-Isopropoxyphenyl methylcarbamate 4.0% Inert Ingredients: 8.0%

*U.S. Patent No. 3,111,539

DIRECTIONS

CROAKS is designed for use by Pest Control or Trained Operators only.

CROAKS is a concentrate and should be diluted before application. Mix I part of concentrate with 3 parts of water, or of solvent if preferred. Apply only as a coarse, wet spray or with a small paint brush as a "spot" treatment. Do not use as a space spray indoors. Heavy application is desirable for the most effective kill and greatest residual effect.

FOR ROACHES, WATERBUGS & SILVERFISH: Apply to infested cracks and other areas where insects have been seen resting, running or hiding. Special attention should be given to treatment of baseboards, areas behind shelving, bookcases and storage areas.

FOR ANTS: Apply to ant trails and areas around door sills, window frames, where pests may enter. Treat openings around water pipes, heat ducts and electrical outlets.

BROWN DOG TICKS: Apply to infested areas around baseboards, window and door frames, wall cracks, sleeping quarters of household pets, and localized areas of floors and floor coverings. Fresh bedding should be placed in animal quarters following treatment. Do not treat pets or other animals.

FOR CADELLES, CIGARETTE BEETLES, DERMESTIDS, DRUGSTORE BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RED FLOUS BEETLES, SAW-TOOTHES GRAIN BEETLES & WEEVILS (exposed stages): Thoroughly treat floors and the walls of the infested area with attention to cracks, crevices, corners and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested products.

FOR CLOVER MITES, CRICKETS, EARWIGS, MILLIPEDES, CENTIPEDES, SOWBUGS, SCORPIONS & SPIDERS: Thoroughly spray interior and exterior perimeter of infested premises. For clover mites give special attention to window sills.

FOR FLIES, MOSQUITOES, GNATS, FLYING MOTHS, WASPS, SMALL FLYING INSECTS OUTDOORS: Apply to localized areas these insects may infest; treat around doors and window frames. For preventing entry into dwellings, apply as a spot treatment thoroughly and repeatedly to screens, window frames, door sills and selected areas on porches where insects alight or congregate. For wasps, spray the nest after dark when all insects have returned to nest.

FOR USE IN MEAT PACKING a FOOD PROCESSING PLANTS: Use only in the inedible product areas of meat packing plants and food processing plants, including those that are federally inspected. When using the spray in the inedible products areas where no exposed food or processing equipment is present, take any needed precautions to avoid contamination of packaged or stored food commodities. Rinse all equipment with water before resuming operations. Treat food processing plants early in the season before canning or other processing begins. Repeat applications, if necessary, when plant is not in operation. Spraying unloading docks and outside areas where insects congregate will aid in preventing re-infestation. Good sanitation and proper disposal of wastes are essential in any effective insect control program. Do not use in edible products areas of food processing plants.

(18 pt.): Keep out of reach of children (12 pt.).

Harmful if swallowed, inhaled or absorbed through the skin. Do not spray on humans or animals or near fish bowls. Avoid contamination of feed and foodstuffs, utensils, dishes or drinking water. Avoid inhalation of mist or contact with skin. In case of contact with skin or eyes, wash with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking. Do not use or store near heat or open flame. Do not smoke white spraying. Do not allow children or pets to contact treated surfaces until dry. Remove caged birds and pets prior to treating. Should illness occur, get prompt medical Lid. Atropine sulfate is antidotal. Use with adequate ventilation. Avoid contact with clothing.

Do not use in edible products areas of food processing plants.

U.S.D.A. Registration No. 541-

Net Vol. Gals.

Quality Control No.

Manufactured by PURITAN CHEMICAL COMPANY Aulanta - Detroit - Houston - No ark St. Louis