

American Fluoride Corporation

17 1948

PCE PROLIN* RAT BAIT, Ready-To-Use

CAUTION: KEEP OUT OF REACH OF CHILDREN

ACTIVE INGREDIENTS:

Sulfaquinoxaline (N'(2-quinoxaliny) sulfanilamide)	0.025%
Warfarin 3(a-acetonylbenzyl)-4-hydroxycoumarin	0.025%

INERT INGREDIENTS:

99.950%

TOTAL 100.000%

*PROLIN is a trademark of Wisconsin Alumni Research Foundation, under U.S. Pat. No. 2427578

NO BAIT SHYNESS - NO PRE-BAITING
INEXPENSIVE AND EASY TO USE
NO TOLERANCE BUILD-UP - SLOW BUT SURE

DIRECTIONS: Place PCE PROLIN RAT BAIT in shallow containers in dry locations where rats and mice ordinarily feed, water, travel or live. Baits should be placed in corners along walls, etc., and protected so that children, pets, or domestic animals cannot reach them.

FOR RATS - use 1 or 2 bait placements (¼-1 pound each) at points of greatest activity for a minimum of 10 days.

FOR MICE - locate tablespoon quantities of bait at 6-10 foot intervals where mice are active for a minimum of 15 days. Inspect feeding stations daily and add bait as needed; never allow a station to remain empty over 48 hours. If bait becomes sour, moldy or otherwise decomposed, it should be replaced promptly.

CAUTION: If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K combined with blood transfusions may be indicated as in the case of hemorrhage caused by overdoses of dihydroxycoumarin.

Manufactured by: PEST CONTROL EQUIPMENT CO. DIV.
American Fluoride Corporation
Mt. Vernon, New York 10550

NET WEIGHT:

LOT NO. _____

USDA REG. NO. 542-31

SEP 1948