

1 of 2

NET WT. 50 LBS.

# MALATHION 5 DUST

AN AGRICULTURAL INSECTICIDE AND MITICIDE

ACTIVE INGREDIENT:

Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate) .....

5%

INERT INGREDIENTS .....

95%

100%

U.S.D.A. Reg. No. 550-45

See rear panel for directions for use and recommendations.

## CAUTION! KEEP OUT OF REACH OF CHILDREN

Avoid breathing dust. Avoid repeated contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs.

**Observe the following use precautions:** Observe use limitations given on the label for specific crops. DO NOT apply during bloom when honey bees or other pollinators are active. Avoid dust drift to adjoining food and forage crops about to be harvested, or crops in bloom. Apply in early morning or evening when there is little or no wind. The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that resi-

dues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified. Before marketing food crops, remove residues by washing, stripping, wiping or other effective means. Consult County Agricultural Extension Agent for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

**Note to Physicians:** Malathion, upon repeated prolonged and careless use, may cause cholinesterase inhibition. Atropine is antidotal.

THE USE OF THIS MATERIAL BEING SUBJECT TO CONDITIONS BEYOND SELLER'S CONTROL, SELLER MAKES NO WARRANTY OF ANY KIND, EXPRESS OR IMPLIED, AS TO THIS MATERIAL OR ITS USE

(10/66)

00-504-5A

## Van Waters & Rogers, Inc.

LOS ANGELES • SAN FRANCISCO • SEATTLE

ALBUQUERQUE • ANCHORAGE • BEAUMONT • BOISE • DALLAS • DENVER • EL PASO • EUGENE • FRESNO • HONOLULU  
HOUSTON • PHOENIX • PORTLAND • SACRAMENTO • SALT LAKE CITY • SAN DIEGO • SPOKANE • TUCSON

## DIRECTIONS FOR USE AND ADDITIONAL PRECAUTIONS

Apply when air is calm, using ground equipment, aircraft, or hand duster. Make thorough coverage of plants with a uniform deposit of dust. Apply at the first sign of infestation and repeat at regular intervals when needed to maintain control, but observe use limits given for specific crops. This product is toxic to fish. Keep out of lakes, streams,

and ponds. Apply this product only as specified on this label. Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. (Keep out of smoke.)

## RECOMMENDATIONS

**Alfalfa:** For the control of Aphids, including Yellow Clover Aphid, Potato Leafhopper, Spider Mites, Spittlebugs, Vetch Bruchid and Alfalfa Weevil Larvae apply at the rate of 30 lbs. per acre. Repeat application as necessary. Note: Do not apply to Alfalfa in bloom. Do not apply within 7 days of harvest or pasturing. **Beans:** Mexican Bean Beetle, Leafhoppers, Black Bean Aphid apply 30 to 35 lbs. per acre. Make two or more applications, as needed. Make no applications within 24 hours of harvest. **Brussels Sprouts:** For the control of Aphids on Brussels Sprouts, apply 30 lbs. per acre. Make full coverage applications and repeat as necessary. Make no application within 7 days of harvest. **Cabbage, Broccoli, Kale, Mustard, Turnips:** For the control of Aphids, Imported Cabbageworm and Cabbage Looper apply 30 lbs. per acre. Make full coverage applications, repeated as necessary. Make no application to Cabbage, Mustard and Kale within 7 days of harvest, or to Broccoli and Turnips within 3 days of harvest. **Cucumber, Squash and Melons:** Aphids and Spider Mites apply 30 to 35 lbs. per acre. Caution: Do not apply unless plants are dry. Make no application within one day of picking. **Lettuce:** For the control of Aphids,

Mites and Cabbage Looper apply 30 to 40 lbs. per acre. Do not apply on head lettuce within 7 days of harvest and on leaf lettuce within 14 days of harvest. **Onion:** For control of Onion Thrips apply at rate of 30 to 40 lbs. per acre. Make full coverage applications and repeat as necessary. Make no application on green onions within 3 days of harvest. **Peas:** Pea Aphid, apply 25 lbs. per acre. Make no application within 3 days of harvest. **Potatoes:** Aphids, apply 30 to 40 lbs. per acre. **Spinach:** For the control of Aphids apply 25 to 30 lbs. per acre. Repeat as necessary. Make no application within 7 days of harvest. **Strawberries:** For the control of Aphids and Spider Mites apply 40 lbs. per acre, repeating as necessary. Make no application within 3 days of harvest. **Tomatoes:** Aphids and Spider Mites apply 35 to 40 lbs. per acre. Make no application within 1 day of harvest. **Beef and Dairy Cattle:** Hornfly, use 3 tablespoonsful per animal, applying to back and neck. Repeat application at 10-14 day intervals. Do not apply to dairy cattle later than 5 hours before milking or during milking. Do not contaminate water troughs. Do not use sprays or dip dairy animals in Malathion.

(10/66)

4-1-67

550-45