

NET WEIGHT 80 POUNDS

GUARDSMAN

SUPREME - D

WEED AND FEED FERTILIZER

ACCEPTED
JUL 9 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC PESTICIDE REGISTERED
UNDER NO. 55074 SUBJECT
TO ATTACHED COMMENTS.

3

COMBINATION LAWN FERTILIZER AND WEED KILLER (HERBICIDE)

FOR USE ON ALL KINDS OF WELL ESTABLISHED GRASS LAWNS.

CONTENTS OF THIS PACKAGE COVERS 8000 SQ. FT.

FEEDS LAWNS AND CONTROLS:

Dandelion, Dog Fennel, Lambsquarter, Pigweed, Plantain,

Sheep Sorrel, Buttercup, Chickweed (Common) and ^{Chickweed} House Lar,

Clover, Japanese Clover, Henbit, Mustard, English Lawn Daisy,

Yarrow, Knotweed.

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic Acid*	0.25%
2-(2-Methyl-4-Chlorophenoxy) Propionic Acid**	0.15%
(Dicamba) Dimethylamine Salt of 2-Methoxy-3,6-Dichlorobenzoic Acid***	0.09%

<u>INERT INGREDIENTS:</u>	99.51%
---------------------------	--------

* Equivalent to 0.25% 2,4-D Acid.

** Equivalent to 0.15% MCPA Acid

*** Equivalent to 0.075% (Dicamba) Banvel D Acid ®

U. S. D. A. REG. NO. 550-

CAUTION! KEEP OUT OF REACH OF CHILDREN

May cause skin irritation.

Avoid contact with eyes, skin and clothing.

Avoid breathing dust.

Induce vomiting if swallowed.

Wash hands after use.

GUARANTEED ANALYSIS

Total Nitrogen (N) 15%

BEST DOCUMENT AVAILABLE

10

REAR PANEL

USE PRECAUTIONS:

SUPREME-D WEED AND FEED FERTILIZER should be used for recommended purposes and at recommended rate only.

Do not apply directly to, or come in contact with, vegetables, flowers, grapes, trees or other desirable plants.

Do not apply to areas underlain by roots of desirable trees and shrubs. Do not apply in a brisk wind.

Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of water and containers.

Do not store near fertilizers, seeds, insecticides, fungicides, feed or foodstuffs.

Use of Supreme-D Weed and Feed Fertilizer may cause injury to creeping grasses such as bentgrass, St. Augustine grass, Dichondra, and carpetgrass.

Do not apply to a new lawn. Wait until lawn is 3 months old.

Do not mow lawn for two days prior to application.

Do not water lawn for 24 hours after application.

Make second application after one month if control of hard-to-kill weeds is not complete.

DIRECTIONS FOR APPLICATION:

WHEN TO APPLY: Apply any time weeds are growing actively - spring, summer or fall. Applications to moist foliage and soil when temperature is between 50° and 80° F. are the most effective.

Applications during hot weather (over 80° F.) or on dry soil are apt to cause some temporary damage to turf.

RATE OF APPLICATION: Apply SUPREME-D uniformly at the rate of 10 lbs. to each 1000 sq. ft.

DISTRIBUTED BY

VAN WATERS & ROGERS
SAN FRANCISCO

BEST DOCUMENT AVAILABLE

12

NET WEIGHT 50 POUNDS

GUARANTEE

SUPPLEMENT

WEED AND FELD FERTILIZER



ACCEPTED
JUL 9 1968
550 74 SUBJECT

COMBINATION LAWN FERTILIZER AND WEED KILLER (HERBICIDE)

FOR USE ON ALL KINDS OF WIDE ESTABLISHED GRASS LAWNS.

CONTENTS OF THIS PACKAGE COVER 5000 SQ. FT.

WEEDS LAWN AND CONTROLS:

Dandelion, Red Fennel, Lambsquarter, Figweed, Plantain,

Sheep Sorrel, Buttercup, Chickweed (Common and Mouse Ear),

Clover, Japanese Clover, Henbit, Mustard, English Lawn Daisy,

Yarrow, Knotweed.

ACTIVE INGREDIENTS:

2,4-Dichlorophenoxyacetic Acid*	0.25%
2-(2-Methyl-4-Chlorophenoxy) Propionic Acid**	0.15%
(Dicamba) Dimethylamine Salt of 2-Methoxy-3,6-Dichlorobenzoic Acid***	0.09%

INERT INGREDIENTS: 99.51%

- * Equivalent to 0.25% 2,4-D Acid.
- ** Equivalent to 0.15% MCPA Acid
- *** Equivalent to 0.075% (Dicamba) Lanval D Acid (P)

U. S. D. A. REG. NO. 550-

CAUTION: KEEP OUT OF REACH OF CHILDREN

- May cause skin irritation.
- Avoid contact with eyes, skin and clothing.
- Avoid breathing dust.
- Induce vomiting if swallowed.
- Wash hands after use.

PEST DOCUMENT AVAILABLE

GUARANTEED ANALYSIS

Total Nitrogen (N) 15%

Supreme-D



Use of Supreme-D

Supreme-D is a selective herbicide which is effective against the majority of weeds and grasses in lawns and gardens. It is safe for use on lawns and gardens.

Do not apply to areas underlain by roots of desirable trees and shrubs. Do not apply to areas underlain by roots of undesirable trees and shrubs.

Do not apply to areas underlain by roots of desirable trees and shrubs.

Do not re-entrain any form of water or irrigation water, drainage ditches, or systems of water and sewerage.

Do not use near fertilizers, seeds, insecticides, fungicides, feed or foodstuffs.

Use of Supreme-D Weed and Feed Fertilizer may cause injury to creeping grasses such as Bentgrass, St. Augustine grass, Dichondra, and carpetgrass.

Do not apply to a new lawn. Wait until lawn is 3 months old.

Do not mow lawn for two days prior to application.

Do not water lawn for 24 hours after application.

Make second application after one month if control of hard-to-kill weeds is not complete.

DIRECTIONS FOR APPLICATION:

WHEN TO APPLY: Apply when weeds are growing actively - in spring, summer or fall. Applications to moist foliage and soil when temperature is between 59° and 86° F. are the most effective.

Applications during hot weather (over 86° F.) or on dry soil are apt to cause some temporary damage to turf.

METHOD OF APPLICATION: Apply SUPREME-D uniformly at the rate of 16 lbs. to each 1000 sq. ft.

DISTRIBUTED BY

VAN COTT & FOSBERG
SAN FRANCISCO

20/1/68

550-74

GUARANTEE ANALYSIS (Cont'd)

Available Phosphoric Acid (P_2O_5)
water soluble Potash (K_2O)

See Rear Panel for Additional Precautions

® Banvel D is the registered trademark of

USE INSTRUCTIONS:

SUPREME-D WEED AND FEED FERTILIZER should be used for recommended purposes and at recommended rate only.

Do not apply directly to, or come in contact with, vegetables, flowers, grapes, trees or other desirable plants.

Do not apply to areas underlaid by roots of desirable trees and shrubs. Do not apply in a brisk wind.

Do not contaminate any body of water by direct application, cleaning of equipment, or disposal of water and containers.

Do not store near fertilizers, seeds, insecticides, fungicides, feed or foodstuffs.

Use of Supreme-D Weed and Feed Fertilizer may cause injury to creeping grasses such as bentgrass, St. Augustine grass, Dichondra, and carpetgrass.

Do not apply to a new lawn. Wait until lawn is 3 months old.

Do not mow lawn for two days prior to application.

Do not water lawn for 24 hours after application.

Make second application after one month if control of hard-to-kill weeds is not complete.

DIRECTIONS FOR APPLICATION:

WHEN TO APPLY: Apply any time weeds are growing actively - spring, summer or fall. Applications to moist foliage and soil when temperature is between 50° and 80° F. are the most effective.

Applications during hot weather (over 80° F.) or on dry soil are apt to cause some temporary damage to turf.

RATE OF APPLICATION: Apply SUPREME-D uniformly at the rate of 10 lbs. to each 1000 sq. ft.

DISTRIBUTED BY

VAN WATLES & ROGERS
SAN FRANCISCO

WEED KILLER

NO. 50

ACTIVE INGREDIENTS	21.0%
Sodium Arsenite	16.5%
Cresylic Acid	4.5%
INERT INGREDIENTS	
Water, Soap	79.0%
	100.0%

Total arsenic as metallic (As) all in water soluble form . . . 9.51%

DIRECTIONS FOR USE

**DO NOT USE OR STORE IN OR AROUND THE HOME
DO NOT ALLOW DOMESTIC ANIMALS TO GRAZE TREATED AREA**

GENERAL WEED CONTROL: For use on bare ground to prevent weed growth and for weed control in enclosed industrial areas such as Tank Farms, Lumber storage Yards and Substations.

Do not contaminate water used for irrigation or domestic purposes. Avoid spray drift to desirable plants. Keep pets and livestock off treated areas until washed into soil. Do not apply within the dripline of desirable trees and shrubs.

TIMING: Where weed growth is present, for best results apply when weeds are less than 6 inches high. For treatment of bare ground apply at any time the ground is not frozen.

APPLICATION: Mix the recommended amount with water and apply with sprayer or sprinkling can.

FOR BARE GROUND AND ENCLOSED AREAS: For use in sprayers mix two gallons of this product in 20 gallons of water. DO NOT use more than 360 gallons of this mixture per acre. For use in sprinkling cans mix 1 pint of this product in 2 gallons of water. DO NOT use more than this amount per 160 square feet.

**MAY BE FATAL IF SWALLOWED, INHALED,
OR ABSORBED THROUGH SKIN.
KEEP OUT OF REACH OF CHILDREN.**

Thoroughly wash hands after making or using the solutions. Avoid skin contact and wash it thoroughly after exposure. If on clothing, remove them and wash. Keep children off treated areas until washed into soil. Avoid inhalation of dust or spray mist. Do not get in eyes. Keep away from foodstuffs.

Antidote if taken internally: Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Then give two tablespoonfuls of Epsom Salts or Milk of Magnesia in water and plenty of milk and water. Have victim lie down and keep quiet. Call a physician immediately.

This product is toxic to wild life. Birds and other wildlife in treated areas may be killed. Do not apply when runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of waste. Discard empty containers in a safe place.



L. L. & A. GUIRE

HOLBROOK, MASS.

EPA REG. NO. 551-1