

## REAR PANEL

### USE PRECAUTIONS:

SUPPLEM-X should be used for recommended purposes and at recommended rates.

Limit use to 6 month intervals.

Do not reseed within 4 months after application.

Sweep up spills on walks or patios and hose area.

Do not allow children or pets on treated lawns until SUPPLEM-X has been watered into the soil and the grass is completely dry.

**Avoid application to adjacent desirable plants.**

To avoid possibility of burning foliage, apply only when blades of grass are dry.

Do not apply in a brisk wind.

Destroy empty container. Do not re-use.

### DIRECTIONS FOR APPLICATION

WHEN TO APPLY: To be effective for weed control SUPPLEM-X must be applied before the emergence of weeds. If weeds are visible above the soil, it is too late. To prevent Smooth Crabgrass, hairy Crabgrass, Goosegrass, watergrass, Lambsquarters and Redroot Pigweed apply in late winter or early spring no later than April. To prevent Annual Bluegrass, Renalnettle, and Shepherds-purse apply in early fall or August. Consult State Agricultural Experiment Stations or other turf and garden authorities for specific information. Because the timing of application will vary with local conditions.

RATE OF APPLICATION: Apply 10 lbs SUPPLEM-X uniformly to each 1,000 sq. ft.

HOW TO APPLY: Before applying remove leaves, dead or tall grass, and other debris.

11-18-79  
550-91  
U.S. DEPARTMENT OF AGRICULTURE  
OFFICE OF PEST CONTROL  
REGISTERED  
SUBJECT  
TO ALL COMMENTS.

AG-SEATTLE

**Guard**smam

# SEED GUARD

|                                      |  |
|--------------------------------------|--|
|                                      | WITH TERRACLOR<br>2 lb. EMULSIFIABLE CONCENTRATE |
| <b>GUARANTEED ACTIVE INGREDIENT:</b> |  |
| Pentachloronitrobenzene              | 24.0%  |
| <b>INERT INGREDIENTS:</b>            | 76.0%  |
| <b>Total</b>                         | <b>100.0%</b>                                    |

Contains 2 lbs. of Pentachloronitrobenzene per U. S. gallon at 68° F.

\*TERRACLOR® TRADEMARK AND PRODUCT OF OLIN CORPORATION  
USDA Reg. No 550-91

**WHEAT SEED TREATMENT**  
Recommendations for Wheat Seed Treatment with  
Guardsmam Seed Guard with Terraclor® 2 lb. Emulsifiable Concentrate  
To control common smut or bunt

Apply undiluted directly to the seed at the rate of 2 oz. per bushel or apply in conventional slurry equipment at these rates: For 23cc cup use a mixture of 1/2 gallon per 1 gallon of water; for 46cc cup use a mixture of 1 quart per 1 gallon of water. This will deposit approximately 1/2 ounce active material per bushel of seed. For mist seed treaters, apply 1cc per pound of seed. Adjust trip weight to deposit amount of seed according to cup used. For example: 2.5 pounds of seed per 2 5cc cup, 5 pounds of seed per 5cc cup, etc.

**DO NOT USE TREATED SEED FOR FOOD OR FEED PURPOSES**  
**COMBUSTIBLE LIQUID**  
**KEEP AWAY FROM HEAT AND OPEN FLAME**  
Do not reuse container. Destroy when empty.

**CAUTION**

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid inhaling spray mist. This material may cause skin irritation. Wash hands and face thoroughly after using. Avoid contact with skin, eyes, or clothing. Use with adequate ventilation. Keep container closed. Avoid contamination of feed and foodstuffs.

This product is toxic to fish. Keep out of any body of water.  
Do not contaminate water by cleaning of equipment, or disposal of wastes

NET CONTENTS 5 GALLONS  
@ 68°F (41½ POUNDS)



Distributed by  
**VAN WATNERS & ROGERS**

SAN FRANCISCO, CALIFORNIA 94119

(1/71)

SELLER MAKES NO WARRANTY, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL OR THE USE OF THIS PRODUCT OTHER THAN INDICATED ON THE LABEL. BUYER ASSUMES ALL RISK OF USE AND/OR HANDLING OF THIS MATERIAL WHEN SUCH USE AND/OR HANDLING IS CONTRARY TO LABEL INSTRUCTIONS.

Printed in U.S.A.