

ACCEPTED
 7-17-72
 550-94
 UNDER THE FEDERAL FOOD, DRUG & COSMETIC ACT
 FOR LABELING PURPOSES
 TO ATTAIN



AAAD
 2010

Supreme-D weed & feed 15-3-6

GUARANTEED FERTILIZER ANALYSIS

TOTAL NITROGEN (N)		15%
Ammonic Nitrogen	1%	
Water Soluble Nitrogen	10%	
Water Insoluble Nitrogen	4%	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)		3%
WATER SOLUBLE POTASH (K ₂ O)		6%
MAGNESIUM (Mg)		1.5%
IRON (Fe)		0.5%

DERIVED FROM: Ammonium Phosphate, Superphosphate, Urea, Ureaformaldehyde, Muriate of Potash, Ferrous Sulfate, Dolomite.

GENERAL INFORMATION

This product contains Trimec[®] herbicide, a new herbicidal formulation designed to keep the herbicide in the weed zone on the surface of a 15-3-6 lawn fertilizer. The weed killing benefits from SUPREME-D are especially important because excellent control results from minimum amounts of chemical. However, if adverse low moisture conditions prevail and weeds are in a state of poor growth, possible retreatment may be necessary when conditions improve.

DIRECTIONS FOR USE

WHEN TO APPLY: For best results apply when weeds are growing vigorously in the spring or early fall after heavy rains.

This material may be used on bluegrass, fescue, rye, Bermuda and zoysia lawns. Normally, at this time there is sufficient moisture for excellent response from the fertilizer as well as very effective weed control.

SUPREME-D is a remarkably effective lawn food which combines the advantages of both fast and long lasting feeding action plus the broad range of weed control of the Trimec[®] herbicide. The fertilizer produces a thick, green turf while killing weeds to make way for turf growth. Weeds contacted by the product stop growing, gradually die and disappear from the lawn.


— COVERS 5,000 SQ. FT. —

HOW MUCH TO USE: This 30 lb. bag of SUPREME-D should be used on 5,000 square feet. Apply evenly with a lawn spreader either as a one-pass application or a two-pass application, criss-crossing the area at half the rate each time.

To get maximum weed killing results, do not rake, mow or water the lawn for at least 24 hours after spreading the product.

Lawns which are heavily overrun by weeds and are in poor condition may require more than one application of SUPREME-D. A suggested second application could be made in approximately 30 days.

CAUTION: Do not use SUPREME-D where desirable clovers are present or allow the product to "wind drift" on flowers, vegetables or ornamental shrubbery. Do not apply to newly seeded lawns. During application close spreader openings when turning. Apply when the air is calm. Keep away from fertilizers, seeds, insecticides, fungicides, feed or foodstuffs. Do not apply in areas underlaid by roots of desirable trees and shrubs. Do not reuse container. Destroy by burying in a safe place. Do not burn.



8



Supreme-D weed & feed 15*3*6

PLUS - TRIMEC HERBICIDAL CONCENTRATE

FEEDS LAWNS AND CONTROLS: Dandelion, Lambsquarter, Pigweed, Plantain, Sheep Sorrel, Thistle, Chickweed, Common and Mouse Ear, Clover, Purslane, Henbit, Mustard, Yarrow, Knotweed, Dock, Daisy.

ACTIVE INGREDIENTS

2,4-Dichlorophenoxyacetic Acid 0.4856
 2-(2-Methyl-4-Chlorophenoxy) Propionic Acid 0.2647
 Dicamba (3,6-Dichloro-o-anisic Acid) 0.0434
INERT INGREDIENTS 99.2663

EQUIVALENT TO:

2,4-Dichlorophenoxyacetic Acid 1.266 lbs. per acre
 2-(2-Methyl-4-Chlorophenoxy) Propionic Acid 0.534 lbs. per acre
 3,6-Dichloro-o-anisic Acid 0.113 lbs. per acre

TOTAL 100.0000

U. S. Patent 3,284,186

CAUTION! - KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Induce vomiting if swallowed. Avoid inhaling dust or contact with skin, eyes and clothing. Wash hands after use.

FOR ADDITIONAL PRECAUTIONS, DIRECTIONS FOR USE, AND FERTILIZER GUARANTEE, SEE REVERSE SIDE.

TRIMEC is a Registered Trademark of PBI Gordon Corp.

COVERS 5000 SQ. FT. ■ NET WT. 30 LBS.



DISTRIBUTED BY
VAN WATERS & ROGERS
 SAN FRANCISCO, CALIF. 94119

NOTE: USER MAKES NO WARRANTY, EXPRESS OR IMPLIED, FOR THE MERCHANTABILITY OF THIS MATERIAL, OR FOR THE RESULTS TO BE OBTAINED FROM THE USE OF THIS MATERIAL, WHEN SUCH USE AND/OR HANDLING IS CONTRARY TO THE LABEL INSTRUCTIONS.

7-17-72
 550-91A

SIDE

2
1.2047



GR
MARK

RED GREEN BLACK 29-1.1160

CORNER



VIGORO

VIGORO LAWN

Contains dimethylamine

Kills Dandelions, Chickweed, Clover, Creeping Beggarweed, Pennywort, and over 100 other common weeds. Read Directions Before Using. Excessive run-off or wetting of the soil may result in loss of turf. Deep-rooted perennial weeds may require several days after mowing. Do not mow for 24 hours.

FOR HOSE END SPRAYERS: Fill bottle to level indicated in chart below.

FILL BOTTLE TO MARK:
5 gal.
10 gal.
15 gal.
20 gal. full

Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals and condition of application equipment will vary, seller makes no claims other than that stated on this label.

TO CONTROL WILD ONION

Add 3 tablespoons of VIGORO LAWN WEED KILLER to a gallon of water and mix well. Spray weeds without excessive run-off or wetting of the soil, using 1 gallon of solution to about 200 sq. ft. of turf.

Application should be made in the very early spring when wild onion is not more than two inches high. Repeat applications should be made as new spikes appear.

LAWN WEED KILLER

SURE DEATH TO DANDELIONS AND MOST BROADLEAF WEEDS - KEEP FROM FREEZING -

ACTIVE INGREDIENTS:

- *Dimethylamine salt of 2,4-dichlorophenoxyacetic acid 62.7%
- **Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy)propionic acid 2.84%
- **Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid) 0.65%

INERT INGREDIENTS 90.24%

- Total 100.00%
- *2,4-dichlorophenoxyacetic acid equivalent 0.45 pounds per gallon or 5.2%
- **2-(2-methyl-4-chlorophenoxy)propionic acid equivalent 0.20 pounds per gallon or 2.35%
- **3,6-dichloro-o-anisic acid equivalent 0.04 pounds per gallon or 0.5%

HARMFUL IF SWALLOWED. AVOID CONTACT. IN CASE OF CONTACT, FLUSH WITH WATER. MEDICAL ATTENTION.

Do not burn poison ivy as smoke contact with the skin. Do not store near shrubs, and other desirable crop plants. Avoid drift or spray mist onto desirable white clover, bent grass and St. Augustine grass. Use separate sprayer for VIGORO LAWN WEED KILLER. Use thoroughly with household ammonia and coarse sprays are less likely to drift onto desirable trees and shrubs.

WARNING: Keep Out of Reach of Children
See Back panel for additional Warnings and Cautions

NET CONTENTS 32 FL. OZ. (1 QT.)

RED GREEN BLACK

1/16" PLAIN TIN
3/32" FOLD

ACCEPTED
4-7-71
55A-1769

ENTER THE
FOLLOWING TO IDENTIFY
FOR RECORDING PURPOSES
ED. ORDER NO.

CUSTOMER	GORDON CORP	ENG 29-1.1160	ICBY 008	SIZE 409X206X614	4
LAYOUT	INDEX NO N-55	D.E. 9-24-71	CE 12.560X7.0	SIZ 208L.5MD. TOP	5
COLORS	RED • GREEN • BLACK ON WHITE COAT			ARTIST 904.1	4

14



STACK

FRONT

CORN H

CORN H

VIGORO

VIGORO LAWN WEED KILLER

Contains dimethylamine salts of 2,4-D, MCPP and Dicamba

Kills Dandelions, Chickweed, Clover, Buckhorn, Ragweed, Plantain, Heal-all, Pigweed, Burdock, Creeping Jenny, Wild Onion, Poison Oak, Poison Ivy, Honeysuckle, Creeping Beggarweed, Pennywort, Creeping Charles, Wandering Jew, Careless Weed, and over 100 other common broadleaf weeds.

Read Directions Before Using: Add 3 tablespoonfuls or 1 1/2 oz. of VIGORO LAWN WEED KILLER to 1 gallon of water and mix well. Spray weeds without excessive run-off or wetting of the soil, using 1 gallon of spray mixture to 275 square feet of turf. Deep-rooted perennial weeds may require repeat application. For best results, apply when weeds are growing actively, when there is ample soil moisture, and several days after mowing. Do not use on Dichondra lawns, Carpet grass, and Centipede grass.

FOR HOSE END SPRAYERS: Pour VIGORO LAWN WEED KILLER into sprayer bottle to level indicated in chart below. Do not add water.

ACTIVE INGREDIENTS:

- *Dimethylamine salt of 2,4-dichlorophenoxyacetic acid 6.27%
- **Dimethylamine salt of 2-(2-methyl-4-chlorophenoxy) propionic acid 2.84%
- ***Dimethylamine salt of Dicamba (3,6-dichloro-o-anisic acid) 0.65%

INERT INGREDIENTS 90.24%

Total 100.00%

- *2,4-dichlorophenoxyacetic acid equivalent 0.45 pounds per gallon or 5.2%
- **2-(2-methyl-4-chlorophenoxy) propionic acid equivalent 0.20 pounds per gallon or 2.35%
- ***3,6-dichloro-o-anisic acid equivalent 0.15 pounds per gallon or 1.8%

FILL BOTTLE TO MARK:

- 5 gal.
- 10 gal.
- 15 gal.
- 20 gal. full

AREA TO SPRAY:

- 1500 sq. ft.
- 3000 sq. ft.
- 4500 sq. ft.
- 6000 sq. ft.

WARNING

HARMFUL IF SWALLOWED. AVOID CONTACT WITH SKIN, EYES, AND CLOTHING. IN CASE OF CONTACT, FLUSH WITH PLENTY OF WATER. FOR EYES GET MEDICAL ATTENTION.

Do not burn poison ivy as smoke from the burning vine is very poisonous on contact with the skin. Do not store near fertilizers, seeds, insecticides, or fungicides.

Avoid drift or spray mist onto vegetables and flowers, ornamental trees and shrubs, and other desirable crop plants. Do not use on newly seeded lawns, desirable white clover, bent grass and St. Augustine grass (except for spot spraying). Use separate sprayer for VIGORO LAWN WEED KILLER, or clean all parts thoroughly with household ammonia and hot water before using for any other purpose. Coarse sprays are less likely to drift. Do not apply to area underlaid by roots of desirable trees and shrubs.

Do not reuse this container. When container is empty, wash thoroughly, and recycle.

DISTRIBUTED BY

SWIFT AGRICULTURAL CHEMICALS CORP.

CHICAGO, ILLINOIS 60606

Keep Out of Reach of Children

Read and Follow All Directions, Warnings, and Cautions

CONTENTS 32 FL. OZ. (1 QT.)

FORMER	GORDON CORP.	ENG 29-1160	FACTY. 008	SIZE 409X206X614
OUT	INDEX NO. N-55	913-24-71	CL. 12.560X7.0	STYLE OBL. SHD. TOP
COBS	RED • GREEN • BLACK ON WHITE COAT			ARTIST 9041

KEY

3/16" FOLD

15

14



ACCEPTED

7-17-72
550-94



Supreme-D weed

GUARANTEED FERTIL

TOTAL NITROGEN (N)
Ammonic Nitrogen
Water Soluble Nitrogen
Water Insoluble Nitrogen
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)
WATER SOLUBLE POTASH (K ₂ O)
MAGNESIUM (Mg)
IRON (Fe)

DERIVED FROM: Ammonium Phosphate, Superphosphate, Ferrous Sulfate, Dolomite.

GENERAL INFO

This product contains Trimec[®] herbicide, a new herbicide in the weed zone on the surface of a 15-3-6 lawn fertilizer. The weed killing benefits from SUPREME-D are especially from minimum amounts of chemical. However, if weeds are in a state of poor growth, possible retreatment may be necessary.

DIRECTIONS

WHEN TO APPLY: For best results apply when weeds are young and fall after heavy rains.

This material may be used on bluegrass, fescue, rye, and other cool season grasses at any time there is sufficient moisture for excellent response and weed control.

SUPREME-D is a remarkably effective lawn food which provides long lasting feeding action plus the broad range of weed control. The fertilizer produces a thick, green turf while killing weeds. Weeds that do not die by the product stop growing, gradually die and disappear.

— COVERS 5,000

HOW MUCH TO USE: This 30 lb. bag of SUPREME-D should be applied evenly with a lawn spreader either as a one-pass application or by spreading the area at half the rate each time.

To get maximum weed killing results, do not rake, mow, or water lawns after spreading the product.

Lawns which are heavily overrun by weeds and are in poor condition should receive application of SUPREME-D. A suggested second application should be made in 4 to 6 weeks.

CAUTION: Do not use SUPREME-D where desirable plants are in bloom. Avoid "drift" on flowers, vegetables or ornamental shrubbery. Close spreader openings when turning. Apply SUPREME-D around trees, shrubs, flower beds, lawns, and lawns. Do not use around desirable trees and shrubs. Do not reuse container.

7-17-72
550-94



Supreme-D weed & feed 15-3-6

GUARANTEED FERTILIZER ANALYSIS

TOTAL NITROGEN (N)		15%
Ammoniac Nitrogen	1%	
Water Soluble Nitrogen	10%	
Water Insoluble Nitrogen	4%	
AVAILABLE PHOSPHORIC ACID (P ₂ O ₅)		3%
WATER SOLUBLE POTASH (K ₂ O)		6%
MAGNESIUM (Mg)		1.5%
IRON (Fe)		0.5%

DERIVED FROM: Ammonium Phosphate, Superphosphate, Urea, Ureaformaldehyde, Muriate of Potash, Ferrous Sulfate, Dolomite

GENERAL INFORMATION

This product contains Trimec herbicide, a new herbicidal formulation designed to keep the herbicide in the weed zone on the surface of a 15-3-6 lawn fertilizer.

The weed killing benefits from SUPREME D are especially important because excellent control results from minimum amounts of chemical. However, if adverse low moisture conditions prevail and weeds are in a state of poor growth, possible retreatment may be necessary when conditions improve.

DIRECTIONS FOR USE

WHEN TO APPLY For best results apply when weeds are growing vigorously in the spring or early fall after heavy rains.

This material may be used on bluegrass, fescue, ryegrass, Bermuda and zoysia lawns. Normally, at this time there is sufficient moisture for excellent response from the fertilizer as well as very effective weed control.

SUPREME D is a remarkably effective lawn food which combines the advantages of both fast and long lasting feeding action plus the broad range of weed control of the Trimec herbicide. The fertilizer produces a thick green turf while killing weeds to make way for turf growth. Weeds contacted by the product stop growing, gradually die and disappear from the lawn.

— COVERS 5,000 SQ. FT.

HOW MUCH TO USE This 30 lb. bag of SUPREME D should be used on 1,000 square feet. Apply evenly with a lawn spreader either as a one pass application or a two pass application, criss crossing the area at half the rate each time.

To get maximum weed killing results, do not rake, mow or water the lawn for at least 24 hours after spreading the product.

Lawns which are heavily overrun by weeds and are in poor condition may require more than one application of SUPREME D. A suggested second application could be made in approximately 30 days.

CAUTION Do not use SUPREME D where desirable clovers are present or allow the product to wind drift on flowers, vegetables or ornamental shrubbery. Do not apply to newly seeded lawns. During application close spreader openings when turning. Apply when the air is calm. Keep away from fertilizers, seeds, insecticides, fungicides, feed or foodstuffs. Do not apply in areas underlaid by roots of desirable trees and/or shrubs. Do not reuse container. Destroy by burying in a safe place. Do not burn.