

Mr. Dennis Prober
Van Waters & Rogers
Division of Univar
2256 Junction Avenue
San Jose, CA 95131

345/122559
17 | 2

16 JUL 1984

ZRP

300/124352
17 | 3

Dear Mr. Prober:

Subject: Amendments - Alternate Sources; Update Label
Guardman Diazinon 4E
EPA Registration No. 550-154
Your Application Dated May 7, 1984

The amendments referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), are acceptable. The formula statement adding the alternate sources of diazinon has been added to this product's registration file. A stamped copy of the labeling bearing the revisions requested in the Notice of Registration is enclosed for your records. It is understood that the appropriate revisions regarding the Storage and Disposal statements, the declaration of the solvent, and farmworker safety will be at the next label printing.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

RD/IRB:LaRocca:DCR-04490:WANG-0027L:eg:Raven:479-2013:6/29/84

BEST DOCUMENT AVAILABLE

ACCEPTED
with COMMENTS
in EPA Label 103-11

JUL 16 1994



Under the
Insecticide
Act and
Registered
550-154

Diazinon® 4E

Only For Sale To, Use and Storage By Pest Control Operators and Other Trained Personnel In Controlling Certain Fruit, Nut, Vegetable, Forage, Forest, Turf, and Ornamental Pests as Listed on the Labeling.

ACTIVE INGREDIENTS:

O, O-diethyl-O-(2-isopropyl-6-methyl-4-pyrimidyl) phosphorothioate	47.5%
Xylene Range Aromatic Solvent	35.1%
INERT INGREDIENTS:	17.4%

Contains 4 pounds of Diazinon per gallon

KEEP OUT OF REACH OF CHILDREN
WARNING
STATEMENT OF PRACTICAL TREATMENT

Do not induce vomiting

First Aid:
If on skin: Wash with plenty of water.
If in eyes: Flush with plenty of water.

NOTE TO PHYSICIAN



Manufactured By
Van Waters & Rogers
division of Univar

BEST DOCUMENT AVAILABLE

05069490

(99-9-81)

EPA Reg. No. 550-154-AA

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
WARNING

Caution: May be harmful if swallowed. Causes eye, nose, and throat irritation. May irritate the respiratory system. Avoid breathing dusts and fumes. Wash thoroughly after handling. Do not get into eyes, nose or mouth. Avoid contact with skin and clothing. Do not eat, drink, or use tobacco products while handling this product. Do not use near open flames or fire. Do not use in confined spaces. Do not use in areas where people, animals, or birds are present. Do not use in areas where people, animals, or birds are present.

DIRECTIONS FOR USE

Fruit and Nut Crops

Note: Treat a fruit or nut crop with Diazinon 4E at a 5-day reentry interval for 10 days after application. For grapes, peaches, and nectarines, five days of reentry is sufficient. Between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage, when a mixture of two or more organic phosphorus pesticides is applied, the reentry interval shall be prolonged by adding to the reentry interval an additional 50% of that interval.

ALMONDS (12 pt)

Mix 1.25 lb Diazinon 4E with 1.25 gal water per 100 gal of water. Apply 1.25 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 1.25 gal per 100 gal of water and apply as a drench to the soil.

APRICOTS (8 pt)

Mix 1 lb Diazinon 4E with 1 gal water per 100 gal of water. Apply 1 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 1 gal per 100 gal of water. Apply when the insects first appear.

BLUEBERRIES (2 pt)

Mix 0.25 lb Diazinon 4E with 0.25 gal water per 100 gal of water. Apply 0.25 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 0.25 gal per 100 gal of water.

CANEBERRIES (2 pt)

Mix 0.25 lb Diazinon 4E with 0.25 gal water per 100 gal of water. Apply 0.25 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 0.25 gal per 100 gal of water. Apply when the insects first appear.

CHERRIES (8 pt)

Mix 1 lb Diazinon 4E with 1 gal water per 100 gal of water. Apply 1 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 1 gal per 100 gal of water. Apply when the insects first appear.

CITRUS FRUITS (20 pt)

Mix 2 lb Diazinon 4E with 2 gal water per 100 gal of water. Apply 2 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 2 gal per 100 gal of water. Apply when the insects first appear.

CRANBERRIES (8 pt)

Mix 1 lb Diazinon 4E with 1 gal water per 100 gal of water. Apply 1 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 1 gal per 100 gal of water. Apply when the insects first appear.

FIGS (8 pt)

Mix 1 lb Diazinon 4E with 1 gal water per 100 gal of water. Apply 1 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 1 gal per 100 gal of water. Apply when the insects first appear.

FILBERTS (4 pt)

Mix 0.5 lb Diazinon 4E with 0.5 gal water per 100 gal of water. Apply 0.5 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 0.5 gal per 100 gal of water. Apply when the insects first appear.

GRAPES (2 pt)

Mix 0.25 lb Diazinon 4E with 0.25 gal water per 100 gal of water. Apply 0.25 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 0.25 gal per 100 gal of water. Apply when the insects first appear.

OLIVES (24 pt)

Mix 3 lb Diazinon 4E with 3 gal water per 100 gal of water. Apply 3 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 3 gal per 100 gal of water. Apply when the insects first appear.

PEACHES, NECTARINES (8 pt)

Mix 1 lb Diazinon 4E with 1 gal water per 100 gal of water. Apply 1 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 1 gal per 100 gal of water. Apply when the insects first appear.

PECANS (5 pt)

Mix 0.5 lb Diazinon 4E with 0.5 gal water per 100 gal of water. Apply 0.5 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 0.5 gal per 100 gal of water. Apply when the insects first appear.

PINEAPPLES (4 qt)

Mix 0.5 lb Diazinon 4E with 0.5 gal water per 100 gal of water. Apply 0.5 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 0.5 gal per 100 gal of water. Apply when the insects first appear.

PLUMS, PRUNES (8 pt)

Mix 1 lb Diazinon 4E with 1 gal water per 100 gal of water. Apply 1 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 1 gal per 100 gal of water. Apply when the insects first appear.

STRAWBERRIES (2 pt)

Mix 0.25 lb Diazinon 4E with 0.25 gal water per 100 gal of water. Apply 0.25 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 0.25 gal per 100 gal of water. Apply when the insects first appear.

WALNUTS (3 qt)

Mix 0.375 lb Diazinon 4E with 0.375 gal water per 100 gal of water. Apply 0.375 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 0.375 gal per 100 gal of water. Apply when the insects first appear.

BEST DOCUMENT AVAILABLE

Vegetable Foliar Insects

BEANS (12 pt)

Mix 1.25 lb Diazinon 4E with 1.25 gal water per 100 gal of water. Apply 1.25 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 1.25 gal per 100 gal of water. Apply when the insects first appear.

CUMBER

Mix 1.25 lb Diazinon 4E with 1.25 gal water per 100 gal of water. Apply 1.25 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 1.25 gal per 100 gal of water. Apply when the insects first appear.

EGGPLANT

Mix 1.25 lb Diazinon 4E with 1.25 gal water per 100 gal of water. Apply 1.25 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 1.25 gal per 100 gal of water. Apply when the insects first appear.

PEPPER

Mix 1.25 lb Diazinon 4E with 1.25 gal water per 100 gal of water. Apply 1.25 gal per 100 gal of water. Apply when the insects first appear. Repeat applications as necessary. Do not apply more than 1.25 gal per 100 gal of water. Apply when the insects first appear.

ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining in the treated area.

PHYSICAL OR CHEMICAL HAZARDS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

To be applied only by or under the supervision of pest control operators or other trained personnel responsible for pest control programs.

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage, disposal of wastes, or cleaning of equipment. Open dumping is prohibited. Do not reuse container.

PESTICIDE DISPOSAL: Pesticides, spray mixture or residue that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

CONTAINER DISPOSAL: A) Reseal container and offer for recycling, reuse, or sale. B) Triple rinse (or equivalent) and offer for recycling, reuse, or sale in an approved landfill or bury in a safe place.

CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES.

Conditions of Sale and Warranty

The Direct Insecticide... Seller makes no warranty, express or implied, concerning the material or the use of this product other than indicated on the label. Buyer assumes all risk of use and repairs caused by material which does not conform to the label or is contrary to label directions.

General Information

Guard... intended... the direct... each applic... Maximum...
Maximum...
Maximum...

... water per acre to the vegetation...

PLUMS, PRUNES (8 pt)

Apply... water per acre to the vegetation...

STRAWBERRIES (2 pt)

Apply... water per acre to the vegetation...

WALNUTS (3 qt)

Apply... water per acre to the vegetation...

BEST DOCUMENT AVAILABLE

Vegetable Foliar Insects

... water per acre to the vegetation...

BEANS (1-12 pt)

Apply... water per acre to the vegetation...

LIMA BEANS (1 pt)

Apply... water per acre to the vegetation...

BROCCOLI, CABBAGE, CAULIFLOWER (1 pt)

Apply... water per acre to the vegetation...

BRUSSELS SPROUTS (1 pt)

Apply... water per acre to the vegetation...

CARROTS (1 pt)

Apply... water per acre to the vegetation...

CELERY (1 pt)

Apply... water per acre to the vegetation...

COLLARDS, KALE, SWISS CHARD, TURNIP TOPS (1 pt)

Apply... water per acre to the vegetation...

CUCUMBERS (1 pt)

Apply... water per acre to the vegetation...

LETTUCE, ENDIVE (1 pt)

Apply... water per acre to the vegetation...

MELONS, SQUASH (1-12 pt)

Apply... water per acre to the vegetation...

MUSHROOM HOUSES

Apply... water per acre to the vegetation...

ONIONS (1 pt)

Apply... water per acre to the vegetation...

PEAS (1 pt)

Apply... water per acre to the vegetation...

PEPPERS (1-2 pt)

Apply... water per acre to the vegetation...

POTATOES (8 pt)

Apply... water per acre to the vegetation...

RADISHES, PARSNIPS, TURNIPS (1 pt)

Apply... water per acre to the vegetation...

SPINACH, BEETS (1 pt)

Apply... water per acre to the vegetation...

SQUASH (Summer and Winter)

Apply... water per acre to the vegetation...

7

SWEET CORN (1-1 1/4 pts)

Apply 1/2 pt per acre of this product to sweet corn plants at the time of silking. This product is effective against the European corn borer, corn earworm, and other insects that feed on the silks and developing ears.

TOMATOES (1-1 1/2 pts)

Apply 1/2 pt per acre of this product to tomato plants at the time of flowering. This product is effective against the tomato pinworm, tobacco hornworm, and other insects that feed on the leaves and fruit.

WATERCRESS (Hawaii only) (1 pt)

Apply 1/2 pt per acre of this product to watercress plants at the time of flowering. This product is effective against the watercress root miner and other insects that feed on the roots.

Vegetable Soil Insects

This product is effective against soil-dwelling insects that damage vegetable roots, including wireworms, cutworms, and mole crickets.

CUTWORMS (Surface and Subterranean)

Apply 1/2 pt per acre of this product to vegetable plants at the time of flowering. This product is effective against cutworms that feed on the leaves and roots.

GARDEN SYMPHYLANS (Centipedes)

Apply 1/2 pt per acre of this product to vegetable plants at the time of flowering. This product is effective against garden symphylans that damage roots.

MOLE CRICKETS

Apply 1/2 pt per acre of this product to vegetable plants at the time of flowering. This product is effective against mole crickets that damage roots.

ONION MAGGOTS

Apply 1/2 pt per acre of this product to onion plants at the time of flowering. This product is effective against onion maggots that damage roots.

ROOT MAGGOTS

Apply 1/2 pt per acre of this product to vegetable plants at the time of flowering. This product is effective against root maggots that damage roots.

WIREWORMS

Apply 1/2 pt per acre of this product to vegetable plants at the time of flowering. This product is effective against wireworms that damage roots.

SOUTHERN POTATO WIREWORMS (Southeastern U.S.)

Apply 1/2 pt per acre of this product to potato plants at the time of flowering. This product is effective against southern potato wireworms that damage roots.

Field and Forage Foliar Insects

This product is effective against foliar-dwelling insects that damage field and forage crops, including alfalfa, clover, and corn.

ALFALFA CLOVER (3 pts)

Apply 1/2 pt per acre of this product to alfalfa and clover plants at the time of flowering. This product is effective against alfalfa cutworms and other insects that feed on the leaves.

CLOVER (8 pts)

Apply 1/2 pt per acre of this product to clover plants at the time of flowering. This product is effective against clover cutworms and other insects that feed on the leaves.

CORN (2 1/2 pts)

Apply 1/2 pt per acre of this product to corn plants at the time of flowering. This product is effective against corn earworms and other insects that feed on the leaves and ears.

COTTON (2 pts)

Apply 1/2 pt per acre of this product to cotton plants at the time of flowering. This product is effective against cotton bollworms and other insects that feed on the leaves and bolls.

GUAR (Texas and Oklahoma only) (1 pt)

Apply 1/2 pt per acre of this product to guar plants at the time of flowering. This product is effective against guar cutworms and other insects that feed on the leaves.

HOPS (2 pts)

Apply 1/2 pt per acre of this product to hop plants at the time of flowering. This product is effective against hop cutworms and other insects that feed on the leaves.

SORGHUM (1 pt)

Apply 1/2 pt per acre of this product to sorghum plants at the time of flowering. This product is effective against sorghum cutworms and other insects that feed on the leaves.

SUGAR BEETS (1 pt)

Apply 1/2 pt per acre of this product to sugar beet plants at the time of flowering. This product is effective against sugar beet cutworms and other insects that feed on the leaves.

SUGARCANE (1 pt)

Apply 1/2 pt per acre of this product to sugarcane plants at the time of flowering. This product is effective against sugarcane cutworms and other insects that feed on the leaves.

TOBACCO (1 pt)

Apply 1/2 pt per acre of this product to tobacco plants at the time of flowering. This product is effective against tobacco hornworms and other insects that feed on the leaves.

Field and Forage Soil Insects

This product is effective against soil-dwelling insects that damage field and forage crops, including wireworms, cutworms, and mole crickets.

TOBACCO (4 qts)

Apply 1/2 pt per acre of this product to tobacco plants at the time of flowering. This product is effective against tobacco hornworms and other insects that feed on the leaves.

Range, Pasture, and Grassland Insects

This product is effective against insects that damage range, pasture, and grassland crops, including cutworms and mole crickets.

BERMUDAGRASS AND FORAGE GRASSES (other than Range and Pasture Grasses) (1 pt)

Apply 1/2 pt per acre of this product to bermudagrass and forage grasses at the time of flowering. This product is effective against cutworms and other insects that feed on the leaves.

RANGELAND, PASTURE DITCH BANKS, ROADSIDES, WASTELAND, NONCROP AREAS, BARRIER STRIPS

Apply 1/2 pt per acre of this product to rangeland, pasture ditch banks, roadsides, wasteland, noncrop areas, and barrier strips at the time of flowering. This product is effective against cutworms and other insects that feed on the leaves.

Forest Insects

PINE (Seed Orchards, Plantations, Forests)

Apply 1/2 pt per acre of this product to pine trees at the time of flowering. This product is effective against pine sawflies and other insects that feed on the needles.

BEST DOCUMENT AVAILABLE

Ornamental Insects

This product is effective against insects that damage ornamental plants, including cutworms and mole crickets.

	Control	Rate
1. Alfalfa	1/2 pt	3 pts
2. Clover	1/2 pt	8 pts
3. Corn	1/2 pt	2 1/2 pts
4. Cotton	1/2 pt	2 pts
5. Guar	1/2 pt	1 pt
6. Hops	1/2 pt	2 pts
7. Sorghum	1/2 pt	1 pt
8. Sugar Beets	1/2 pt	1 pt
9. Sugarcane	1/2 pt	1 pt
10. Tobacco	1/2 pt	1 pt

Mimosa Webworms

Apply 1/2 pt per acre of this product to mimosa trees at the time of flowering. This product is effective against mimosa webworms that feed on the leaves.

Lawn Insects

This product is effective against insects that damage lawns, including cutworms and mole crickets.

	Control	Rate
1. Alfalfa	1/2 pt	3 pts
2. Clover	1/2 pt	8 pts
3. Corn	1/2 pt	2 1/2 pts
4. Cotton	1/2 pt	2 pts
5. Guar	1/2 pt	1 pt
6. Hops	1/2 pt	2 pts
7. Sorghum	1/2 pt	1 pt
8. Sugar Beets	1/2 pt	1 pt
9. Sugarcane	1/2 pt	1 pt
10. Tobacco	1/2 pt	1 pt

Boxelder Bugs

Apply 1/2 pt per acre of this product to boxelder trees at the time of flowering. This product is effective against boxelder bugs that feed on the leaves.

Hyperodes Weevil (New York only)

Apply 1/2 pt per acre of this product to hyperodes weevils at the time of flowering. This product is effective against hyperodes weevils that damage roots.

B. Bugs, White Grubs (such as Japanese Beetle Larvae, European Chafar, Southern Chafar)

Apply 1/2 pt per acre of this product to b. bugs, white grubs, and other insects that damage roots.

Fire Ants

Apply 1/2 pt per acre of this product to fire ant colonies at the time of flowering. This product is effective against fire ants that damage roots.

Note

NAMECO Van Waters & Rogers

1000 University Ave. New York, N.Y. 10017

GUAR (Texas and Oklahoma only) (1 pt)
 HOPS (2 pts)
 SORGHUM (1 pt)
 SUGAR BEETS (1 pt)
 SUGARCANE (1 pt)
 TOBACCO (1 pt)

Field and Forage Soil Insects

ALFALFA, CLOVER (4 qts)
 CORN (4 qts)
 COWPEAS (4 qts)
 LESPEDEZA (4 qts)
 PEANUTS (4 qts)
 SORGHUM (4 qts)
 SOYBEANS (4 qts)
 SUGAR BEETS (1 gal)
 SUGARCANE (4 qts)


Hyperodes Weevil (New York only)
 Billbugs White Grubs (such as Japanese Beetle Larvae European Chaffer Southern Chaffer)

Fire Ants

Note

Registered Trademark of CIBA GEIGY Corporation

NAMCO trademark of Van Waters & Rogers
 Division of Union



BEST DOCUMENT AVAILABLE

AGRICULTURAL CHEMICAL
Do not ship or store with food, feeds or clothing

05075299-CA

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
WARNING**

May be fatal if swallowed. Do not swallow. May be absorbed through skin. May be dangerous to eyes and skin. Do not get in eyes or on skin. Wash thoroughly after handling and before smoking or eating. Wash contaminated clothing before reuse. Avoid breathing vapors and spray mist. Handle concentrate in a ventilated area. Avoid contamination of food, food utensils, domestic water, or feedstuffs.

Do not use on humans, pets or livestock. Do not apply if unprotected persons or animals, including children or pets, are in the area to be treated. Cover aquariums. Do not contaminate ornamental fishponds. Do not allow unprotected persons or animals to re-enter treated areas until spray has dried. When used in dwellings, care should be taken to avoid deposits which could be frequently contacted by children.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply to water that will be used for recreational purposes and human and live stock consumption. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. This product must not be used in areas where impact on threatened and endangered species is likely. Contact your State Fish and Game Agency before applying this product. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds flowering in the treated area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

PHYSICAL OR CHEMICAL HAZARDS

COMBUSTIBLE MIXTURE. Do not use, pour, spray, or store in heat or open flame. Keep out of fire.

DIRECTIONS FOR USE

If application of this material for use of this product is required, it must be done with the following: To be applied only by or under the supervision of pest control operators or other trained personnel responsible for pest control programs

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage, disposal, or waste. Do not use equipment or containers previously used for other pesticides. Do not use for other purposes.

PESTICIDE DISPOSAL: Pesticides should be disposed of in accordance with the following: when empty, triple rinse and dispose of in a safe manner. Do not use for other purposes. Do not contaminate water, food, or feed.

CONTAINER DISPOSAL: Wash, reuse, or recycle. Do not use for other purposes. Do not use for other purposes. Do not use for other purposes. Do not use for other purposes.

CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES

See attached pamphlet for specific use directions and precautions. SELLER MAKES NO WARRANTY EXPRESSED OR IMPLIED CONCERNING THIS MATERIAL OR THE USE OF THIS PRODUCT OTHER THAN INDICATED ON THE LABEL. BUYER ASSUMES ALL RISK OF USE AND OR HANDLING OF THIS MATERIAL WHEN SUCH USE OR HANDLING IS CONTRARY TO LABEL INSTRUCTIONS

 **Guardsman***
Diazinon 4E**

Only For Sale To, Use and Storage By Pest Control Operators and Other Trained Personnel In Controlling Certain Fruit, Nut, Vegetable, Forage, Forest, Turf, and Ornamental Pests as Listed on the Labeling.

ACTIVE INGREDIENTS:

O, O-diethyl-O-(2-isopropyl-6-methyl-4-pyrimidyl) phosphorothioate 47.5%
Xylene Range Aromatic Solvent 35.1%

INERT INGREDIENTS:

17.4%

Contains 4 pounds of Diazinon per gallon

* trademark of Univar Corp.

** trademark of CIBA-GEIGY Corp.

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Get medical aid immediately. Induce vomiting if the patient is conscious and if water is available. Do not induce vomiting unless other treatment is not available. Vomiting may cause aspiration pneumonia. If this reflex is likely to occur, give one or two glasses of water and insert finger in back of throat repeatedly until fluid is clear. Do not induce emesis if the patient is unconscious or if a serious poison.

If inhaled: Remove victim to fresh air. Apply respiration if indicated.

If on skin: Remove contaminated clothing and other personal articles; wash thoroughly with soap and water.

If in eyes: Flush eyes thoroughly for at least 15 minutes with plenty of lukewarm water. Get medical aid immediately.

NOTE TO PHYSICIAN: Diazinon is a cholinesterase inhibitor. If a cholinergic crisis has developed, atropine should be used. If a cholinergic crisis has developed, atropine should be used. If a cholinergic crisis has developed, atropine should be used.

SEE STATE PAMPHLET FOR ALTERNATIVE PROCEDURES AND STATEMENTS



Manufactured By
Van Waters & Rogers
division of Univar

San Mateo, CA 94403

99 9 81

EPA Reg. No. 550-154-AA

EPA Est. No. 550-CA-1