

*Do not spray for 14 days	
5 days*	Strawberries, Figs
7 days*	Blueberries, Raspberries
10 days*	Apricots, Cherries, Grapes, Nectarines, Plums, Prunes
14 days*	Apples, Pears
21 days*	Grapefruit, Lemons, Oranges, Peaches

Thoroughly spray all plant parts. Generally, applications at 7 to 10 day intervals may be required. Do not feed treated vines to livestock.

Bean Pods	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Mites 4 ft. oz. in 3 gal. of water covers approximately 1,000 sq. ft. of garden
Cabbage	Aphids, Diamondback Moths, Flea Beetles, Imported
Cauliflower	Cabbage Worms, Dipterous Leaf Miners
Corianders	
Broccoli	4 ft. oz. in 3 gal. of water covers approximately 1,000 sq. ft. of garden
Kale, Radishes	
Turnips	
Carrots, Cherry	Aphids, Flea Beetles, Leafhoppers, Dipterous Leaf
Endive, Lettuce	Miners, Mites
Peking Spinach	4 ft. oz. in 3 gal. of water covers approximately 1,000 sq. ft. of garden
Cantaloupes	Aphids, Thrips, Mites
Cucumbers	4 ft. oz. in 3 gal. of water covers approximately 1,000 sq. ft. of garden
Muskmelon	
Squash	
Watermelon	
Peppers	Aphids, Colorado Potato Beetles, Ear Armyworms, Tomato
Potatoes	Flea Beetles, Leafhoppers, Southern Armyworms,
Tomatoe	tomato and tomato Vinegar Flies, <i>Drosophila</i> spp. (tomato)
	4 ft. oz. in 3 gal. of water covers approximately 1,000 sq. ft. of garden
Onions	Onion Thrips
	4 ft. oz. in 3 gal. of water covers approximately 1,000 sq. ft. of garden
Sweet Corn	Aphids, Flea Beetles, Grasshoppers, Mites
	4 ft. oz. in 3 gal. of water covers approximately 1,000 sq. ft. of garden

Cabbage	Cutworms (Surface Subterranean)
Corn	24 in sq per 1000 sq ft sufficient water to achieve through coverage
Letuce	
Pole Beans	
Radishes	Apply broadcast just before parting immediate
Red beets	work into soil 2-3 inches for Surface Cutworms 3-4 inches for Subterranean Cutworms
Snap Beans	
Tomatoes	
Turnips	

1 day - Vegetables with
 1 day - Fresh Tomatoes Sweet Corn
 3 day - Le. Laloups Mushrooms Squash Watermelons
 5 day - Broccoli Cauliflower
 7 day - Beans Cabbage Cucumbers Peppers
 10 day - Carrots Mary Culiards Endive Kale Lettuce
 10 day - Onions Parsnips Radishes Spinach Turnips
 36 day - Potatoes
 * OF HARVEST

ATC82

32 fl oz n3 gal of water

Thoroughly spray window and door screens, sills, foundations, porch and patio walls and ceilings, entrance ways, walks, garbage cans, fire trucks and into cracks and other places where insects can hide.

Do not treat plants or animals

*Spray a 5 ft band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet

*See table recommendations for other control measures for these pests.

To Aid in the control of fire ants in lawns and other recreation areas, apply 4 fl oz (18 g) per 1 m minimum of 1 gal of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal per mound. High pressure sprays may disturb the ants and cause migration reducing product effectiveness.

Note: For best results apply in cool weather 65-80°F in early morning or late evening hours. Treat new mounds as they appear.

ACCEPTED
20 JAN 1984
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended for the pesticide registered under EPA Reg. No. 655-476