

WEED KILLER

NET CONTENTS **No. 250**

20% SOLUTION

A Water-soluble Herbicide containing 2, 4-D

Active Ingredients:

• Mixed Ethanolamine Salts of 2, 4-Dichlorophenoxyacetic Acid	33.9%
Inert Ingredients:	66.1%
TOTAL	100.0%

• Equivalent to 2, 4-Dichlorophenoxyacetic Acid - 20.0%

KILLS

~~DANDELION - POISON IVY - POISON OAK - RAGWEED - PLANTAIN, (BROAD AND NARROW LEAF) - THISTLES - DOCKS - WILD MORNING-GLORY - SHEPHERD'S PURSE - WILD CARROT, NASTUR~~

and many other broad-leaved weeds without injury to grasses.

For lawns and ornamental turf in golf courses, parks, playgrounds and recreational areas, non-crop areas such as along highways, railroad rights-of-way, airfields and around farm buildings, and spot treatments in pastures.

CAUTION

KEEP OUT OF REACH OF CHILDREN

May cause skin irritation. Avoid contact with eyes, skin and clothing. Avoid spray drift to susceptible plants as this product may injure cotton, beans, ornamentals, etc. (Course sprays are less likely to drift.) Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. Do not store near fertilizers, seeds, insecticides or fungicides.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes.

Manufactured by

BAIRD & McGUIRE, INC.,
Holbrook, Mass.

USDA Reg.
#551-2




DIRECTIONS

Use at the rate of one gallon with water to make 250 gallons of spray, equivalent to eight teaspoons to 2 1/2 gallon water. Apply with hand or power sprayer in sufficient quantity to give thorough coverage of the weeds without excessive runoff or wetting of the soil. For very dense foliage approximately 2 1/2 gallons per acre (one gallon to 200 square feet) will be required. More luxuriant foliage will require more spray. If extremely efficient spray equipment is available, it may be possible to obtain thorough coverage with a smaller volume of spray mixture, in which case the concentration of Weed Killer in the spray should be proportionately increased.

The recommended concentration should be doubled for the difficult to control weeds, such as yellow and green foxtail. Double the concentration also for weeds growing in hard locations. Control of the species in these locations is doubtful. Deep-rooted perennial weeds such as bindweed, Canada thistle and hairy vetch are more difficult to kill and may require repeated applications.

Apply with pressures of 5 to 250 pounds per square inch where dense foliage occurs. Higher pressure may cause matting of vegetation and make thorough coverage impossible. Apply during warm weather when temperature is at least 60°F, and when weeds are growing actively. Apply to lawns between mowings so that a maximum of foliage is present.



WEED KILLER

NET CONTENTS No. 250

20% SOLUTION

A Water-soluble Herbicide containing 2, 4-D

Active Ingredient:

• Mixed Ethanolamine Salts of	
2, 4-Dichlorophenoxyacetic Acid	33.3%
Inert Ingredients:	66.7%
TOTAL	100.0%

* Equivalent to 2, 4-Dichlorophenoxyacetic Acid - 20.0%

KILLS

~~DANDELION - POISON IVY - POISON OAK - RAGWEED - PCNTAIN~~
~~- (BROAD AND NARROW LEAF) - THISTLES - DOCKS - WILD MORRIS~~
~~INC. GLORY - SHEPHERD PURSE - WILD CARROT, MALLOW~~

and many other broad leaved weeds without injury to grasses.

For lawns and ornamental turf in golf courses, parks, playgrounds and recreational areas, non-crop areas such as along highways, railroad rights of way, airfields and around farm buildings, and spot treatments in pastures.

CAUTION

KEEP OUT OF REACH OF CHILDREN

May cause skin irritation. Avoid contact with eyes, skin and clothing. Avoid spray drift to susceptible plants as this product may injure cotton, beans, ornamentals, etc. (Course sprays are less likely to drift.) Thoroughly clean spray equipment with a suitable chemical cleaner before using for other purposes. Do not store near fertilizers, seeds, insecticides or fungicides.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

Manufactured by
PAIRD & McGUIRE, INC.
Belbrook, Mass.

USDA Reg.
#551-2



IMPORTANT

Avoid application to vegetables, flowering plants, shrubbery, trees or other desirable type of plants. This is particularly necessary when treating weeds in orchards, vineyards and nurseries.

Do not use on bent grass lawns, except for spot spraying.

Always clean spraying equipment thoroughly before using for any other purpose. Rinse several times with hot water containing soap or trisodium phosphate (one pound in twenty-five gallons of water). Allowing the tank to stand filled with water overnight is recommended. Be sure to clean out hose and spray nozzle.

WARNING

Do not dispose of poison ivy by burning after it has been killed. Smoke from the burning vine is very poisonous on contact with the skin.

CAUTION

Do not graze treated areas by dairy animals within 7 days after treatment.

Do not use on newly seeded turf.

Do not contaminate irrigation ditches or water used for domestic purposes.

Do not apply to clover as injury may result.

60